# Transcriptome analysis offers a comprehensive illustration of the genetic background of pediatric acute myeloid leukemia

Norio Shiba,<sup>1-3</sup> Kenichi Yoshida,<sup>4</sup> Yusuke Hara,<sup>2,3,5</sup> Genki Yamato,<sup>2,3,5</sup> Yuichi Shiraishi,<sup>6</sup> Hidemasa Matsuo,<sup>7</sup> Yusuke Okuno,<sup>8</sup> Kenichi Chiba,<sup>6</sup> Hiroko Tanaka,<sup>9</sup> Taeko Kaburagi,<sup>2,5</sup> Masanobu Takeuchi,<sup>1</sup> Kentaro Ohki,<sup>10</sup> Masashi Sanada,<sup>3</sup> Jun Okubo,<sup>2,11</sup> Daisuke Tomizawa,<sup>12</sup> Tomohiko Taki,<sup>13</sup> Akira Shimada,<sup>14</sup> Manabu Sotomatsu,<sup>2</sup> Keizo Horibe,<sup>3</sup> Takashi Taga,<sup>15</sup> Souichi Adachi,<sup>7</sup> Akio Tawa,<sup>16</sup> Satoru Miyano,<sup>17</sup> Seishi Ogawa,<sup>4,18,19</sup> and Yasuhide Hayashi<sup>2,20</sup>

<sup>1</sup>Department of Pediatrics, Yokohama City University Hospital, Yokohama, Japan; <sup>2</sup>Department of Hematology/Oncology, Gunma Children's Medical Center, Shibukawa, Japan; <sup>3</sup>Clinical Research Center, National Hospital Organization Nagoya Medical Center, Nagoya, Japan; <sup>4</sup>Department of Pathology and Tumor Biology, Graduate School of Medicine, Kyoto University, Kyoto, Japan; <sup>5</sup>Department of Pediatrics, Gunma University Graduate School of Medicine, Maebashi, Japan; <sup>6</sup>Center for Cancer Genomics and Advanced Therapeutics, National Cancer Center, Tokyo, Japan; <sup>7</sup>Department of Human Health Sciences, Kyoto University Graduate School of Medicine, Kyoto, Japan; <sup>9</sup>Center for Advanced Medicine and Clinical Research, Nagoya University Hospital, Nagoya, Japan; <sup>9</sup>Laboratory of Sequence Analysis, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan; <sup>10</sup>Department of Pediatric Hematology and Oncology Research, National Research Institute for Child Health and Development, Tokyo, Japan; <sup>11</sup>Department of Pediatrics, Hokkaido University Graduate School of Medical Technology, Kyorin University Faculty of Health Sciences, Mitaka, Japan; <sup>14</sup>Department of Pediatrics, Okayama University Graduate School of Medicine, Okayama, Japan; <sup>15</sup>Department of Pediatrics, Shiga University of Medical Science, Ohtsu, Japan; <sup>16</sup>Department of Pediatrics, National Hospital Organization Osaka National Hospital, Osaka, Japan; <sup>17</sup>Laboratory of DNA Information, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan; <sup>18</sup>Institute for the Advanced Study of Human Biology (WPI-ASHBi), Kyoto University, Kyoto, Japan; <sup>19</sup>Department of Medicine, Centre for Haematology and Regenerative Medicine, Karolinska Institute, Stockholm, Sweden; and <sup>20</sup>Institute of Physiology and Medicine, Jobu University, Takasaki, Japan

#### **Key Points**

- Using RNA-seq in pediatric AML patients,
  5 gene rearrangements were newly identified, including NPM1 and RUNX1 gene rearrangements.
- RNA-seq unmasked the complexity of gene alterations in pediatric AML by identifying disease-causing alterations in nearly all patients.

Recent advances in the genetic understanding of acute myeloid leukemia (AML) have improved clinical outcomes in pediatric patients. However,  $\sim$ 40% of patients with pediatric AML relapse, resulting in a relatively low overall survival rate of  $\sim$ 70%. The objective of this study was to reveal the comprehensive genetic background of pediatric AML. We performed transcriptome analysis (RNA sequencing [RNA-seq]) in 139 of the 369 patients with de novo pediatric AML who were enrolled in the Japanese Pediatric Leukemia/ Lymphoma Study Group AML-05 trial and investigated correlations between genetic aberrations and clinical information. Using RNA-seq, we identified 54 in-frame gene fusions and 1 RUNX1 out-of-frame fusion in 53 of 139 patients. Moreover, we found at least 258 gene fusions in 369 patients (70%) through reverse transcription polymerase chain reaction and RNA-seq. Five gene rearrangements were newly identified, namely, NPM1-CCDC28A, TRIP12-NPM1, MLLT10-DNAJC1, TBL1XR1-RARB, and RUNX1-FNBP1. In addition, we found rare gene rearrangements, namely, MYB-GATA1, NPM1-MLF1, ETV6-NCOA2, ETV6-MECOM, ETV6-CTNNB1, RUNX1-PRDM16, RUNX1-CBFA2T2, and RUNX1-CBFA2T3. Among the remaining 111 patients, KMT2A-PTD, biallelic CEBPA, and NPM1 gene mutations were found in 11, 23, and 17 patients, respectively. These mutations were completely mutually exclusive with any gene fusions. RNA-seq unmasked the complexity of gene rearrangements and mutations in pediatric AML. We identified potentially disease-causing alterations in nearly all patients with AML, including novel gene fusions. Our results indicated that a subset of patients with pediatric AML represent a distinct entity that may be discriminated from their adult counterparts. Based on these results, risk stratification should be reconsidered.

Submitted 7 May 2019; accepted 9 July 2019. DOI 10.1182/ bloodadvances.2019000404. The full-text version of this article contains a data supplement. Downloaded from http://ashpublications.net/bloodadvances/article-pdf/3/20/3157/1328998/advancesadv2019000404.pdf by guest on 01 June 2024

#### Introduction

Recent advances in the genetic understanding of cancer have drastically improved clinical outcomes in pediatric patients. However, ~40% of patients with childhood acute myeloid leukemia (AML) relapse, resulting in a relatively low overall survival (OS) rate of  $\sim$ 70%. AML is caused by various chromosomal aberrations, gene mutations/epigenetic modifications, and deregulated/ overregulated gene expression, leading to increased proliferation and decreased differentiation of hematopoietic progenitor cells.<sup>1-4</sup> Several clinically important molecular markers have been discovered in patients with AML, which assisted in improving risk stratification. However, the evaluation of risk remains challenging even after incorporating these molecular markers in clinical practice.

Patients with biallelic CEBPA mutations, NPM1 mutations, RUNX1-RUNX1T1 fusions, or CBFB-MYH11 fusions have relatively good outcomes in response to treatment with chemotherapy-based consolidation regimens.<sup>5,6</sup> However, the prognosis of patients with FUS-ERG, NUP98-NSD1, CBFA2T3-GLIS2, NUP98-KDM5A, or KMT2A-MLLT3 with high MDS1 and EVI1 complex locus protein EVI1 (MECOM, synonyms EVI1 and PRDM3) expression was dismal, because many of these patients could not be during the first complete remission (CR), even with allogeneic transplantation.<sup>7-10</sup>

FMS-like tyrosine kinase 3-internal tandem duplication (FLT3-ITD) is a major alteration in pediatric and adult AML that is significantly associated with poor prognosis. However, the prognosis of patients with this alteration is not necessarily dismal, depending on concomitant genetic alterations.11-13 In our previous studies, the overexpression of PRD1-BF1-RIZ1 homologous domain containing 16 (PRDM16, also known as MEL1) was highly recurrent in patients with both pediatric AML and adult AML with intermediate- and highrisk cytogenetic profiles while being independently associated with an adverse outcome.14,15

Recently, Bolouri et al reported the seminal comprehensive molecular landscape and characteristics of pediatric AML in Children's Oncology Group AML trials.<sup>16</sup> However, the biologic basis differs among pediatric AML patients in the United States, Europe, and Japan. The objective of this study was to reveal the genetic alterations of all Japanese patients with pediatric AML and the differences in the genetic background between favorable and adverse groups using RNA sequencing (RNA-seq) by identifying new cytogenetic aberrations and comparing gene expression.

#### **Methods**

#### Study design and participants

In this study, we enrolled 485 patients with de novo AML who registered in the Japanese Pediatric Leukemia/Lymphoma Study Group (JPLSG) AML-05 trial between November 2006 and December 2010. The trial was registered with the University Hospital Medical Information Network Clinical Trials Registry (#UMIN000000511; http://www.umin.ac.jp/ctr/index.htm), and details of the schedules and treatment regimens have been described previously (supplemental Figure 2).<sup>17</sup> Patients diagnosed with acute promyelocytic leukemia or AML with Down syndrome were excluded from this study. Among the 485 patients with AML who were enrolled in the AML-05 trial, 116 were excluded because of sample insufficiency for diagnosis (n = 7), not meeting the criteria

(n = 6), misdiagnosis (n = 25), lack of RNA samples (n = 74), or other reasons (n = 4). Therefore, a total of 369 patients were genetically analyzed. In these 369 analyzed patients, the age and initial white blood cell count were higher, whereas the proportion of M7 patients was lower compared with the 74 nonanalyzed patients. However, there were no significant differences observed between these groups in terms of mortality (20% vs 24%, log-rank P = .84) (supplemental Table 1). The present study was conducted in accordance with the Declaration of Helsinki and approved by the institutional review boards of Gunma Children's Medical Center and the participating institutes and the ethical review board of the JPLSG AML-05 trial. Written informed consent was provided by all patients or their parents/guardians.

#### Sample preparation

All leukemic samples were obtained from the bone marrow or peripheral blood at the time of diagnosis. DNA and total RNA samples were prepared using the AllPrep DNA/RNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Complementary DNA was prepared using 0.8 to 1.0 µg of total RNA and Ready-To-Go reverse transcription polymerase chain reaction (RT-PCR) beads (GE Healthcare, Buckinghamshire, United Kingdom).

#### **RNA-seq**

We performed RNA-seq for 139 out of the 369 patients with pediatric AML in order to obtain a complete registry of gene rearrangements, other genetic lesions, and gene expressions in pediatric AML. The RNA-seg data were available at the European Genome-Phenome Archive (EGAS00001003701). The cvtogenetic characteristics of the analyzed patients are shown in supplemental Table 2. The study population mainly included patients with a normal karyotype (60/70 patients), FLT3-ITD (33/47 patients), KMT2A-PTD (12/13 patients), and high PRDM16 expression (65/ 84 patients). All 137 patients with core binding factor AML (CBF-AML) and many patients with KMT2A rearrangements were excluded from RNA-seq, because these fusions are already known to affect leukemogenesis. The main reasons behind the inability to perform analyses were assay failure, sample availability, and/or sample quality. The quality of the extracted RNA was assessed using TapeStation system (Agilent Technologies, Santa Clara, CA). Sequencing libraries were prepared using an NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA), and prepared libraries were run on a HiSeq 2500 high-throughput sequencing system. Sequencing reads were aligned using bowtie and blat, and fusion genes were analyzed using Genomon-fusion.<sup>18</sup> Candidate gene fusions were validated by RT-PCR. Obtained reads were also analyzed using an in-house pipeline, GenomonExpression (https://github.com/Genomon-Project/GenomonExpression), to obtain fragments per kilobase million (FPKM) values. We used Cluster 3.0 to perform hierarchical clustering.<sup>19</sup> Briefly, genes that were not expressed (FPKM = 0) in >20% of samples and genes with a standard deviation <10 were excluded. For the remaining genes, their FPKM values were log transformed. Finally, an average linkage was calculated. Results were visualized using Java TreeView.<sup>20</sup>

#### **Targeted gene mutation analysis**

We performed targeted gene mutation analysis of KIT (exons 8 and 17), CEBPA (all coding regions), NPM1 (exon 12), WT1 (exons 7-10), RUNX1 (exons 2-6), NRAS (exons 1 and 2), KRAS (exons 1 and 2), ASXL1 (exon 11), ASXL2 (exons 11 and 12), and GATA2

Downloaded from http://ashpublications.net/bloodadvances/article-pdf/3/20/3157/132898/advancesadv2019000404.pdf by guest on 01 June 2024

(exons 4-6). The allelic ratio (AR) of *FLT3*-ITD to wild-type alleles was further examined using GeneScan (Applied Biosystems, Foster City, CA), and a defined mutant/wild-type >0.40 was considered as high AR according to previous studies.<sup>12,21-26</sup> *KMT2A*-PTD was analyzed using previously described multiplex ligation-dependent probe amplification methods.<sup>27</sup> Gene fusions, namely, *RUNX1-RUNX1T1*, *CBFB-MYH11*, *KMT2A* rearrangements, *NUP98-NSD1*, *CBFA2T3-GLIS2*, *NUP98-KDM5A*, *FUS-ERG*, and *DEK-NUP214*, were also analyzed in all patients. In addition to these conventional gene fusions, we performed RT-PCR using samples from all patients to detect the *MYB-GATA1*, *NPM1-CCDC28A*, *MLF1-NPM1*, *NPM1-TRIP12*, *TBL1XR1-RARB*, *ETV6-CTNNB1*, *ETV6-NCOA2*, *ETV6-MECOM*, *RUNX1-FNBP1*, *RUNX1-CBFA2T2*, and *RUNX1-CBFA2T3* alterations identified through RNA-seq.

# Quantification of *PRDM16* and *MECOM* gene expression

Quantitative RT-PCR analysis of the *PRDM16* and *MECOM* genes was performed using the 7900HT Fast Real Time PCR System, TaqMan Gene Expression Master Mix, and TaqMan Gene Expression Assay (Applied Biosystems), which have been previously described.<sup>28</sup>

#### Data collection

A standardized form was used to record clinical variables, including patient demographic information. Every 6 months, data forms were forwarded to the JPLSG AML-05 trial data coordination center at the National Center for Child Health and Development (Tokyo, Japan), reviewed for internal consistency and face validity, and transferred into an Excel database (Microsoft Corporation, Redmond, WA). The clinical data of patients in each risk group were followed up until December 2013 (censored for 3 years from the date of final registration). The JPLSG conducted a central review of morphological classification and karyotyping based on the World Health Organization's classification, French-American-British (FAB) classification and cytogenetic analysis using conventional G-banding. OS was defined as the time from AML diagnosis to death or censorship at the last follow-up. Event-free survival (EFS) was defined as the time from AML diagnosis to treatment failure, relapse, death, or last follow-up.

#### Statistical analysis

Univariate and multivariate Cox regression analyses were calculated using EZR (version 1.37; Saitama Medical Center, Jichi Medical University, Saitama, Japan), which is a graphical user interface for R (version 3.4.1; The R Foundation, Vienna, Austria)<sup>29</sup> to extract the adverse and favorable risk factors related to OS. We defined the risk factors with a hazard ratio (HR) of <0.20 as favorable and those with a ratio of >2.0 as adverse. Continuous variables are presented as means  $\pm$  standard deviations and/or medians with ranges. Categorical variables are represented by frequencies and percentages. For all analyses, the P values were 2 tailed, and P < .05 was considered statistically significant. A multivariate Cox regression analysis was used to identify independent genetic factors related to OS. Initially, we included all genetic variables in the first model and then sequentially removed the nonsignificant variables ( $P \ge .05$ ). To test the probability of cooccurrence and mutual exclusivity of 2 parameters, we performed a Monte Carlo simulation using the total number of patients and the number of patients who were positive for each parameter.  $^{\rm 30}$ 

#### Results

#### **Overview of RNA-seq**

We performed RNA-seq in 139 patients with pediatric AML and identified a total of 54 in-frame fusions and 1 *RUNX1* out-of-frame gene fusion in 53 patients. Many of the recurrent gene fusions identified in this study have been previously reported as targets in AML (Figure 1). Among these 54 gene fusions, 5 gene rearrangements were newly identified, namely, *NPM1-CCDC28A*, *TRIP12-NPM1*, *MLLT10-DNAJC1*, *TBL1XR1-RARB*, and *RUNX1-FNBP1*. We also identified 17 rare gene rearrangements, including *MYB-GATA1*, *NPM1-MLF1*, *ETV6-NCOA2*, *ETV6-MECOM*, *ETV6-CTNNB1*, *RUNX1-PRDM16*, *RUNX1-CBFA2T2*, and *RUNX1-CBFA2T3* (Figures 1 and 2). The characteristics of the patients with these fusions are shown in Table 1. We identified 5 *RUNX1* rearrangements other than *RUNX1-RUNX1T1*, 4 *NPM1* rearrangements, 3 *ETV6* fusions, and 1 *TBL1XR1-RARB* fusion in FAB-M3 patients without the *PML-RARA* fusion.

# Complete genetic landscape of the JPLSG AML-05 trial

We showed the demographic characteristics and genetic landscape of 369 patients with AML enrolled in the JPLSG AML-05 trial (Figure 1). RUNX1-RUNX1T1 and KIT mutations were more frequently observed and CBFB-MYH11, trisomy 8, and NPM1 were less frequently observed as a molecular characteristic in the Japanese cohort compared with the Children's Oncology Group (supplemental Figure 1). We also showed the result of univariate and multivariate Cox regression analysis in Table 2. In the univariate analysis, factors significantly associated with OS included CBFB-MYH11 (HR, 0.11, 95% confidence interval [CI], 0.02-0.80), biallelic CEBPA (HR, 0.16; 95% CI, 0.02-1.13), RUNX1-RUNX1T1 (HR, 0.19; 95% Cl, 0.09-0.39), NPM1 (HR, 0.19; 95% Cl, 0.03-1.35), FLT3-ITD (HR, 2.92; 95% Cl, 1.83-4.65), CBFA2T3-GLIS2 (HR, 3.00; 95% Cl, 2.54-6.48), PRDM16 overexpression (HR, 3.09; 95% CI, 2.06-4.65), and NUP98-NSD1 (HR, 5.07; 95% CI, 2.54-10.1). The multivariate analysis revealed that CBFB-MYH11 (HR, 0.08; 95% Cl, 0.01-0.58), biallelic CEBPA (HR, 0.10; 95% CI, 0.01-0.69), RUNX1-RUNX1T1 (HR, 0.17; 95% CI, 0.08-0.35), NPM1 (HR, 0.09; 95% CI, 0.01-0.64), FLT3-ITD (HR, 2.53; 95% Cl, 1.57-4.08), FUS-ERG (HR, 4.54; 95% Cl, 1.81-11.4), and RPN1-MECOM (HR, 5.44; 95% Cl, 1.32-22.4) were independent prognostic factors associated with OS. Concerning CBF-AML, 9 of 137 patients died of primary disease or other causes (4 and 5 patients, respectively) (Table 3). Although CBF-AML patients with the KIT mutation exhibited a higher relapse rate than those without the KIT mutation, most of these patients were rescued by hematopoietic stem cell transplantation.<sup>31</sup> No fusion genes were identified in patients with the NPM1 or biallelic CEBPA mutation, and their prognosis was absolutely favorable irrespective of the presence of FLT3-ITD. Among the 41 patients with the biallelic CEBPA or NPM1 mutation, only 2 patients died, and the causes of death were acute respiratory disorder syndrome (ARDS) and primary disease (Table 3). On the other hand, the prognosis of patients with adverse gene fusions (eg, FUS-ERG, NUP98-NSD1, RPN1-MECOM, KMT2A-AFDN, DEK-NUP214, CBFA2T3-GLIS2, and NUP98-KDM5A) and adverse mutations (eg, FLT3-ITD, RUNX1, and



Figure 1. Clinical and genetic profiles of the 369 patients enrolled in the JPLSG AML-05 trial. (A) Pie chart represented the frequency of each gene alterations. (B) Landscape of 369 de novo pediatric AML patients. Each column indicates 1 patient. \*\* Both gene fusions were observed in the same patient. *RUNX1*-R, *RUNX1* rearrangement; *NPM1*-R, *NPM1*-rearrangement.

*KMT2A*-PTD) was dismal (5-year OS, 31%; EFS, 9%). The characteristics of patients with unfavorable gene fusions are shown in supplemental Table 3. Although all of patients with *FUS-ERG*, *RPN1-MECOM*, *KMT2A-AFDN*, and *DEK-NUP214* were predicted by chromosomal G-banding, *NUP98-NSD1*, *NUP98-KDM5A*, and *CBFA2T3-GLIS2* were cryptic. No disease-causing genetic alterations were identified in 45 patients out of 369 patients with AML (29 patients with FAB non-M7 and 16 patients with FAB-M7) (supplemental Table 4). Ten patients were cytogenetically normal, whereas the remaining 35 patients had structural alterations in chromosomes, including complex karyotype (n = 23), monosomy 7 (n = 4), and trisomy 8 (n = 3). The prognosis of these patients tended to be dismal (3-year OS, 67%; 3-year EFS, 42%); thus, they should be treated as high-risk patients. Further research is warranted to confirm these observations.

## Characteristics of patients with pediatric AML with normal karyotype

Sixty out of 70 patients with a normal karyotype were analyzed using RNA-seq. Although some cryptic gene fusions were identified (eg, *NUP98-NSD1* and *CBFA2T3-GLIS2*), fewer gene fusions were identified in this subset of patients. In particular, gene fusions were not identified in patients with normal karyotype with biallelic *CEBPA*, *NPM1*, and *KMT2A*-PTD aberrations, which are frequently observed in the normal karyotype (supplemental Figure 3).

## Characteristics of patients with pediatric AML with *KMT2A* rearrangements

Sixty-one *KMT2A* rearrangements were identified in 369 patients with AML (16%). Among them, 7 *KMT2A* fusions not identified through conventional chromosomal G-banding were detected through RNA-seq. *KMT2A-MLLT3* was the most commonly observed alteration (30/61 patients). Intriguingly, the prognosis of these patients was highly dependent on the expression of the *MECOM* gene. Seven of 11 patients with high *MECOM* expression died compared with only 1 of 19 patients with low *MECOM* 

expression (P < .001). *KMT2A-ELL* and *KMT2A-MLLT1* were the second most commonly observed alterations in patients with pediatric AML (supplemental Table 5).

### Characteristics of patients with pediatric AML with FAB-M7

In this study, we performed RNA-seq in only 9 of 34 patients with acute megakaryoblastic leukemia. Among them, we confirmed 3 types of recurrent gene fusions identified through RT-PCR, including 2 *RBM15-MKL1*, 3 *CBFA2T3-GLIS2*, and 1 *NUP98-KDM5A*. In the remaining 3 patients with acute megakaryoblastic leukemia, there were no gene fusions identified. Among 34 patients registered and treated in the JPLSG AML-05 trial, 19 of 34 patients showed high expression of the *MECOM* gene. Although high *MECOM* gene expression was associated with poor prognosis in other FAB patients, this association was not observed in patients with FAB-M7 (Figure 1). Interestingly, patients with the *CBFA2T3-GLIS2* and *NUP98-KDM5A* alterations, which were associated with poor prognosis, showed low *MECOM* gene expression, indicating the presence of a different pathogenesis mechanism.

#### Gene expression profiling

The RNA-seq expression profiles of 139 patients were subjected to all-gene-based unsupervised hierarchical clustering, and 3 clusters were generated (blue, orange, and green) (Figure 3A). Thirteen out of 14 patients with *KMT2A* rearrangement were classified into the blue cluster, and all patients with *FLT3*-ITD with high AR were classified into the orange or green cluster. On the contrary, 6 of 7 patients with *NUP98-NSD1* and 2 patients with *DEK-NUP214* were classified into the green cluster. The gene expression pattern of *NUP98-KDM5A* was different from the other *NUP* fusions in this study (Figure 3A). Furthermore, *HOXA7*, *HOXB3*, *HOXB6*, *HOXB7*, *RUNX1*, *FLT3*, *PRDM16*, *HIST1H1C*, and *HIST1H2BD* were overexpressed in patients with *FLT3*-ITD (Figure 3B-C), and *HOX* genes were underexpressed in patients with biallelic *CEBPA*. Intriguingly, *FLT3* and *HOXAs* were

Figure 2. Novel/rare fusion genes identified in patients with de novo pediatric AML. (A-D) The structure of the detected fusion proteins. (A) Five RUNX1 rearrangements. (B) Three ETV6 rearrangements. (C) Three NPM1 rearrangements. (D) Other novel gene fusions. R1, R2, and R3 are 51  $\pm$  52 amino acid tandem repeats that comprise the DNA-binding domain Ext-leukemiaassociated protein (LAP). 14-3-3 BD, 14-3-3 binding domain; AD, acidic domains; AT-hook, adenine-thymine hook; CCD, coiled-coil domain; CDC15, CDC15 homology region; CID, CBP interaction domain; CtBP, C-terminal binding protein; ETS, ETS domain; FCHD, FER-CIP1 homology domain; HLH, helix-loop-helix; ID, transcription inhibition domain; LiSH, Lis homology domain; MBD, metal-binding domain; MBR, moderately basic region; MTG16, myeloid translocation gene 16; NES, nuclear export signal; NHR2, nervy homology region 2; NLS, nuclear localization signal; PRBR, putative ρ-binding region; RD, runt domain; RHD, runt homolog domain; SH3D, Src homology 3 domain; TAD, transcription activation domain; TAFH, TAF homology; ZF, zinc finger; ZFD, zinc-finger domain; Znf-MYND, MYND zinc-finger domain.



RNA no.	Novel/rare	Fusion gene	Sex	Age, y	WBC, 10 <sup>9</sup> /L	FAB	Risk	Chromosome	<i>FLT</i> 3-ITD	SCT	Relapse	Outcome
273	Rare	RUNX1-CBFA2T2	Σ	1.5	43.3	M1	Int	47,XY,+8,t(14;20)(q11.2;q11.2),del(21)(q21q22)[20]	I	T	I	Alive
227	Rare	RUNX1-CBFA2T3	ш	4.4	5.0	M1	Int	46,XX,t(16;21)(q24;q22)[4]/46,s ,t(1;16)(q32;p13.3)[16]	I	I	I	Alive
308	Rare	RUNX1-CBFA2T3	ш	1.6	35.4	M4	Ħ	46,XX,add(2)(q31),t(16;21)(q24;q22)[5]/45,sl,+2,-add(2), -11,add(18)(p11.2)[12//91,sl,×2,-11[2]	I	T	I	Alive
177	Rare	RUNX1-PRDM16	Σ	2.9	25.2	RAEB-T	Non-CR	45,XY,-7[1]/45, sl, t(11;21)(q13;q22)[19]	I	+	I	Dead
440	Novel	RUNX1-FNBP1	ш	0.2	10.0	M1	ц	46,XX,add(7)(q11.2),add(8)(q34),add(12)(p13)[20]	I	I	I	Alive
104	Rare	KAT6A-CREBBP	ш	17	10.6	M4	Int	46,XX,t(8;16)(p11.2:p13.3)[19]/46,XX[1]	I	I	I	Alive
216	Rare	KAT6A-CREBBP	Σ	15.3	18.1	M5a	ц	46,XY,t(8;16)(p11.2;p13.3)[20]	I	I	+	Dead
271	Rare	KAT6A-CREBBP	Σ	1.8	9.8	M5b	Int	46,XY,add(7)(q22),t(8;16)(p11.2;p13.3)[17]/46,XY[3]	I	I	I	Alive
331	Rare	ETV6-CTNNB1	Σ	4.8	2.2	M5a	Ţ	46,XY, add(3)(p21),inv(9)(p12q13),add(12)(p11.2)[7]/ 46,idem.del(20)(q11.2q13.3),add(22)(q11.2)[11]/ 46,XY,inv(9)(p12q13)[2]	I	+	+	Dead
45	Rare	ETV6-MECOM	Σ	4.8	33.9	M5a	Non-CR	45,XY,-7[1]/45,sl,t(3;12)(q26;p13)[18]/46,XY[1]	I	+	+	Dead
333	Rare	ETV6-NCOA2	ш	9.8	4.1	MO	ц	46,XX,t(8;12)(q11.2;p11.2)[20]	I	I	I	Alive
73	Rare	MYB-GATA1	Σ	0.3	20.2	M5a	Щ	47,Y,del(X)(p11.2),-6,+8,+r[15]/46,Y,del(X)(p11.2),-6, add(19)(p13.1),+r[2]/46, XY[3]	I	I	I	Alive
171	Novel	NPM1-CCDC28A TRIP12-NPM1	ш	1.4	16.6	M2	High	46,XX,add(2)(q31),add(5)(q35),add(6)(q15)[13]/46,XX[7]	+	+	I	Dead
58	Rare	NPM 1-MLF1	Σ	12.9	14.4	RAEB-T	Int	46,XY,t(3;5)(q24;q33)[20]	I	+	I	Alive
274	Rare	PICALM-MLLT10	ш	8.2	0.9	M1	<u>I</u>	46,XX,t(10;11)(p12;q14)[19]/46,XX[1]	I	+	+	Dead
30	Rare/novel	PICALM-MLLT10 MLLT10-DNAJC1	ш	4.1	26.5	M5a	Int	46,XX,add(9)(q27),add(8)(p11.2),t(10;11)(p12;q14), -16,add(19)(p13),+r1[13]/46,XX[7]	I	I	I	Alive
436	Rare	PICALM-MLLT10	Σ	10.2	29.9	M5a	Ţ	46,XY, add(1)[p,36,1), add(5)(q,31), add(7)(q,22), del(9)(q), r(10;11) (p1 2;q14), add(17)(p11.2)[1]/46,s1, add(3)(q11.2), -9, + mar1 [4]/ 46, sdl1, -4, -15, + der(?)t(?;-4)(?;q12), + mar2[11]/46, XY [4]	I	+	+	Alive
161	Novel	TBL 1XR1-RARB	ш	0.9	30.5	M3	Int	47,XX,+6[2]/46,XX	I	+	+	Alive
ນ	Rare	RBM 15-MKL 1	Σ	0.8	12.0	M7	Ţ	61,XXX,der(1)t(1;22)(p13;q13),t(1;22)(p13;q13),-3;-4,-5,+7, -9,-11,-12,-13,-15,-18,+19,-22[10]/46,XX[10]	I	+	+	Dead
162	Rare	RBM15-MKL1	ш	0.1	42.2	M7	Ъ	46,XY,der(1)t(1;22)(p13;q13)add(1)(q32),der(22) t(1;22)(add(1)(p22)(15)/46,XY[5]	I	I	I	Alive
					:							

Table 1. The clinical features of de novo pediatric AML patients with novel/rare fusions

F, female; Int, intermediate; M, male; SCT, stem cell transplantation; WBC, white blood cell count.

3162 SHIBA et al

				Univaria	te analysis			Multivari	ate analysis	
	No. of deat	hs/patients			95%	% CI			95%	6 CI
os	Negative	Positive	HR	P	Lower	Upper	HR	P	Lower	Uppe
CBFB-MYH11	93/338	1/31	0.11	.03	0.02	0.80	0.08	.01	0.01	0.58
Biallelic CEBPA	93/346	1/23	0.16	.07	0.02	1.13	0.10	.02	0.01	0.69
RUNX1-RUNX1T1	86/263	8/106	0.19	<.001	0.09	0.39	0.17	<.001	0.08	0.35
NPM1	93/351	1/18	0.19	.10	0.03	1.35	0.09	.02	0.01	0.64
KIT	80/283	14/86	0.43	.02	0.21	0.85				
KMT2A rearrangement	75/308	19/61	1.10	.73	0.64	1.88				
PICALM-MLLT10	93/366	1/3	1.22	.84	0.17	8.78				
WT1	86/345	8/24	1.33	.44	0.64	2.74				
KAT6A-CREBBP	93/366	1/3	1.40	.74	0.19	10.1				
RBM15-MKL1	93/367	1/2	1.76	0.57	0.24	12.7				
MECOM overexpression	71/312	23/57	1.91	.007	1.20	3.07				
<i>KMT2A</i> -PTD	89/356	5/13	2.28	.04	1.06	4.94				
FLT3-ITD	70/322	24/47	2.92	<.001	1.83	4.65	2.53	<.001	1.57	4.08
CBFA2T3-GLIS2	87/358	7/11	3.00	.005	2.54	6.48				
PRDM16 overexpression	52/285	42/84	3.09	<.001	2.06	4.65				
KMT2A-AFDN	91/365	3/4	3.15	.051	1.00	9.97				
NUP98-KDM5A	91/363	3/6	3.20	.049	1.01	10.2				
RUNX1	89/361	5/8	3.80	.004	1.54	9.38				
DEK-NUP214	91/365	3/4	4.91	.007	1.39	15.6				
NUP98-NSD1	85/358	9/11	5.07	<.01	2.54	10.1				
FUS-ERG	89/364	5/5	6.80	<.001	2.75	16.8	4.54	<.001	1.81	11.4
RPN1-MECOM	92/367	2/2	7.84	.04	1.92	32.0	5.44	.02	1.32	22.4

#### Table 2. Univariate and multivariate Cox regression analysis of OS in AML-05 trial

GVHD, graft-versus-host disease; TMA, thrombotic microangiopathy.

overexpressed in patients with *KMT2A*-rearrangements. Herein, we showed the detail of gene expression pattern focused on *HOXAs*, *HOXBs*, *PRDM16*, *FLT3*, *RUNX1*, and *HIST1s* in Figure 3B.

#### Discussion

Using RNA-seq, we identified 54 in-frame gene fusions and 1 out-offrame fusion in 53 of 139 patients with pediatric AML. Moreover, we found  $\geq$ 258 gene fusions in 369 patients (70%) through RT-PCR and RNA-seq. Only a few fusion genes were detected in patients with AML with a normal karyotype. Only 2 patients had 2 in-frame gene fusions, which resulted from 3-way translocations (*PICALM-MLLT10* and *MLLT10-DNAJC1*; *NPM1-CCDC28A* and *TRIP12-NPM1*). This finding suggested that most of the in-frame gene fusions are able to facilitate AML. In the remaining 111 patients, *KMT2A*-PTD, biallelic *CEBPA*, and *NPM1* mutations were found in 11, 23, and 17 patients, respectively. These mutations were completely mutually exclusive with any gene fusions. The *FLT3*-ITD alteration was more frequently observed in patients with adverse risk factors than in those with favorable risk factors, such as co-occurring mutations, and was considered to amplify the malignant potential.

In this study, we identified 5 novel gene rearrangements, namely, NPM1-CCDC28A, TRIP12-NPM1, NPM1-DNAJC1, TBL1XR1-RARB, and RUNX1-FNBP1, and 17 rare fusions. Concerning patients with NPM1 rearrangements, TRIP12 encodes E3 ligase, which ubiquitinates CDKN2A. NPM1 is a major component of the nucleolus, with important functions in regulating cell growth, proliferation, and transformation. Intriguingly, as an additional important effect, NPM1 stabilizes the protein levels of CDKN2A.32 CDKN2A is known to suppress aberrant cell growth by activating the p53 response.<sup>33</sup> The function of CDKN2A may be inhibited by fusing NPM1 and TRIP12. NPM1-MLF1 has been associated with therapy-related AML and myelodysplastic syndrome with good prognosis.<sup>34,35</sup> Our patient was diagnosed as RAEB-T and maintained CR for >3 years (Table 1). Regarding the roles of NPM1-MLF1 and NPM1-CCDC28A, the fusion resulted in the loss of DNA/RNA-binding domains. This may deregulate the function of NPM1 and facilitate leukemogenesis because the mutational hotspot is located within the DNA/RNA binding domain at exon 12. Interestingly, CCDC28A has been reported as the partner gene of NUP98 in T-ALL.36 Regarding the RUNX1-CBFA2T2 gene fusion, the expression of the CBFA2T2 gene was high in AML cells following the activation of the RUNX1 promoter compared with that observed in normal bone marrow. Thus, the RUNX1-CBFA2T2 gene fusion may be associated with leukemogenesis.<sup>37</sup> In addition, RUNX1-CBFA2T3 showed a gene expression pattern similar to that of RUNX1-RUNX1T1.38 Moreover, we identified novel out-offusion of RUNX1-FNBP1 (Figure 2). The affected RUNX1 is truncated at the end of the runt homolog domain, forming a stop codon within the FNBP1 gene, 88 to 90 bp downstream of the

able 3.	ne cause or dearr	n or pequatri	с раце	ents witr	n ravorable al	terations				
RNA no.	Alterations	Mutations	Sex	Age, y	WBC, 10 <sup>9</sup> /L	FAB	Chromosome	Risks	Relapse	Cause of death
13	RUNX1-RUNX1T1	KIT	Σ	15.7	29.2	M2	46,XY,t(8;21)(q22;q22) [20]	Low	+	Acute GVHD grade
31	RUNX1-RUNX1T1	I	Σ	17.9	6.9	M2	45,X,-Y,t(8;21)(q22;q22)[18]/46,XY[2]	Low	+	Dyspnea due to GVHD and TMA
134	RUNX1-RUNX1T1	KIT	ш	3.6	11.7	M2	46,XX,t(8;21)(q22;q22)[12]/46,XX[8]	Low	+	Infection, cardiac failure
146	RUNX1-RUNX1T1	I	Σ	5.4	9.8	RAEB-T	45,X,-Y,t(8;21)(q22;q22)[20]	Low	+	Primary disease
172	RUNX1-RUNX1T1	Ι	Σ	7.8	9.3	RAEB-T	45,X,-Y,t(8;21)(q22;q22)[19]/46,XY[1]	Low	+	Primary disease
200	RUNX1-RUNX1T1	КIT	ш	6.1	18.7	M2	46,XX,t(8;14;21)(q22;q24;q22)[20]	Low	+	Primary disease
355	RUNX1-RUNX1T1		Σ	14.5	19.2	RAEB-T	46,XY,t(8;18;21)(q22;q21;q22)[12]/46,XY[2]	Low	+	Sepsis
383	RUNX1-RUNX1T1	KIT, RUNX1	Σ	9.9	38.7	M2	46,XY,t(8;19)(q22;p13) or t(8;19;21)(q22;p13;q22)[20]	Non-CR	I	Primary disease
209	CBFB-MYH11	FL 73-ITD	Σ	16.2	172.2	M4Eo	$\begin{array}{l} 48,XY,+6,+8,inv(16)(p13q22)[8]/49,sl,+21[5]/50,sl1,+10[5]/\\ 49,sl2,+8[2]/49,sl3,+11[2] \end{array}$	Induction death	I	Intracranial hemorrhage (day 16)
26	biallelic CEBPA	Ι	Σ	6.8	10.7	M	46,XY[20]	Intermediate	I	ARDS
97	NPM1	I	ш	11.6	38.2	RAEB-T	47,XX,+18[1]/46,XX[19]	Intermediate	+	Primary disease

break point (Figure 2A). In this study, patients with RUNX1-CBFA2T3, RUNX1-CBFA2T2, and RUNX1-FNBP1 maintained CR without relapse for >3 years. On the other hand, the patient with RUNX1-PRDM16 did not achieve CR and eventually expired owing to primary disease following hematopoietic stem cell transplantation in non-CR. This patient showed the highest PRDM16 gene expression among the 369 patients with AML, whereas other patients with RUNX1 fusions showed low PRDM16 expressions. This finding indicates that the RUNX1-PRDM16 fusion gene has a function distinct from those of other RUNX1 fusions (eg, RUNX1-RUNX1T1) in terms of pathogenesis. RUNX1-PRDM16 has already been reported in myelodysplastic syndrome, de novo AML, therapyrelated AML, and chronic myelogenous leukemia and is associated with high PRDM16 gene expression and dismal outcomes.<sup>39-43</sup> We showed the genetic landscape of patients with AML with high levels of PRDM16 expression (supplemental Figure 4). Many adverse risk factors were observed in patients with high PRDM16 gene expression, and numerous patients died, indicating that high PRDM16 expression is associated with high-risk factors and dismal outcomes.4

Previous studies suggested *MYB-GATA1* as the cause of acute basophilic leukemia.<sup>44</sup> This fusion gene commits myeloid cells to the granulocyte lineage, blocking their differentiation.<sup>44</sup> This fusion gene has been reported to be a cause of infant AML FAB-M5 as a result of reducing the expression of the *GATA1* gene.<sup>45</sup> Recently, this fusion was also reported in a pediatric patient with acute erythroid leukemia.<sup>46</sup> The break point was quite the same as in our patient, and the characteristics were very similar, including age, FAB-M5, and favorable outcomes. Thus, the phenotype may depend on the co-occurrence of genomic alterations.

It has been reported that ETV6-NCOA2 in acute leukemia is associated with the coexpression of T-lymphoid and myeloid markers. Our patient was diagnosed with peroxidase-negative acute leukemia including minimally differentiated FAB-MO. As blast cells were positive for CD7, CD13, CD33, CD34, CD56, CD117, and HLA-DR by surface marker analysis, myeloid-natural killer cell acute leukemia should be considered. Consistent with the results of previous studies, this finding suggests that this fusion gene may be specific for a common myeloid-lymphoid progenitor cell.47,48 Intriguingly, similar to ETV6-NCOA2, ETV6-CTNNB1 has been previously reported in patients with T-cell acute lymphoblastic leukemia.<sup>49</sup> ETV6-MECOM was previously reported in patients with chronic myelogenous leukemia or therapy-related leukemia.50 Our patient with this fusion showed the eighth highest MECOM expression among the 369 patients. ETV6 may promote the expression of the MECOM gene and turn on the critical role in progression to the blast crisis of AML. Interestingly, all 3 patients with MECOM rearrangements and 1 patient with PRDM16 rearrangement harbored monosomy 7, suggesting that PRDM alterations may be associated with monosomy 7, and showed dismal outcomes (supplemental Table 3).

Of the 61 patients (16%) with *KMT2A* rearrangements, 16 patients did not show 11q23 breakage through conventional chromosomal G-banding. This suggested that fluorescence in situ hybridization analysis of *KMT2A* may be useful for the detection of rare *KMT2A* rearrangements, because >100 partner genes are already known (supplemental Table 5). *KMT2A-MLLT3* was the most common *KMT2A* rearrangement. Moreover, *MECOM* gene expression was





Downloaded from http://ashpublications.net/bloodadvances/article-pdf/3/20/3157/1328998/advancesadv2019000404.pdf by guest on 01 June 2024



Figure 3. (Continued).

useful in stratifying the prognosis of patients with KMT2A-MLLT3 (low expression [3-year OS, 92%; 3-year EFS, 86%] vs high expression [3-year OS, 39%; 3-year EFS, 9%]). However, the expression of the MECOM gene was not correlated with poor prognosis in patients with FAB-M7. One of the reasons for this may be that MECOM expression is also associated with the differentiation of megakaryocytes, despite the association of high MECOM expression with stem cell maintenance through the expression of the GATA2 gene in other FAB subtypes.<sup>51</sup> KMT2A-MLLT10, ELL, and MLLT1 were common partner genes found in 9, 7, and 5 patients with de novo pediatric AML, respectively (supplemental Table 5). Of the 7 patients with KMT2A-ELL, 3 patients expired. Intriguingly, these 3 patients were <1 year old, and 2 of them died due to ARDS. On the other hand, the prognosis of patients with KMT2A-MLLT10 tended to be worse than that of patients with other KMT2A gene fusions (ie, 3 of 9 patients died due to primary disease) (supplemental Table 5). Consistent with the findings of previous studies,<sup>52</sup> the prognosis of patients with KMT2A-AFDN was extremely dismal, indicating a different pathogenesis from other KMT2A rearrangements.

The co-occurrence and mutual exclusivity test identified that various clinical biomarkers may reflect different aspects of the same poor

prognostic subgroup of patients with AML (Figure 4). *RUNX1-RUNX1T1* and *CBFB-MYH11* formed a unique entity that was mutually exclusive of any fusions and mutations other than *KIT* mutation. High *MECOM* expression was significantly associated with *KMT2A*-R and monosomy 7, whereas high *PRDM16* gene expression was significantly associated with *FLT3*-ITD, *KMT2A*-PTD, and trisomy 8. These could potentially be used as surrogate biomarkers for this distinct subgroup, which shares several molecular signatures and predict the prognosis.

In view of the gene expression results, *HOX* genes were upregulated in patients with *FLT3*-ITD or *KMT2A*-rearrangements. Moreover, overexpression of some *HOX* is known to enhance the self-renewal of hematopoietic stem and progenitor cells and perturb differentiation.<sup>53</sup> Additionally, *FLT3*, *RUNX1*, and some *HIST1*s were upregulated in these patients, suggesting that the aberrant expression of these genes plays a crucial role in leukemogenesis and indicates poor prognosis. Tiberi et al reported that the expression of *HIST1* was significantly higher in patients with AML with low levels of H3K27me3 than in those with high levels of H3K27me3.<sup>54</sup> Repressing H3K27me3 levels affects the transcriptional repression in patients with AML harboring aberrant expression of *HOX*, *FLT3*, *RUNX1*, and *HIST1*.



Figure 4. Correlations between established risk factors for pediatric AML. The diagram indicates co-occurrence or mutual exclusivity between pairs of clinical features, genetic alterations, gene expressions, and outcomes. The circles indicate that crossed factors are statistically co-occurrent (yellow) or exclusive (green).

In the present study, it was shown that RNA-seq unmasked the complexity of gene alterations in pediatric AML. We identified potentially disease-causing alterations in nearly all patients with AML, including novel gene fusions. Our results indicated that a subset of patients with pediatric AML represent a distinct entity that may be discriminated from their adult counterparts through an investigation of the spectrum of gene rearrangements and mutations. Furthermore, we found that a complex interplay of genetic events contributes to the pathogenesis of AML in individual patients. These findings suggest that gene rearrangements in conjunction with mutations play essential roles in pediatric AML. Based on the results of the RNA-seq, the prognostic factors currently used for risk stratification in pediatric AML may be reconsidered, particularly for intermediate/high-risk patients.

#### Acknowledgments

The authors thank Yuki Hoshino for her valuable technical assistance. The authors would like to thank Enago (www.enago.jp) for the English language review. Supercomputing resources were provided by the Human Genome Center, Institute of Medical Science, The University of Tokyo.

This work was supported by a Grant-in-Aid for Scientific Research on Innovative Areas from the Ministry of Health, Labor and Welfare of Japan (15H05909) (S.O.), a grant for project for development of innovative research on cancer therapeutics (P-DIRECT) from the Japan Agency for Medical Research and Development (AMED; JP18ck0106250h0002) (S.O.), the Practical Research for Innovative Cancer Control from AMED (15ck0106066h0002) (M.S.), a grant for project for cancer research and therapeutics evolution (P-CREATE) from AMED (JP18cm0106501h0003) (S.O.), the Japan Society for the Promotion of Science (KAKENHI grants 25893028 [N.S.], 15H05909 [S.O.], 16K20951 [N.S.], and 17K10130 [N.S.]), a research grant from the Japanese Society of Hematology, and the Kawano Memorial Public Interest Incorporated Foundation for Promotion of Pediatrics.

#### Authorship

Contribution: N.S. and Y. Hayashi designed the study; N.S., K.Y., Y. Hara, G.Y., T.K., H.M., K.O., and A.S. performed the experiments; Y. Hayashi and S.O supervised the work; N.S., K.Y., Y. Hara, G.Y., Y.S., T.K., M.T., Y.O., and K.O. analyzed the results; N.S. constructed the figures; N.S., K.Y., and Y. Hayashi wrote the paper; and all authors critically reviewed the draft and approved the final version for publication.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: G.Y., 0000-0003-4000-0753; Y.O., 0000-0003-3139-9272; H.T., 0000-0001-9634-8922; D.T., 0000-0003-1520-7007; K.H., 0000-0002-6251-6059; A.T., 0000-0003-4520-7905; S.O., 0000-0002-7778-5374.

Correspondence: Yasuhide Hayashi, Institute of Physiology and Medicine, Jobu University, 270-1, Shinmachi, Takasaki 370-1393, Gunma, Japan; e-mail: hayashiy-tky@umin.ac.jp.

#### References

- 1. Fröhling S, Scholl C, Gilliland DG, Levine RL. Genetics of myeloid malignancies: pathogenetic and clinical implications. J Clin Oncol. 2005;23(26): 6285-6295.
- Marcucci G, Haferlach T, Döhner H. Molecular genetics of adult acute myeloid leukemia: prognostic and therapeutic implications. J Clin Oncol. 2011; 29(5):475-486.
- 3. Pui CH, Carroll WL, Meshinchi S, Arceci RJ. Biology, risk stratification, and therapy of pediatric acute leukemias: an update. J Clin Oncol. 2011;29(5): 551-565.
- 4. Patel JP, Gönen M, Figueroa ME, et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. N Engl J Med. 2012;366(12): 1079-1089.
- 5. Preudhomme C, Sagot C, Boissel N, et al; ALFA Group. Favorable prognostic significance of CEBPA mutations in patients with de novo acute myeloid leukemia: a study from the Acute Leukemia French Association (ALFA). *Blood*. 2002;100(8):2717-2723.
- Hollink IH, Zwaan CM, Zimmermann M, et al. Favorable prognostic impact of NPM1 gene mutations in childhood acute myeloid leukemia, with emphasis on cytogenetically normal AML. *Leukemia*. 2009;23(2):262-270.
- 7. Kong XT, Ida K, Ichikawa H, et al. Consistent detection of TLS/FUS-ERG chimeric transcripts in acute myeloid leukemia with t(16;21)(p11;q22) and identification of a novel transcript. *Blood*. 1997;90(3):1192-1199.
- de Rooij JD, Hollink IH, Arentsen-Peters ST, et al. NUP98/JARID1A is a novel recurrent abnormality in pediatric acute megakaryoblastic leukemia with a distinct HOX gene expression pattern. Leukemia. 2013;27(12):2280-2288.
- Hollink IH, van den Heuvel-Eibrink MM, Arentsen-Peters ST, et al. NUP98/NSD1 characterizes a novel poor prognostic group in acute myeloid leukemia with a distinct HOX gene expression pattern. *Blood.* 2011;118(13):3645-3656.
- 10. Matsuo H, Kajihara M, Tomizawa D, et al. EVI1 overexpression is a poor prognostic factor in pediatric patients with mixed lineage leukemia-AF9 rearranged acute myeloid leukemia. *Haematologica*. 2014;99(11):e225-e227.
- 11. Shimada A, Taki T, Tabuchi K, et al. Tandem duplications of MLL and FLT3 are correlated with poor prognoses in pediatric acute myeloid leukemia: a study of the Japanese childhood AML Cooperative Study Group. *Pediatr Blood Cancer*. 2008;50(2):264-269.
- 12. Taketani T, Taki T, Sugita K, et al. FLT3 mutations in the activation loop of tyrosine kinase domain are frequently found in infant ALL with MLL rearrangements and pediatric ALL with hyperdiploidy. *Blood*. 2004;103(3):1085-1088.
- Pratcorona M, Brunet S, Nomdedéu J, et al; Grupo Cooperativo Para el Estudio y Tratamiento de las Leucemias Agudas Mieloblásticas. Favorable outcome of patients with acute myeloid leukemia harboring a low-allelic burden FLT3-ITD mutation and concomitant NPM1 mutation: relevance to postremission therapy. *Blood.* 2013;121(14):2734-2738.
- Shiba N, Ohki K, Kobayashi T, et al. High PRDM16 expression identifies a prognostic subgroup of pediatric acute myeloid leukaemia correlated to FLT3-ITD, KMT2A-PTD, and NUP98-NSD1: the results of the Japanese Paediatric Leukaemia/Lymphoma Study Group AML-05 trial. Br J Haematol. 2016; 172(4):581-591.
- 15. Yamato G, Yamaguchi H, Handa H, et al. Clinical features and prognostic impact of PRDM16 expression in adult acute myeloid leukemia. *Genes Chromosomes Cancer*. 2017;56(11):800-809.
- Bolouri H, Farrar JE, Triche T Jr, et al. The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and agespecific mutational interactions [published correction appears in Nat Med. 2018;24(4):526]. Nat Med. 2018;24(1):103-112.
- 17. Tomizawa D, Tawa A, Watanabe T, et al. Excess treatment reduction including anthracyclines results in higher incidence of relapse in core binding factor acute myeloid leukemia in children. *Leukemia*. 2013;27(12):2413-2416.
- 18. Shiraishi Y, Fujimoto A, Furuta M, et al. Integrated analysis of whole genome and transcriptome sequencing reveals diverse transcriptomic aberrations driven by somatic genomic changes in liver cancers. *PLoS One*. 2014;9(12):e114263.
- 19. de Hoon MJ, Imoto S, Nolan J, Miyano S. Open source clustering software. Bioinformatics. 2004;20(9):1453-1454.
- 20. Saldanha AJ. Java Treeview--extensible visualization of microarray data. Bioinformatics. 2004;20(17):3246-3248.
- Yamato G, Shiba N, Yoshida K, et al. RUNX1 mutations in pediatric acute myeloid leukemia are associated with distinct genetic features and an inferior prognosis. Blood. 2018;131(20):2266-2270.
- 22. Meshinchi S, Alonzo TA, Stirewalt DL, et al. Clinical implications of FLT3 mutations in pediatric AML. Blood. 2006;108(12):3654-3661.
- Gale RE, Green C, Allen C, et al; Medical Research Council Adult Leukaemia Working Party. The impact of FLT3 internal tandem duplication mutant level, number, size, and interaction with NPM1 mutations in a large cohort of young adult patients with acute myeloid leukemia. *Blood.* 2008;111(5): 2776-2784.
- 24. Linch DC, Hills RK, Burnett AK, Khwaja A, Gale RE. Impact of FLT3(ITD) mutant allele level on relapse risk in intermediate-risk acute myeloid leukemia. Blood. 2014;124(2):273-276.
- Xu F, Taki T, Yang HW, et al. Tandem duplication of the FLT3 gene is found in acute lymphoblastic leukaemia as well as acute myeloid leukaemia but not in myelodysplastic syndrome or juvenile chronic myelogenous leukaemia in children. Br J Haematol. 1999;105(1):155-162.
- Beillard E, Pallisgaard N, van der Velden VH, et al. Evaluation of candidate control genes for diagnosis and residual disease detection in leukemic patients using 'real-time' quantitative reverse-transcriptase polymerase chain reaction (RO-PCR): a Europe against cancer program. *Leukemia*. 2003;17(12): 2474-2486.

- 27. Balgobind BV, Hollink IH, Reinhardt D, et al. Low frequency of MLL-partial tandem duplications in paediatric acute myeloid leukaemia using MLPA as a novel DNA screenings technique. *Eur J Cancer.* 2010;46(10):1892-1899.
- Jo A, Mitani S, Shiba N, et al. High expression of EVI1 and MEL1 is a compelling poor prognostic marker of pediatric AML. *Leukemia*. 2015;29(5): 1076-1083.
- 29. Kanda Y. Investigation of the freely available easy-to-use software 'EZR' for medical statistics. Bone Marrow Transplant. 2013;48(3):452-458.
- 30. Murakami N, Okuno Y, Yoshida K, et al. Integrated molecular profiling of juvenile myelomonocytic leukemia. Blood. 2018;131(14):1576-1586.
- 31. Tokumasu M, Murata C, Shimada A, et al. Adverse prognostic impact of KIT mutations in childhood CBF-AML: the results of the Japanese Pediatric Leukemia/Lymphoma Study Group AML-05 trial. *Leukemia*. 2015;29(12):2438-2441.
- 32. Grisendi S, Mecucci C, Falini B, Pandolfi PP. Nucleophosmin and cancer. Nat Rev Cancer. 2006;6(7):493-505.
- 33. Collado M, Serrano M. The TRIP from ULF to ARF. Cancer Cell. 2010;17(4):317-318.
- Lim G, Choi JR, Kim MJ, et al. Detection of t(3;5) and NPM1/MLF1 rearrangement in an elderly patient with acute myeloid leukemia: clinical and laboratory study with review of the literature. Cancer Genet Cytogenet. 2010;199(2):101-109.
- Falini B, Nicoletti I, Bolli N, et al. Translocations and mutations involving the nucleophosmin (NPM1) gene in lymphomas and leukemias. *Haematologica*. 2007;92(4):519-532.
- Romana SP, Radford-Weiss I, Ben Abdelali R, et al; Groupe Francophone de Cytogénétique Hématologique. NUP98 rearrangements in hematopoietic malignancies: a study of the Groupe Francophone de Cytogénétique Hématologique. Leukemia. 2006;20(4):696-706.
- Guastadisegni MC, Lonoce A, Impera L, et al. CBFA2T2 and C20orf112: two novel fusion partners of RUNX1 in acute myeloid leukemia. *Leukemia*. 2010;24(8):1516-1519.
- Quentin S, Cuccuini W, Ceccaldi R, et al. Myelodysplasia and leukemia of Fanconi anemia are associated with a specific pattern of genomic abnormalities that includes cryptic RUNX1/AML1 lesions. *Blood.* 2011;117(15):e161-e170.
- Deluche L, Joha S, Corm S, et al. Cryptic and partial deletions of PRDM16 and RUNX1 without t(1;21)(p36;q22) and/or RUNX1-PRDM16 fusion in a case of progressive chronic myeloid leukemia: a complex chromosomal rearrangement of underestimated frequency in disease progression? Genes Chromosomes Cancer. 2008;47(12):1110-1117.
- 40. Sakai I, Tamura T, Narumi H, et al. Novel RUNX1-PRDM16 fusion transcripts in a patient with acute myeloid leukemia showing t(1;21)(p36;q22). Genes Chromosomes Cancer. 2005;44(3):265-270.
- 41. Roche-Lestienne C, Deluche L, Corm S, et al; Fi-LMC group. RUNX1 DNA-binding mutations and RUNX1-PRDM16 cryptic fusions in BCR-ABL+ leukemias are frequently associated with secondary trisomy 21 and may contribute to clonal evolution and imatinib resistance. *Blood.* 2008;111(7): 3735-3741.
- Hazourli S, Chagnon P, Sauvageau M, Fetni R, Busque L, Hébert J. Overexpression of PRDM16 in the presence and absence of the RUNX1/PRDM16 fusion gene in myeloid leukemias. Genes Chromosomes Cancer. 2006;45(11):1072-1076.
- Stevens-Kroef MJ, Schoenmakers EF, van Kraaij M, et al. Identification of truncated RUNX1 and RUNX1-PRDM16 fusion transcripts in a case of t(1; 21)(p36;q22)-positive therapy-related AML. *Leukemia*. 2006;20(6):1187-1189.
- 44. Quelen C, Lippert E, Struski S, et al. Identification of a transforming MYB-GATA1 fusion gene in acute basophilic leukemia: a new entity in male infants. Blood. 2011;117(21):5719-5722.
- 45. Belloni E, Shing D, Tapinassi C, et al. In vivo expression of an aberrant MYB-GATA1 fusion induces leukemia in the presence of GATA1 reduced levels [published correction appears in *Leukemia*. 2011;25(12):1922]. *Leukemia*. 2011;25(4):733-736.
- 46. lacobucci I, Wen J, Meggendorfer M, et al. Genomic subtyping and therapeutic targeting of acute erythroleukemia. Nat Genet. 2019;51(4):694-704.
- 47. Strehl S, Nebral K, König M, et al. ETV6-NCOA2: a novel fusion gene in acute leukemia associated with coexpression of T-lymphoid and myeloid markers and frequent NOTCH1 mutations. *Clin Cancer Res.* 2008;14(4):977-983.
- Weerkamp F, Baert MR, Brugman MH, et al. Human thymus contains multipotent progenitors with T/B lymphoid, myeloid, and erythroid lineage potential. Blood. 2006;107(8):3131-3137.
- 49. Liu Y, Easton J, Shao Y, et al. The genomic landscape of pediatric and young adult T-lineage acute lymphoblastic leukemia. *Nat Genet.* 2017;49(8): 1211-1218.
- 50. Kim YJ, Yang JJ, Han Y, et al. A rare case of ETV6/MECOM rearrangement in therapy-related acute myeloid leukemia with t(3;12) and monosomy 7. *Clin Lab.* 2017;63(2):415-418.
- 51. Yuasa H, Oike Y, Iwama A, et al. Oncogenic transcription factor Evi1 regulates hematopoietic stem cell proliferation through GATA-2 expression. *EMBO J.* 2005;24(11):1976-1987.
- 52. Balgobind BV, Raimondi SC, Harbott J, et al. Novel prognostic subgroups in childhood 11q23/MLL-rearranged acute myeloid leukemia: results of an international retrospective study. *Blood.* 2009;114(12):2489-2496.
- 53. Grier DG, Thompson A, Kwasniewska A, McGonigle GJ, Halliday HL, Lappin TR. The pathophysiology of HOX genes and their role in cancer. *J Pathol.* 2005;205(2):154-171.
- 54. Tiberi G, Pekowska A, Oudin C, et al. PcG methylation of the HIST1 cluster defines an epigenetic marker of acute myeloid leukemia. *Leukemia*. 2015; 29(5):1202-1206.