

Identification of germline variants in adults with hemophagocytic lymphohistiocytosis

Peter G. Miller,¹⁻³ Abhishek Niroula,^{1,3} John J. Ceremski,⁴ Christopher J. Gibson,¹ Martin S. Taylor,⁵ Sebastian Birndt,⁶ Florian Perner,¹ Jon Arnason,⁷ Adam S. Sperling,^{1,3} Mridul Agrawal,^{1,2} Alison M. Schram,⁸ Sarah Nikiforow,¹ German Pihan,⁷ Robert P. Hasserjian,⁵ Jon C. Aster,⁹ Paul La Rosée,¹⁰ Elizabeth A. Morgan,⁹ Nancy Berliner,² and Benjamin L. Ebert^{1,3,11}

¹Department of Medical Oncology, Dana-Farber Cancer Institute, and ²Division of Hematology, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA; ³Broad Institute of MIT and Harvard, Cambridge, MA; ⁴Harvard-MIT Program in Health Sciences and Technology, and ⁵Department of Pathology, Massachusetts General Hospital, Harvard Medical School, Boston, MA; ⁶Klinik für Innere Medizin II, Universitätsklinikum Jena, Jena, Germany; ⁷Department of Medical Oncology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA; ⁸Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; ⁹Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA; ¹⁰Klinik für Innere Medizin II, Schwarzwald-Baar-Klinikum, Villingen-Schwenningen, Germany; and ¹¹Howard Hughes Medical Institute, Dana-Farber Cancer Institute, Boston, MA

Key Points

- Some germline variants are predicted to disrupt protein function in HLH-associated genes.
- Such variants are neither enriched in adult-onset HLH nor associated with specific clinical or laboratory features of HLH.

Introduction

Hemophagocytic lymphohistiocytosis (HLH) is a life-threatening disorder of immune system over-activation that occurs as familial and acquired forms.¹ HLH is characterized by excessive cytokine production and inflammation, mediated by multiple immune cells including persistently activated macrophages. Familial HLH (FHL) is typically diagnosed in childhood and is often caused by inherited biallelic, deletion, or truncating variants in genes regulating the cytotoxic function of T lymphocytes and natural killer cells.^{2,3} By contrast, acquired HLH usually occurs in the setting of malignancy, infection, or autoimmune disease, and may be diagnosed at any age. Prior studies using in silico prediction algorithms have concluded that germline HLH-associated variants are enriched in adult patients with HLH but have been limited in the number of genes analyzed, incomplete clinical annotation to confirm true HLH diagnoses, and the relatively small size of the adult cohorts. Finally, the comparatively young ages at the time of HLH onset have made distinguishing FHL that occurs in early adulthood from true adult-onset HLH difficult.⁴⁻⁶ To overcome these issues, we sought to identify potential pathogenic germline variants in 17 genes implicated in FHL or other inherited immune disorders in a highly annotated cohort of patients diagnosed with HLH in adulthood.⁷

Methods

After obtaining institutional review board approval, genomic DNA was isolated from bone marrow aspirate or peripheral blood samples from adults treated in our hospital systems between 2001 and 2018 who met HLH-2004 diagnostic criteria; although there was incomplete information for some patients (particularly natural killer cell activity; see supplemental Figure 1A), all patients reported here met HLH-2004 criteria.⁸ Hybrid capture was performed on the samples using a custom SureSelect system (Agilent Technologies) targeting the exons of 17 genes implicated in FHL that occur in the context of other inherited immune disorders and have been described in the literature as associating with HLH, or are tested on commercial HLH panels (*AP3B1*, *BLOC1S6*, *BTK*, *CD27*, *IL2RG*, *ITK*, *LYST*, *MAGT1*, *PRF1*, *RAB27A*, *SH2D1A*, *SLC7A7*, *STX3*, *STX11*, *STXBP2*, *UNC13D*, *XIAP*).^{2,7,9} Sequencing was performed on the Illumina platform, and germline variant calling was done using the GATK Haplotype caller pipeline.¹⁰ Common variants were excluded by removing alterations with a genome aggregation database (gnomAD)¹¹ frequency of >0.05. We predicted the effect of variants on protein function using 5 in silico tools (PON-P2, PROVEAN, FATHMM, M-CAP, and REVEL). Variants classified as disruptive (predicted to impair normal protein function) by at least 2 tools were considered disruptive if none of the tools classified the variant to be nondisruptive. Variants classified as nondisruptive by a single tool were still considered disruptive if 3 or more tools classified the variant as

Submitted 25 November 2019; accepted 29 January 2020; published online 9 March 2020. DOI 10.1182/bloodadvances.2019001272.

The original sequencing data are available upon request to the corresponding author (benjamin_ebert@dfci.harvard.edu).

The full-text version of this article contains a data supplement.
© 2020 by The American Society of Hematology

Table 1. Characteristics of 88 patients studied in cohort

Total patients	N = 88
Median age (range), y	54 (18-81)
Median age, n (%), y	
<40	23 (26)
40-59	33 (38)
>60	32 (36)
Sex, n (%)	
Female	40 (45)
Male	48 (55)
Self-reported ethnicity, n (%)	
White or Caucasian	68 (77)
Hispanic or Latino	5 (6)
Asian	5 (6)
Black or African American	2 (2)
South Asian/Indian	1 (1)
Unknown/not reported	7 (8)
Precipitating etiology, n (%)	
Malignancy	43 (49)
Infection	26 (30)
Autoimmune disease	15 (17)
Multiple	14 (16)
Idiopathic	20 (23)
HLH-04 criteria, n (%)	
5*	26 (30)
6	38 (43)
7	21 (24)
8	3 (3)

*Three patients were determined to meet at least 5 HLH-04 criteria by the treating clinician but had 4 documented criteria in the clinical database due to incomplete information (see "Methods" and supplemental Figure 1A).

likely disruptive.¹² The effect of splice site variants was predicted as previously described.¹³ Statistical analysis for the enrichment of variants in the HLH cohort compared with the control population was performed using a Fisher's exact test with $P < .05$ considered significant.

Results and discussion

Our cohort included 88 adult patients with HLH (Table 1). The median age at diagnosis was 54 years old (range, 18-81 years) and the majority were male (55%) (supplemental Figure 1B). All patients met HLH-2004 criteria.^{8,14,15} A precipitating etiology was identified in most patients (77%) including malignancy (49%), infection (30%), and autoimmune disease (17%).

Among the 17 genes sequenced, a total of 42 variants with a gnomAD frequency of <0.05 were identified in 45 patients. Of these, 7 variants were considered disruptive (predicted to have an impact on protein function) in 18 patients (20% of the cohort) (Figure 1A). The most common variant was *PRF1* A91V ($n = 12$; 14%), an alteration commonly reported in adult HLH that has been shown to impair lymphocyte cytotoxicity.^{16,17} Three patients had

SLC7A7 A91V, whereas the disruptive variants were restricted to individuals, including *PRF1* H222Q, *ITK* R581Q, *LYST* R2624W, *STX3* splicing, and a male patient with *SH2D1A* R55Q (*SH2DA1* is on chromosome X). Three patients harbored multiple disruptive variants (*PRF1* A91V and *PRF1* H222Q, *PRF1* A91V and *SLC7A7* A91V, and biallelic *PRF1* A91V). We did not identify disruptive variants in the other FHL genes (*UNC13D*, *STX11*, or *STXBP2*) (Figure 1B).

We next examined whether the 7 unique disruptive variants were enriched in our cohort relative to whole-exome data from 2504 patients in the 1000 Genome Project (TGP; Figure 1C). Disruptive variants were identified in the TGP using the same gene set, pipeline, filtering, and algorithms used in our cohort. The specific variants in our cohort were significantly less common in the TGP (20% vs 5%; $P < .001$), an unsurprising result given the rarity of individual variants and requirement for a gnomAD frequency of <0.05 . To avoid this selection bias, we compared the frequency of disruptive variants in our cohort to the frequency of any disruptive variant in the 17 tested genes in the TGP. Again, our cohort had a higher frequency compared with the TGP (20% vs 9%; $P < .01$). However, this difference was no longer significant when restricting our cohort to patients who reported as "white" or "Caucasian" ($n = 68$, the most common ancestry in our cohort, hereafter referred to as "white") and comparing those to European populations in the TGP ($n = 503$; 20% vs 13%; $P = .13$). Furthermore, although the *PRF1* A91V variant was more common in our cohort compared with the TGP (14% vs 3%; $P < .001$), it was not more common when restricting to European populations (17% vs 9%; $P = .08$). Furthermore, after excluding patients with a *PRF1* A91V variant, there was no difference in the frequency of patients with disruptive variants between our cohort and the TGP (4.4% in both groups; Figure 1C). Of note, there was no difference in the frequency of variants among whites (14 of 68; 21%) and nonwhites (4 of 20; 20%) in the cohort.

Finally, we investigated whether the presence of a disruptive variant was associated with any clinical feature. The presence of a disruptive variant was not associated with age (mean age, 53 vs 51 years; $P = .64$), sex (33% vs 49% female; $P = .22$; supplemental Figure 2), or a specific precipitating etiology including malignancy (19% vs 22%; $P = .79$). Nor was there a significant difference in peak ferritin levels, soluble interleukin 2 receptor (sIL-2R) levels, or the sIL-2R-to-ferritin ratio (a metric often elevated in patients with lymphoma-associated HLH).¹⁸ We did not find a difference in the frequency of variants between patients meeting 5 to 6 HLH-04 criteria (13 of 64; 20%) and those meeting 7 to 8 criteria (5 of 24; 21%). Similarly, no differences were noted in those with and without a *PRF1* A91V variant (supplemental Figure 3). Survival analysis was not possible given the number of potential covariates (such as age, sex, ancestry, precipitating etiology, prior/subsequent treatment).

Our data suggest that disruptive germline variants do not drive adult HLH, unlike FHL. In contrast to cases of FHL, biallelic alterations, deletions, nonsense, and frameshift mutations were extremely rare, consistent with data from animal models that either biallelic loss or polygenic mutations can drive disease.¹⁹ Furthermore, disruptive variants were not enriched in adult HLH compared with control when accounting for ancestry, particularly when considering




Figure 1. Frequency and distribution of predicted disruptive variants in the HLH cohort. (A) Schematic of germline variant frequency within cohort among 17 HLH-associated genes sequenced. (B) Frequency of total and predicted disruptive germline variants observed in FHL genes in cohort. (C) Percentage of individuals in HLH and the 1000 Genomes Project (TGP) cohort carrying variants in HLH-associated genes. TGP var indicates the percentage of individuals in the TGP cohort carrying variants identified in the HLH cohort; TGP rule indicates the percentage of individuals in the TGP cohort carrying any variants identified by the same rules as for the HLH cohort. Percentages compared whole data (HLH [n = 88] and TGP [n = 2503]) and white or Caucasian individuals (HLH [n = 68] and TGP [n = 503]). Among white or Caucasian individuals, similar analysis was performed by excluding the *PRF1* A91V variant. The percentage of individuals carrying predicted disruptive variants is shown in red. **P* < .05; •*P* ≥ .05. The significance test result for carriers of all variants is indicated in black; the same for carriers of predicted disruptive variants is shown in red. + indicates the *SH2D1A* R55Q mutation, which is encoded on the X chromosome, was present in a male patient.

non-*PRF1* A91V variants. Additionally, there were no clinical differences between patients with and without disruptive variants. Finally, the contrast between the high frequency of disruptive variants in the control population (9%) and the rarity of adult HLH suggests that either the in silico algorithms overestimate the disruptive effects of variants or these disruptive variants are not strong drivers of disease. Either interpretation raises questions about the role and routine testing of germline variants in adults with HLH. Indeed, the fact that some of the predicted disruptive mutations including *PRF1* A91V are considered of uncertain significance by other metrics like the American College of Medical Genetics (ACMG) further calls into question the role of these mutations in adult HLH pathogenesis.²⁰ Our data, however, do not negate the importance of testing for genetic variants in cases in which FHL is highly suspected (including strong family history and young age of onset) and may influence treatment decisions (such as donor selection for allogeneic hematopoietic stem cell transplant).

Our study has several limitations. Like most prior studies, our cohort is limited in size given the rarity of HLH and difficulty obtaining usable biospecimens. However, this cohort contrasts with prior adult HLH reports including strict diagnostic inclusion criteria, significantly higher median age (at least 10 years older), expanded gene panel, and incorporation of ancestry (an essential consideration when studying germline genetics). Furthermore, whereas prior studies compared the frequency of the specific disruptive variants in their cohort to control populations, we compared the frequency of any disruptive variant in a gene previously associated with HLH

in our cohort to a control population, and further assessed the importance of ancestry and *PRF1* A91V in this comparison. Another limitation is the use of in silico algorithms to predict the effect of mutations on protein function, an approach used in nearly all prior studies of adult HLH. However, we used newer algorithms that, although still limited, have shown better performance than those previously used.¹² We also note that both ACMG and commercial testing often incorporate these predictive algorithms so understanding their relevance has clinical implications.²⁰ To highlight the importance of these issues and the variability across studies, we contrast our results with a recent report of 112 adults with HLH in China that concludes that germline variants are enriched in this population (43%); though in agreement with our study, they were predominantly missense and monoallelic.⁶ The studies differ in the ancestral composition of the cohort, older age of our population (which excludes cases of pediatric HLH), differing algorithms to predict disruption of protein function, and our approach to look for any disruptive variant in control populations (not just the frequency of those found in the group with HLH). In contrast, another study from China found that just 5.1% of adults with HLH had a germline variant; but again, the average patient age, gene panel sequenced, and disruptive prediction algorithm differed, and there was no formal comparison with a control population.²¹ Ultimately, additional efforts to better define disruptive variants in larger numbers of genes, including whole-exome/genome analyses, combined with expanded cohorts of adults with HLH of varying ancestries, will further refine our understanding of genetic contribution to the disease. Our data suggest that familial and

adult-onset HLH have a distinct pathophysiology and do not support routine germline testing of adults with HLH.

Acknowledgments

The authors thank Aaron Thorner and Anwesha Nag from the Dana-Farber Center for Cancer Genome Discovery for their help in sequencing the sample and David Steensma for review of the manuscript.

P.G.M. was supported by a National Institutes of Health (NIH), National Heart, Lung, and Blood Institute fellowship training grant (4T32HL116324-04) and an American Society of Hematology Research Training Award for Fellows. A.N. was supported by a Knut and Alice Wallenberg Foundation postdoctoral scholarship (2017.0436). A.S.S. was supported by a Ruth L. Kirschstein National Research Service Award and a Conquer Cancer Foundation Young Investigator Award. M.A. received support from the Deutsche Forschungsgemeinschaft (AG252/1-1). This work was supported by NIH, National Heart, Lung, and Blood Institute grant R01HL082945 and NIH, National Cancer Institute grant P01CA108631, the Howard Hughes Medical Institute, the Edward P. Evans Foundation, and the Leukemia & Lymphoma Society (B.L.E.).

References

1. Schram AM, Berliner N. How I treat hemophagocytic lymphohistiocytosis in the adult patient. *Blood*. 2015;125(19):2908-2914.
2. Brisse E, Wouters CH, Matthys P. Advances in the pathogenesis of primary and secondary haemophagocytic lymphohistiocytosis: differences and similarities. *Br J Haematol*. 2016;174(2):203-217.
3. Cetica V, Sieni E, Pende D, et al. Genetic predisposition to hemophagocytic lymphohistiocytosis: report on 500 patients from the Italian registry. *J Allergy Clin Immunol*. 2016;137(1):188-196.e4.
4. Zhang K, Jordan MB, Marsh RA, et al. Hypomorphic mutations in PRF1, MUNC13-4, and STXBP2 are associated with adult-onset familial HLH. *Blood*. 2011;118(22):5794-5798.
5. Tesi B, Lagerstedt-Robinson K, Chiang SC, et al. Targeted high-throughput sequencing for genetic diagnostics of hemophagocytic lymphohistiocytosis. *Genome Med*. 2015;7:130.
6. Miao Y, Zhu HY, Qiao C, et al. Pathogenic gene mutations or variants identified by targeted gene sequencing in adults with hemophagocytic lymphohistiocytosis. *Front Immunol*. 2019;10:395.
7. Chandrakasan S, Filipovich AH. Hemophagocytic lymphohistiocytosis: advances in pathophysiology, diagnosis, and treatment. *J Pediatr*. 2013;163(5):1253-1259.
8. Henter JI, Horne A, Aricó M, et al. HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48(2):124-131.
9. Cincinnati Children's Hospital Laboratory of Genetics and Genomics. Hemophagocytic Lymphohistiocytosis Gene Sequencing Panel. <https://www.cincinnatichildrens.org/-/media/cincinnati%20childrens/home/service/d/dchi/default/familial-hemophagocytic-lymphohistiocytosis.pdf>. Accessed 25 October 2019.
10. McKenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010;20(9):1297-1303.
11. Lek M, Karczewski KJ, Minikel EV, et al; Exome Aggregation Consortium. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*. 2016;536(7616):285-291.
12. Niroula A, Vihinen M. How good are pathogenicity predictors in detecting benign variants? *PLOS Comput Biol*. 2019;15(2):e1006481.
13. Jian X, Boerwinkle E, Liu X. In silico prediction of splice-altering single nucleotide variants in the human genome. *Nucleic Acids Res*. 2014;42(22):13534-13544.
14. Henter JI, Aricó M, Egeler RM, et al. HLH-94: a treatment protocol for hemophagocytic lymphohistiocytosis. HLH study Group of the Histiocyte Society. *Med Pediatr Oncol*. 1997;28(5):342-347.
15. Henter JI, Elinder G, Ost A; The FHL Study Group of the Histiocyte Society. Diagnostic guidelines for hemophagocytic lymphohistiocytosis. *Semin Oncol*. 1991;18(1):29-33.
16. Jordan MB, Hildeman D, Kappler J, Marrack P. An animal model of hemophagocytic lymphohistiocytosis (HLH): CD8+ T cells and interferon gamma are essential for the disorder. *Blood*. 2004;104(3):735-743.
17. Voskoboinik I, Sutton VR, Ciccone A, et al. Perforin activity and immune homeostasis: the common A91V polymorphism in perforin results in both presynaptic and postsynaptic defects in function. *Blood*. 2007;110(4):1184-1190.

Authorship

Contribution: P.G.M., N.B., and B.L.E. initiated the project, designed the research, and wrote the paper with input from other authors; P.G.M., A.N., J.C., C.J.G., A.S.S., M.A., and A.M.S. performed the research; P.G.M., A.N., and C.J.G. analyzed the data; and S.B., F.P., J.A., G.P., R.P.H., M.S.T., J.C.A., E.A.M., P.L.R., and S.N. contributed vital new reagents or analytical tools.

Conflict-of-interest disclosure: B.L.E. has received research funding from Celgene and Deerfield, consulting fees from GRAIL, and is on the scientific advisory boards for Exo Therapeutics and Skyhawk Therapeutics. P.G.M. has received consulting fees from Foundation Medicine, Inc. The remaining authors declare no competing financial interests.

ORCID profiles: P.L.R., 0000-0002-6758-7809; E.A.M., 0000-0001-5880-9337.

Correspondence: Benjamin L. Ebert, Dana-Farber Cancer Institute, 450 Brookline Ave, D1610A, Boston, MA 02215; e-mail: benjamin_ebert@dfci.harvard.edu.

18. Tsuji T, Hirano T, Yamasaki H, Tsuji M, Tsuda H. A high sIL-2R/ferritin ratio is a useful marker for the diagnosis of lymphoma-associated hemophagocytic syndrome. *Ann Hematol*. 2014;93(5):821-826.
19. Sepulveda FE, Garrigue A, Maschalidi S, et al. Polygenic mutations in the cytotoxicity pathway increase susceptibility to develop HLH immunopathology in mice. *Blood*. 2016;127(17):2113-2121.
20. Richards S, Aziz N, Bale S, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
21. Wang Y, Wang Z, Zhang J, et al. Genetic features of late onset primary hemophagocytic lymphohistiocytosis in adolescence or adulthood. *PLoS One*. 2014;9(9):e107386.