

Population pharmacokinetic modeling of factor concentrates in hemophilia: an overview and evaluation of best practice

Tine M. H. J. Goedhart,¹ Laura H. Bukkems,² C. Michel Zwaan,¹ Ron A. A. Mathôt,^{2,*} and Marjon H. Cnossen,^{1,*} for the OPTI-CLOT study group and SYMPHONY consortium

¹Department of Pediatric Hematology and Oncology, Erasmus MC Sophia Children's Hospital, University Medical Center Rotterdam, Rotterdam, The Netherlands; and ²Department of Clinical Pharmacology - Hospital Pharmacy, Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands

The accuracy of pharmacokinetic (PK)-guided dosing depends on the clinical and laboratory data used to construct a population PK model, as well as the patient's individual PK profile. This review provides a detailed overview of data used for published population PK models for factor VIII (FVIII) and factor IX (FIX) concentrates, to support physicians in their choices of which model best suits each patient. Furthermore, to enhance detailed data collection and documentation, we do suggestions for best practice. A literature search was performed; publications describing prophylactic population PK models for FVIII and FIX concentrates based on original patient data and constructed using nonlinear mixed-effect modeling were included. The following data were collected: detailed demographics, type of product, assessed and included covariates, laboratory specifications, and validation of models. Included models were scored according to our recommendations for best practice, specifically scoring the quality of data documentation as reported. Respectively, 20 models for FVIII and 7 for FIX concentrates were retrieved. Although most models (22/27) included pediatric patients, only 4 reported detailed demographics. The wide range of body weights suggested that overweight and obese adults were represented. Twenty-six models reported the assay applied to measure factor levels, whereas only 16 models named reagents used. Eight models were internally validated using a data subset. This overview presents detailed information on clinical and laboratory data used for published population PK models. We provide recommendations on data collection and documentation to increase the reliability of PK-guided prophylactic dosing of factor concentrates in hemophilia A and B.

Introduction

Patients with moderate and severe hemophilia receive replacement therapy with factor VIII (FVIII) and factor IX (FIX) concentrates to prevent spontaneous bleeding and bleeding after minor trauma.¹ Ahlberg et al reported fewer joint bleeds in patients with trough levels >0.01 IU/mL.² Therefore, clinicians historically use prophylactic replacement therapy to maintain trough factor levels of ≥ 0.01 IU/mL or higher in case of a severe bleeding phenotype by body weight-based dosing. However, body weight-based dosing does not account for inter-individual variability in pharmacokinetics (PK) of respective factor concentrates, that affect the achieved factor levels.³ As an example, body weight-based dosing may lead to higher dosing than necessary in obese patients because FVIII concentrate is distributed in the blood plasma, which does not increase proportionally with body weight.⁴ To address these inter-individual differences, PK-guided dosing using a posteriori Bayesian estimation is increasingly applied.⁵ This methodology relates an individual's

Submitted 28 April 2021; accepted 10 June 2021; prepublished online on *Blood Advances* First Edition 8 September 2021; final version published online 26 October 2021. DOI 10.1182/bloodadvances.2021005096.

*M.H.C. and R.A.A.M. are joint last authors.

© 2021 by The American Society of Hematology. Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

measured factor levels to the PK observed in a population to obtain more reliable individual PK parameters.⁶ To perform PK-guided dosing, accurate and validated population PK models are obligatory to enable the ability to prospectively predict data reliably.⁷

The accuracy of PK-guided dosing of factor concentrates is both dependent on the clinical and laboratory data used to construct the individual PK profile of the patient and on the quality of the underlying population PK model if Bayesian forecasting is applied. The clinical characteristics of the individual patient undergoing PK-guided dosing should preferably correspond to the population used to develop the population PK model because allometric exponents may otherwise not be accurate. If the characteristics do not correspond, the model may need to be further validated or enriched with such a specific patient group. However, population PK models are generally constructed with data from drug trials that often do not include specific patient populations such as children or older and obese patients. Consequently, estimation of their PK characteristics may be less accurate. In the same way, if factor levels of an individual patient are measured with different assays than with which the factor levels were measured within a certain population PK model, this may affect estimation of the individual PK parameters. As an example, a specific B-domain depleted recombinant FVIII should be measured by the chromogenic substrate assay (CSA) because factor levels are significantly lower when measured by one-stage assay (OSA), therefore affecting PK parameters and subsequently population PK modeling.⁸ Discrepancies between results obtained by CSA and OSA have been extensively discussed in literature and are especially relevant when extended half-life (EHL) FVIII and FIX concentrates are considered.⁹⁻¹¹ In addition, variation in assay results may also be caused by variation in reagents used within an assay. Moreover, the source of the applied factor-deficient plasma, and instruments such as analyzers and the material used to calibrate, may be of influence on measured factor activity levels, in particular levels when measuring below 0.10 IU/mL.^{12,13}

Therefore, clinicians and clinical pharmacologists should be aware of the specifications with regard to clinical and laboratory data of the individual patient undergoing PK-guided treatment, as well as the data with which the population PK model has been constructed. Recently, Preijers et al discussed available population PK models for FVIII and FIX concentrates, focusing on the methods used to construct these models, key features, and established covariate relationships.¹⁴ Our review aims to provide a detailed overview of the clinical and laboratory data used to construct available population PK models for both standard half-life (SHL) and EHL FVIII and FIX concentrates, as reported in literature. In this way, we aim to support physicians in their choices that model best suits each individual patient. Moreover, we propose some suggestions for best practice with regard to data collection and documentation after evaluating these models in detail to increase the reliability of PK-guided dosing and to aid in future population PK model construction.

Methods

Literature search strategy

We performed the following literature search to identify publications found on PubMed: (search date January 26, 2021): (haemophilia* OR hemophilia*) AND ("VIII"[Tiab] OR "IX"[Tiab]) AND ("population pharmacokinetic*" [Tiab] OR "pharmacokinetic model*" [Tiab] OR

"population pharmacokinetic analysis" [Tiab] OR "population PK" [Tiab]) NOT ("monkey*" [Tiab] OR "mice*" [Tiab] OR "dog*" [Tiab] OR "rabbit*" [Tiab] OR "rat*" [Tiab]).

First, publications were selected by title and abstract. Reading full text, we included publications that described a new population PK model for prophylactic treatment with both SHL and EHL FVIII or FIX concentrates. In addition, models were required to be based on original patient data and had to be constructed using nonlinear mixed-effect modeling. Backward citations were screened to include additional studies. Population PK models for von Willebrand factor (VWF) containing concentrates were excluded. The screening process was performed by 2 independent authors (M.G., L.B.).

Data collection

We collected data from both the publications describing the development of the population PK model as well as from publications of the underlying clinical trials. The following clinical data of the study populations were collected: total number of patients, number of children (age < 18 years), age; morphometric variables: body weight, body mass index (BMI), ideal body weight (IBW), lean body weight (LBW), body surface area (BSA), fat-free mass (FFM); endogenous factor level; factor concentrate; and assessed and included covariates and number of patients used for validation by data splitting. The collected laboratory data included the following items: type of assay used (OSA/CSA) and further specifications, namely reagent, calibrator, deficient plasma, and analyzer. Missing data were kindly requested from respective pharmaceutical companies or by correspondence with authors of the included publications.

Criteria to establish best practice

Based on our expert opinion and literature, we defined 12 clinical criteria of data documentation that should be reported in publications of population PK models. Thereafter, our included models were evaluated accordingly as complete (+), incomplete (\pm), absent (-), or not applicable. To specifically evaluate the quality of data documentation of included publications, only data were used as reported in the respective publications for this separate best practice evaluation. This is in contrast to the overview tables (Tables 1-4), in which all available and requested data are included. Documentation of 3 main demographic characteristics (number, age, and body weight) were evaluated for both the total population and pediatric populations separately. The first criterion (number) was scored as complete when the exact number of included (pediatric) patients was reported. If a minimum of pediatric patients was reported, we evaluated this item as incomplete. The criterion's age, body weight, and other morphometric variables such as BMI, LBW, FFM, and IBW were scored "complete" if both measures of location (eg, median, mean) and dispersion (eg, range, standard error) were available as is scientifically common practice, and "incomplete" and "absent," respectively, if 1 or both of those elements were lacking. In addition, if no exact age of the included children was given but only numbers of patients per age group (eg, between 6 and 12 years) was reported, age was defined as incomplete. As data splitting is a powerful method for model evaluation,¹⁵ we scored the criterion validation as complete if a data subset was used (comprising patients not used in the calibration population) to validate the constructed model. If other internal validations were executed, such as simulations and/or goodness-of-fit plots, we labeled the criteria as absent. Next, covariate analysis was evaluated as complete if the performance of a covariate analysis was reported (or the

Table 1. Literature search and detailed data on patient populations used to construct available FVIII population PK models

Study	Factor concentrate	Endogenous FVIII, IU/mL	Total population				Pediatrics		
			N	Age, y (range)	BW, kg (range)	Morphometric variables (range)	N	Age, y (range)	BW, kg (range)
Standard half-life factor concentrates									
Stass, 2006 ¹⁷	Kogenate-FS	<0.01-0.05	19	Mean 13 (4-12)	Mean 56 (21-96)	LBW mean 44.8 (18-66) BSA mean 1.5 (21-96)	19	Mean 13 (4-12)	Mean 56 (21-96)
Bolon-Larger, 2007 ³⁹	NA	<0.01-0.19	33	40 (7-77)	68 (21-120)	BSA 1.76 (0.85-2.32)	NA	NA	NA
Bjorkman, 2009 ¹⁹	Several*	<0.01-0.05	34	24 (7-74)	68 (26-124)	NA	11	12 (7-17)	44 (26-82)
Bjorkman, 2012 ²⁹	Advate	≤0.02	100	19 (10-66)	68 (35-108)	NA	45	NA (1-6)	>35
			52	4 (1-6)	16 (11-27)	NA	52	4 (1-6)	16 (11-27)
Karafoulidou, 2009 ⁴²	ReFacto	<0.01-0.17	28	34 (18-70)	75 (54-104)	NA	–	–	–
Abrantes, 2017 ¹¹	ReFacto, Xyntha	>0.01-0.40	754	23 (0-73)	69 (3-134)	NA	234	NA (0-17)†	NA
Jiménez-Juste, 2015 ⁴³	NovoEight	<0.01	76	21 (1-60)	57‡ (12-107)	BMI 21.4 (12-34)§	30	6‡ (1-17)	21 (12-66)‡
Tiede, 2020 ⁴⁴	NovoEight	<0.01	231	20 (1-60)	63 (12-120)	BMI 23.7 (14-40)	87	8 (1-17)	27 (12-95)
Garmann, 2017 ³¹	Kovaltry	<0.01	183	22 (1-61)	60 (11-124)	BMI 20.4 (13-38) LBW 54.1 (9-79)	>51¶	6 (1-11)	23 (11-59)
Shah, 2017 ⁴⁵	Kovaltry	<0.01	18	36 (19-64)	80 (55-99)	BMI 26.1 (19-29)	–	–	–
McEneny-King, 2019 ²⁰	All SHL	0.05	310	NA (1-62)	660 (10-132)	NA	NA	NA	NA
Standard and extended factor concentrates									
Allard, 2020 ¹⁸	Several#	0.01-mild	258	30 (3-77)	64 (15-130)	BMI 21.3 (13-45)	87	10 (3-17)‡	37 (15-109)‡
Extended half-life factor concentrates									
Zhang, 2017 ⁴⁶	Afstyla**	<0.01	91	29 (12-60)	72 (38-106)	BMI 23.7 (15-38)	10‡	16 (12-17)‡	50 (37-100)‡
			39	5 (1-11)	23 (10-87)	BMI 16.5 (13-30)	39	5 (1-11)	23 (10-87)
Delavenne, 2018 ⁴⁷	Nuwiq	<0.01	115	31 (2-67)	70 (13-140)	NA	29	Otherwise†	NA
Nesterov, 2015 ²³	Elocta	<0.01	180	NA (12-65)	(42-127)‡‡	NA	NA ^a	NA	NA
Bukkems, 2020 ²⁴	Elocta	≤0.02	43	28 (5-70)	72 (20-113)	BMI 23.73 (13-34)‡	13‡	10 (5-17)‡	34 (20-113)‡
Shah, 2019 ²¹	Jivi, Elocta	<0.01	18	34 (22-65)	NA	BMI 25.0 (19-30)	–	–	–
Solms, 2020 ³⁷	Jivi	<0.01	198	29 (2-62)	67 (12-126)	BMI 22.0 (13-42) LBW 49.0 (10-75)	153	NA (2-17) ^b	NA
Solms, 2020 ²²	Jivi and Adynovi	<0.01	18	34 (23-56)	NA	BMI 24.45 (18-30)	–	–	–
Chelle, 2020 ³⁸	Adynovi	<0.01	154	19 (3-72)	70 (15-150)	BMI 23.2 (14-5)	NA	NA ^c	NA

All data are reported in median (range) unless otherwise specified. NA, not available.

*Kogenate, Immunate, Helixate Nex Gen, Monoclate, Octanativ-MK.

†Number of patients per age group: age 0-1 y, N = 62; age 1-2 y, N = 21; age 2-6 y, N = 8; age 6-12 y, N = 25; age 12-17 y, N = 118.

‡Data provided by authors/pharmaceutical company on request.

§BMI <18 y 16.2 (12.4-25.6), ≥18 y 23.5 (15.3-33.8).

||BMI based on 168 patients (24 children <18 y and 144 adults).

¶Possibly 20 extra children from Leopold 1 and 2 study aged ≥12 y but not reported how many were included in final model.

#Factane, Advate, Kogenate, Kovaltry, Afstyla, Refacto, NovoEight, Elocta.

**PK model pooled from the 2 PK studies; data shown separately.

††Number of patients per age group: age 2-5 y, N = 29; age 6-12 y, N = 30; age 12-18 y, N = 2.

‡‡Number of patients and median body weight per subgroup: subgroup 1, N = 16, median 82.7; subgroup 2, N = 164, median 71.6.

^aA maximum of 14 children aged ≥12 y until 18 y was included. No exact number available.

^bTherapeutic indicated and administered to patients ≥12 y. Number of patients per age group: age < 6 y, N = 32; age 6-12 y, N = 29; age >12 y, N = 13.

^cTherapeutic indicated and administered to patients ≥12 y.

explanation why analysis was not performed), otherwise as absent. “Laboratory assay” was scored as complete, if the applied assay (OSA or CSA) was reported. If the other 4 further laboratory test specifications (applied reagent, calibrator, deficient plasma, and analyzer) were reported, this criterion was assessed as complete. If at least 1 of the 4 or no specification was reported, the publication was scored

as incomplete or absent, respectively. Finally, we assessed whether the applied assays and reagents corresponded to the recommendations for that factor concentrate according to UK Haemophilia Centre Doctors' Organization guidelines.¹⁶ This guideline describes, for each licensed and available product in the United Kingdom, specific reagents that are regarded as suitable, or should be avoided. If the

Table 2. Literature search and detailed data on patient populations used to construct available FIX population PK models

Author, y	Factor concentrate	Endogenous FIX, IU/mL	Total population				Pediatrics		
			N	Age, y (range)	BW, kg (range)	Other morphometric variables (range)	N	Age, y (range)	BW, kg (range)
Standard half-life factor concentrates									
Bjorkman, 2012 ³⁰	pdFIX*	≤0.02	26	39 (16-65)	70 (47-115)	NA	NA	NA	NA
Brekkan, 2016 ⁴⁸	pdFIX†	≤0;02	34	Mean 28 (10.9 SD)	Mean 67 (13.5 SD)	NA	>16‡	NA‡	NA
Bjorkman, 2013 ⁴⁹	BeneFIX	<0.01-0.05	56	23 (4-56)	NA (18-133)	NA	22	Otherwise§	NA
Suzuki, 2016 ²⁸	BeneFIX	0.01-0.02	201	12 (0-69)	44 (1-173)	NA	>88	NA	NA
Extended half-life factor concentrates									
Diao, 2014 ²⁷	Alprolix	≤0.02	135	31 (12-76)	73 (45-187)	NA	11	NA	NA
Collins, 2012 ²⁵	Refixia	<0.013	15	30 (21-55)	78 (47-101)	BMI 25.2 (18-29)	–¶	–	–
Zhang, 2016 ²⁶	Idelvion	≤0.02	104	26 (1-61)	64 (11-132)	BMI 21.8 (13-63)	38	8 (1-17)	28 (11 –71)

All data are reported in median (range) unless otherwise specified.

pdFIX, plasma-derived FIX.

*Alphanine, Mononine, Preconativ, Nanotiv, Replenine-VF.

†Alphanine, Mononine, Preconativ, Nanotiv, factor IX Grifols, Immunine, Octanin.

‡At least 16 pediatric patients with minimal age of 12 y.

§Number of patients and mean age per age group: age 4-9 y, N = 11, mean 6; age 10-19 y, N = 10, mean 13.

||Data provided by authors/pharmaceutical company on request.

¶Therapeutic indicated as administered to patients ≥12 y.

model was constructed using data obtained, by applying a suitable reagent, or if no specific recommendations were given by the guidelines for the specific factor concentrate, we evaluated this item as complete (+). If specific recommendations were given by Gray et al, but the applied reagent was neither defined as suitable, nor as a reagent that should be avoided, the criterion was evaluated criteria as incomplete (±). If models used factor concentrates that were not included in the UK guideline at time of publication, or if the reagents were not specified, scoring this criterion was not applicable (–).

Results

The initial search yielded 85 publications, of which 26 were included and 59 were excluded based on the set inclusion criteria. In addition, we included 1 supplementary article by backward citation,¹⁷ leading to a total of 27 included publications.

FVIII population PK models

A population PK model for FVIII concentrates was reported in 20 publications (Table 1). Of these 20, 6 combined data of multiple concentrates.^{11,18-22} Eleven publications described a population PK model for a SHL FVIII concentrate, 8 publications for an EHL FVIII concentrate, and 1 publication for both SHL and EHL FVIII concentrates.

Patient characteristics

Pediatric patients (<18 years) were included in most population PK models (16/20), although the exact number of included children could not be found or obtained in 5 population PK models. Details on the age of the children of 12 publications were collected and showed that all pediatric age groups were represented. Of the remaining 4 models that included children, only Nesterov et al restricted inclusions to children ≥12 years of age.²³ Strikingly, we were only able to retrieve body weight of pediatric patients in 9/20 models.

Overweight and obese patients seem to be included in most studies based on the total maximum reported body weight. Details on other morphometric variables of the total population were reported in 13/20 publications (Table 1). BMI was mostly presented (11 publications), followed by LBW (3 publications) and BSA (2 publications). Of the 7 studies that did not report other morphometric variables, 3 assessed such a variable as a covariate but did not incorporate it in the model.

Model covariates and validation

Table 3 depicts all evaluated and included covariates. The following other morphometric variables were assessed and included as covariates: BMI by respectively 4 and no publication(s), BSA by 3 and 1 publication(s), LBW by 5 and 3 publications, and FFM by 2 and 2 publications. Remarkably, 5/7 models for EHL factor concentrates evaluated the incorporation of VWF levels, in contrast to 1 model for a SHL factor concentrate. Of these 6 models, 4 models included VWF as a covariate.

Table 3 also shows that 6 models have been validated using a subset of the data, in addition to validation by simulations. The ratio between the number of included patients used to develop the model (calibration dataset) and the number of patients used to validate the model (validation dataset) varied between 0.17 and 1.27. Bukkems et al²⁴ externally validated and enriched the model by Nesterov et al.²³

Laboratory data

Laboratory data are presented in Table 4. Both OSA and CSA were used to measure the FVIII levels. Moreover, 4 models applied both methods. The reagents used were described in 12/20 models, resulting in a total of 14 different reagents. Only 4 publications reported all 4 laboratory specifications.

Table 3. Assessed and included covariates in population PK models

Author, y	Validation, N*	Assessed covariates	Included covariates
FVIII population PK models			
Stass, 2006 ¹⁷	–	Age, BW, HT, BSA, LBW, HCT, HG, WBC count, platelets, BG, PTT, INR, AST, ALT, Sodium, MCV, MCH, VWF:Ag, comedication	LBW
Bolon-Larger, 2007 ³⁹	18	Age, BW, BSA	BW, BSA
Bjorkman, 2009 ¹⁹	16	Age, BW, baseline FVIII, preparation and BG	Age, BW, baseline FVIII
Bjorkman, 2012 ²⁹	–	Age, BW	Age, BW
Karafoulidou, 2009 ⁴²	–	Age, BW, viral status	BW, viral status
Abrantes 2017 ¹¹	–	Age, BW, race, inhibitor status and titer, assay, LBW, TBW	Age, BW, race, inhibitor, assay
Jiménez-Juste, 2015 ⁴³	–	Age, BW	Age, BW
Tiede, 2020 ⁴⁴	–	Age, BW	Age, BW
Garmann 2017 ³¹	–	Age, BW, HT, BMI, LBW, race	LBW
Shah, 2017 ⁴⁵	–	–	–
McEneny-King, 2019 ²⁰	394	Age, BW, FFM, brand	Age, BW, FFM, brand
Allard, 2020 ¹⁸	–	Age, BW, brand, structure of FVIII, EHL	Age, BW, EHL
Zhang, 2017 ⁴⁶	–	Age, BW, BMI, AST, ALT, CrCl, VWF, HCT, HCV status, antidrug antibody, geographical region, race, initial PK assessment vs repeat	BW, VWF
Delavenne, 2018 ⁴⁷	20	Age, BW, HT, IBW, BSA	Age, BW
Nesterov, 2015 ²³	28†	Age, BW, HT, race, BG, HCT, VWF:Ag, albumin, NNA presence, IgG1 concentrations, HCV status, HIV status	Age, BW, VWF:Ag, HCV status, HCT
Bukkems, 2020 ²⁴	–	Age, BW, BG 0, HCT, VWF:Ag, presence of target joints	VWF:Ag, BW
Shah 2019 ²¹	–	LBW, VWF	LBW
Solms, 2020 ³⁷	–	Age, BW, HT, BMI, LBW, VWF:Ag, race	LBW, VWF:Ag
Solms, 2020 ²²	–	–‡	–
Chelle, 2020 ³⁸	26	Age, BW, HT, FFM, assay	Age, FFM, assay
FIX population PK models			
Bjorkman, 2012 ³⁰	–	Age, BW, FIX concentrate preparation	Age, BW, FIX concentrate
Brekkan, 2016 ⁴⁸	–	Age, BW, FIX concentrate, baseline FIX activity	BW, baseline FIX activity
Bjorkman, 2013 ⁴⁹	–	Age, BW	BW
Suzuki, 2016 ²⁸	72	Age, BW, race	BW
Diao, 2014 ²⁷	100	BW, Albumin, race	BW
Collins 2012 ²⁵	–	Unknown	Unknown
Zhang, 2016 ²⁶	–	Age, BW, dose, BMI, AST, ALT, CrCl, HCV status, antidrug antibody, region	BW, weight adjusted dose

ALT, alanine transaminase levels; AST, aspartate transaminase levels; BW, body weight; BG, blood group; CrCl, creatinine clearance; FFM, fat-free mass; HCT, hematocrit; HCV, hepatitis C virus; HT, height; HG, hemoglobin; IgG1, immunoglobulin; INR, international normalized ratio; MCH, mean corpuscular hemoglobin; MCV, mean corpuscular volume; NNA, non-neutralizing antibody; PTT, prothrombin time; TBW, total body water; VWF:Ag, von Willebrand factor antigen; WBC, white blood cell.

*Validation using a data subset.

†Externally validated by Bukkems et al.

‡Because of the small study size, no additional covariate search was conducted.

FIX population PK models

Four population PK models for SHL FIX concentrates and 3 population PK models for EHL FIX concentrates were reviewed (Table 2). The models included a single recombinant FIX concentrate, with the exception of 2 models which combine multiple plasma-derived FIX concentrates.

Patient characteristics

Except for the nonacog beta pegol (Refixia) population PK model,²⁵ all other 6 models included pediatric patients (<18 years), of which 3 only included children 12 years and older. However, only 3 of the 6 models reported the exact number children, although 2 models showed the number of included pediatric patients per age group.

Morphometric variables for pediatric patients were only available for the population model reported by Zhang et al. In descriptions of the total population, body weight was presented for all models and BMI for 2 models. Furthermore, only 1 model by Zhang et al assessed the inclusion of other morphometric variables as covariates (Table 3). Despite this lack of data, the parameter “patient’s body weight” showed a wide inter-individual variation in all studies. Strikingly, Zhang et al included patients up to a BMI of 63.1.²⁶

Model covariates and validation

As shown in Table 3, only Collins et al did not report assessed and included covariates. As also depicted in Table 3, only 2 of the 7 models were internally validated by data splitting.^{27,28} The number of patients included in the calibration and validation dataset in the model

Table 4. Laboratory data specifications used in population PK models

Author, y	Assay	Reagent	Calibrator	Deficient plasma	Analyzer
FVIII population PK models					
Stass, 2006 ¹⁷	OSA	NA	NA	NA	MDA 180 (BioMerieux)
Bolon-Larger, 2007 ³⁹	NA	NA	NA	NA	NA
Bjorkman, 2009 ¹⁹	CSA	Coatest FVIII kit	Coatest FVIII kit	Coatest FVIII kit	KabiVitrium
Bjorkman, 2012 ²⁹	OSA	NA	NA	20th British Standard	NA
Karafoulidou, 2009 ⁴²	CSA	Coamatic FVIII	RLS (Wyeth)	Diagnostica Stago	Chromogenix instrumentation Laboratory SpA
Abrantes 2017 ¹¹	OSA and CSA	NA	NA	NA	NA
Jiménez-Juste, 2015 ⁴³	OSA	SynthASil	NA	Standard Human Plasma	Coasys Plus, Roche
Tiede, 2020 ⁴⁴	OSA	SynthASil	NA	Standard Human Plasma	Coasys Plus, Roche
Garmann 2017 ³¹	CSA	NA	NA	NA	NA
Shah, 2017 ⁴⁵	CSA	NA	NA	Standard Human Plasma	NA
McEneny-King, 2019 ²⁰	OSA	NA	NA	NA	NA
Allard, 2020 ¹⁸	OSA	STA R-STA-CK Prest, Pathrombin, SynthAsil	NA	NA	NA
Zhang, 2017 ⁴⁶	CSA	Coamatic FVIII	NA	Coamatic FVII kit	Behring Coagulation System
Delavenne, 2018 ⁴⁷	OSA and CSA	Trinity Biotech & Coamatest SP	NA	NA	Siemens BCS-XP
Nesterov, 2015 ²³	OSA	Actin FSL (Dade)	NA	SHP (Precision Biologics CRYOcheck)	Siemens BCS XP
Bukkems, 2020 ²⁴	OSA and CSA	Several*,†	Several*,†	Several*,†	Several*,†
Shah 2019 ²¹	OSA	HemosL SynthASil	NA	Standard Human Plasma	ACL Advance System
Solms, 2020 ³⁷	OSA and CSA	HemosL SynthAFax & BIOPHEN chromogenic assay kit (HYPHEN)	NA	NA	ACL Advance System
Solms, 2020 ²²	OSA	HemosL SynthASil	NA	Standard Human Plasma	ACL Advance System
Chelle, 2020 ³⁸	OSA and CSA	Several*,‡	Several*,‡	Several*,‡	Several*,‡
FIX population PK models					
Bjorkman, 2012 ³⁰	OSA	NA	NA	NA	NA
Brekkan, 2016 ⁴⁸	OSA	Organon, Diagnostika Stago, Dade Behring Marburg GmbH	NA	NA	NA
Bjorkman, 2013 ⁴⁹	OSA	NA	NA	Standard Human Plasma	NA
Suzuki, 2016 ²⁸	OSA	NA	NA	NA	NA
Diao, 2014 ²⁷	OSA	From Trinity Biotech	NA	Precision Biologic	NA
Collins 2012 ²⁵	OSA	Trinity Automated aPTT	N9-Gp-calibrator	Precision Biologic	Siemens BCS-XP
Zhang, 2016 ²⁶	OSA	Pathromtin SL	NA	Standard Human Plasma	Behring Coagulation System

*Data provided by authors/pharmaceutical company on request.

†Specifications of model by Bukkems et al: reagents: HemosL SynthASil, HS Tcoag, Actin FS, Biophen Kit,

Tcoag Trinichrom, Siemens Bovine, Hemosll Electrachrome, Siemens Chromogenic; calibrator: CRYOcheck, Stago calibrator, Siemens, Hyphen Biomed-Biophen, HemosL; deficient plasma: HemosL, Stago, Immunodeficient plasma, Siemens; analyzer: ACL TOP (Werfen-IL), Star Max Stago, Sysmex (Siemens).

‡All levels have been extracted from the Web-Accessible Population Pharmacokinetic Service Hemophilia (WAPPS-Hemo). All reagents were used with no 1 covering a significant larger size of sample than any other.

by Suzuki et al were, respectively, 201 and 72, and 135 and 100 in the model by Diao et al.

Laboratory data

All FIX levels were measured using the OSA for both SHL and EHL FIX concentrates (Table 4). The used reagent was stated in 4 of the

7 population PK models. Only Collins et al reported all 4 laboratory specifications.

Best practice

Figure 1 shows the evaluation of best practice of the included publications in this review. The scored items have been discussed

Author, year	Total population			Pediatric population			Other morphometric variables	Validation using data subset	Covariate analysis	Laboratory assay	Laboratory specifications	Recommended assay & reagent used *
	Number	Age (years)	Body weight (kg)	Number	Age (years)	Body weight (kg)						
Stass, 2006 ¹⁷	+	+	+	+	+	+	+	-	+	+	+	-
Bolon-Larger, 2007 ³⁹	+	+	+	-	-	-	+	+	+	-	-	-
Bjorkman, 2009 ¹⁹	+	+	+	+	+	+	-	+	+	+	+	-
Bjorkman, 2012 ²⁹	+	+	+	+	+	+	-	-	+	+	+	+
Karafoulidou, 2009 ⁴²	+	+	+	-	-	-	-	-	+	+	+	+
Abrantes, 2017 ¹²	+	+	+	+	+	-	-	-	+	+	-	+
Jiménez-Juste, 2015 ⁴³	+	+	+	+	+	+	+	-	+	+	+	+
Tiede, 2020 ⁴⁴	+	+	+	+	+	+	+	-	+	+	+	+
Garmann, 2017 ³²	+	+	+	+	+	+	+	-	+	+	-	-
Shah, 2017 ⁴⁵	+	+	+	-	-	-	+	-	+	+	+	-
McEneny-King, 2019 ²⁰	+	+	+	-	-	-	-	+	+	+	-	+
Allard, 2020 ¹⁸	+	+	+	+	-	-	+	-	+	+	+	+
Zhang, 2017 ⁴⁶	+	+	+	+	+	+	+	-	+	+	+	+
Delavenne, 2018 ⁴⁷	+	+	+	+	+	-	-	+	+	+	+	+
Nesterov, 2015 ²³	+	+	+	-	-	-	-	+	+	+	+	+
Bukkems, 2020 ²⁴	+	+	+	+	-	-	-	+	+	+	-	-
Shah, 2019 ²¹	+	+	-	-	-	-	+	-	+	+	+	+
Solms, 2020 ³⁷	+	+	+	+	+	-	+	-	+	+	+	+
Solms, 2020 ²²	+	+	-	-	-	-	+	-	-	+	+	+
Chelle, 2020 ³⁸	+	+	+	-	-	-	+	+	+	+	-	+
Bjorkman, 2012 ³⁰	+	+	+	-	-	-	-	-	+	+	-	-
Brekkan, 2016 ⁴⁸	+	+	+	+	-	-	-	-	+	+	+	-
Bjorkman, 2013 ⁴⁹	+	+	+	+	+	-	-	-	+	+	+	+
Suzuki, 2016 ²⁸	+	+	+	+	-	-	-	+	+	+	-	+
Diao, 2014 ²⁷	+	+	+	+	-	-	-	+	+	+	+	+
Collins, 2012 ²⁵	+	+	-	-	-	-	-	-	-	+	+	+
Zhang, 2016 ²⁶	+	+	+	+	+	-	-	-	+	+	+	+

Figure 1. Evaluation of included models according to our personal recommendations of best practice, specifically on the quality of data documentation in the publications. Results presented as complete (green, +), incomplete (orange, ±), absent (red, -) or not applicable (-). * According to UK guidelines by Grey et al. Haemophilia 2020. † Bukkems et al. have externally validated the model by Nesterov et al.

previously. As is clearly visible, most publications cannot be scored as complete according to our strict set of criteria because of a lack of detailed information in several domains.

Discussion

In this overview, we have collected detailed information on the demographic, clinical, and laboratory data used to construct 27 published population PK models for both SHL and EHL FVIII and FIX concentrates. We also introduce recommendations for best practices of data collection and documentation after evaluating the publications of included population PK models accordingly.

Patient characteristics

Importantly, because children have other PK characteristics than do adults, pediatric patients (mostly) of all ages were included in 22/27 PK populations. Body weight adjusted clearance decreases during normal development from child until adulthood, automatically affecting elimination half-life time because it is inversely related to clearance.²⁹ Notably, approximately one-half of these age-related differences disappear when an identical sampling strategy is performed in children as in adults, instead of a reduced sampling strategy. Therefore, we advise, if possible, measuring FVIII and FIX levels 15 to 30 minutes after infusion.³ When both pediatric and adult patients are included

in the data collection on the basis of which a population PK model is constructed, an exponential relationship between body weight, both total body weight and/or LBW, and clearance is introduced. This is also called allometric scaling. As a result, a population PK model can be used for patients of all ages.^{24,26,30} Allometric scaling is applied in all published models in this review, which is a positive finding. Historically, Bjorkman et al illustrates the importance of sufficient variation and volume of the included pediatric participation, as a poor population prediction for children of ages 3 to 10 years was observed with predicted FVIII levels 31% lower than observed FVIII levels because of inclusion of only 4 children from this age group.¹⁹ In Bukkems et al, it was shown that population clearance and central volume of distribution were underpredicted in patients aged < 12 years when applying a population PK model based solely on patients 12 years and older.²⁴

Overweight and obese adults are represented in most models. This is clinically relevant because of the increasing prevalence of overweight/obesity in 31% of hemophilia patients in Europe and North American, with concomitant risks of under- or overdosing of factor concentrates.^{31,32} With increasing BMI, studies show an increasing in vivo recovery of FVIII concentrate both in children and adults, with a decreasing steady state distribution volume.³³⁻³⁵ Unfortunately, no guidelines yet exist to optimize factor concentrate dosing in this patient group.

Van Moort et al reported that IBW-based dosing most accurately calculates doses of FVIII concentrate in comparison with other morphometric variables.³⁶ This approach led to both better targeting of FVIII levels, as well as a mean reduction of 48.9% in prophylactic FVIII concentrate consumption during a 3-month period.⁴ Interestingly however, in the included population PK models in this review, IBW was neither evaluated nor included as a covariate. Contrastingly, LBW was evaluated and included as a covariate in 4 and 5 publications in our review, respectively. LBW showed a linear correlation with central volume of distribution and a positive nonlinear relationship with clearance.^{11,17,21,31,37} Stass et al reported that scaling to LBW is superior to scaling to actual body weight. A third morphometric variable that was included in several models as a covariate was FFM, which seemed to correlate better to plasma volume than total body weight.^{20,38}

Model covariates and validation

Covariate analysis was described in all models, among which we specifically highlight the inclusion of VWF in the FVIII population PK models. Overall, a negative association between VWF and FVIII clearance is reported.^{23,24,37} In other words, the higher VWF level, the lower FVIII clearance. This is explained by the fact that VWF protects FVIII from proteolytic degradation and rapid clearance from the blood circulation.²³ In our review, VWF was evaluated and included as covariate in 6 and 4 FVIII population PK models, respectively. Stass et al reported VWF as an important covariate, although it was not identified as one in the performed analyses. This contradiction can be attributed to the fact that only sparse data were available to adequately test covariate analysis,¹⁷ as also reported by Chelle et al.³⁸ Therefore, when possible, we advise measurement of VWF antigen (VWF:Ag) levels during PK profiling and monitoring of FVIII concentrates to improve both PK guidance and model development.

Interestingly, only 8 models were validated by data splitting, a powerful, advanced method of internal validation. Although, the decision to

split available data into a calibration and a validation set of course depends on the number of included study patients.⁷ Notably, in the FVIII population PK models by Bolon-Larger and Bjorkman et al, the number of included patients in the calibration dataset was 33 and 34, respectively. These were relatively small numbers compared with the patients in the validation sets of these studies (eg, 18 and 16 patients). In contrast, McEneny-King et al included 310 and 394 patients in the calibration and validation dataset, respectively.

Laboratory data

All publications described which assay was used to measure the factor levels, except for the model by Bolon-Larger et al.³⁹ Because of the discrepancies between OSA and CSA and the effects on PK parameter estimation, it is essential that population PK models include results of both assays when both are used in clinical practice. Solms et al constructed separate population PK models based on OSA and CSA results and described that PK parameters were similar.³⁷ However, application of correction factors for OSA and CSA is also generally applied when other assays are used.^{24,38} For example, Bukkems et al calculated an exponential function of 1.06 to correct for FVIII levels in an enriched FVIII-Fc Fusion protein population PK model. This resulted in an increase of FVIII levels from 100 IU/dL when measured by OSA compared with 131 IU/dL when measured by CSA.²⁴

Because reagents have been identified as the most important cause of assay variability,¹⁶ it is striking that we were not able to identify the used reagent of 11/27 models. Interestingly, we identified 16 different documented reagents. Five models used several reagents, therefore increasing accuracy of PK guidance because of the included inter-reagent variability. To illustrate the importance of this covariate, external quality organizations for laboratories have reported variation between FIX measurements when using different OSA reagents between 2.0 and 19.0 IU/mL and when applying a golden standard of spiked plasma with recombinant FIX Fc fusion protein level of 6.0 IU/mL FIX.⁴⁰ These variations may lead to underestimations that amount to up to 75%.^{16,40} Understandingly, these differences may lead to errors in PK guidance of dosing, especially when targeting FVIII or FIX trough levels, if reagent variability is not captured in the population PK model.

Therefore, UK Haemophilia Centre Doctors' Organization guidelines provide helpful advice on the appropriate choice of assays and reagents for each factor concentrate, which is licensed for use in the United Kingdom.¹⁶ Recommendations are based on safeguarding of assay results within 20% and 30% of FVIII and/or FIX target levels based on potency label in samples spiked as >30 IU/dL and <10 IU/dL. In addition, with each recommendation, it is noted if sufficient data are available to substantiate the guidelines. No population PK models in this review used assays or reagents that were advised to avoid.¹⁶

Strengths and limitations

The importance of more insight into the construction of population PK models is highlighted by the comparative study by Preijers and van Moort et al. For identical patients, 3 available PK tools produced significantly different PK parameters and clinically relevant variation in doses of recombinant FVIII concentrate. This was due to the influence of the patient data on which the population PK model was based and the resulting PK parameters that affected individual estimations.⁴¹

Our review has some limitations. First, we may have missed population PK models because we only performed the literature search using

PubMed and reference searches; however, we do believe that we have included most population PK models used in clinical practice with this approach. Second, we did not succeed in contacting all authors, which may have affected the completeness of presented results. Next, in the formulation of best practice, we did not integrate the number of patients in the validation dataset relative to the number patients in the calibration set because experts have not reached consensus on aspect of validation.⁷ Finally, we did not apply ranking to the items incorporated in our recommendations for best practice. Despite these limitations, we believe this review provides a valuable and objective overview of the most relevant aspects needed to ensure reliability and feasibility of PK guidance of factor concentrates in hemophilia care.

Recommendations

Clinicians or clinical pharmacologists, who treat patients with PK-guided dosing, should evaluate whether the population PK model is suitable for each individual patient, based on the data used to construct the model. According to our evaluation of best practice in hemophilia, improvements can be made on data collection and documentation. If a factor concentrate is approved and registered for all age groups, we recommend that the corresponding population PK model includes pediatric patients and that their characteristics are described. Also, patients with both underweight and overweight/obesity should be included with documentation of morphometric variables such as BMI, LBW, IBM, or FFM. Next, new population PK models should be validated by a subset of the dataset or even more optimally by external validation to ensure optimal safety in standard clinical practice. Also, when performing PK guidance, it is essential that laboratory specifications, both of the patient populations used to construct the models as well as the patients who receive PK guidance, are reported and taken into account, preferably in accordance with international guidelines.

These recommendations with regard to detailed information on patient characteristics, laboratory assays and reagents, and validation strategies may of course also apply to PK guidance of novel and upcoming therapies in hemophilia with alternative administration routes, as well as expected combined treatment modalities such as emicizumab and on-demand FVIII concentrate, gene therapy, and on-demand FVIII or FIX concentrates.

Conclusion

This overview presents the wide variation in detail of included clinical and laboratory data used to construct 27 population PK models. By providing detailed information on these population PK models, the applicability and reliability of PK-guided prophylactic dosing of factor concentrates in hemophilia A and B can be investigated by (pediatric) hematologists. We also recommend best practice with regard to data collection to enable reliable PK guidance of treatment in patients with bleeding disorders as a whole for current and future therapies.

Acknowledgments

The SYMPHONY consortium, which aims to orchestrate personalized treatment in patients with bleeding disorders, is a unique collaboration between patients, health care professionals, and translational and fundamental researchers specializing in inherited bleeding disorders, as well as experts from multiple disciplines. It aims to identify best treatment choice for each individual based on bleeding phenotype. To achieve this goal, work packages

(WP) have been organized according to 3 themes (eg, Diagnostics [WPs 3 and 4], Treatment [WPs 5-9], and Fundamental Research [WPs 10-12]). This research received funding from the Netherlands Organization for Scientific Research (NWO) in the framework of the NWA-ORC Call grant agreement NWA.1160.18.038. Principal investigator: M.H. Cnossen; project manager: S.H. Reitsma. Beneficiaries of the SYMPHONY consortium: Erasmus MC and Erasmus MC Sophia Children's Hospital, University Medical Center Rotterdam, project leadership and coordination, Sanquin Diagnostics, Sanquin Research, Amsterdam University Medical Centers, University Medical Center Groningen, University Medical Center Utrecht, Leiden University Medical Center, Radboud University Medical Center, Netherlands Society of Hemophilia Patients, Netherlands Society for Thrombosis and Hemostasis, Bayer B.V., CSL Behring B.V., and Swedish Orphan Biovitrum (Belgium) BVBA/SPRL. This study was also performed as part of the OPTI-CLOT international multicenter research consortium, "Patient Tailored Pharmacokinetic (PK) Guided Dosing of Clotting Factor Concentrates and Desmopressin in Bleeding Disorders," which is currently WP 6 within the SYMPHONY consortium. This paper is written on behalf of the international multicenter OPTI-CLOT and To WiN studies that aim to implement a PK-guided approach for the treatment of bleeding disorders using population PK models for desmopressin, factor concentrates, and other alternative drugs. OPTI-CLOT and To WiN study group members are: Steering committee: M.H. Cnossen (principal investigator and chair), F.W.G. Leebeek, Erasmus MC Sophia Children's Hospital and Erasmus MC, University Medical Center Rotterdam, Rotterdam; R.A.A. Mathôt (co-leading investigator), K. Fijnvandraat, M. Coppens, Amsterdam University Medical Center, Amsterdam, University Medical Center, Amsterdam; K. Meijer, University Medical Center Groningen, Groningen; S.E.M. Schols, Radboud University Medical Centre, Nijmegen; H.C.J. Eikenboom, Leiden University Medical Centre, Leiden; R.E.G. Schutgens, University Medical Center Utrecht, Utrecht; E.A.M. Beckers, Maastricht University Medical Center, Maastricht; and P. Ypma, Haga Hospital, The Hague. Principal investigators and local collaborators in the Netherlands: M.J.H.A. Kruip, S. Polinder, Erasmus MC, University Medical Center Rotterdam, Rotterdam; R.Y.J. Tamminga, University Medical Centre Groningen, Groningen; P. Brons, Radboud University Medical Centre, Nijmegen; K. Fischer, K.P.M. van Galen, University Medical Centre Utrecht, Utrecht; F.C.J.I. Heubel-Moenen, Maastricht University Medical Centre, Maastricht; L. Nieuwenhuizen, Maxima Medical Centre, Eindhoven; M.H.E. Driessens, The Netherlands Hemophilia Patient Society; I. van Vliet, Erasmus MC, University Medical Centre Rotterdam, Rotterdam. OPTI-CLOT/To WiNs: J. Lock, H.C.A.M. Hazendonk, I. van Moort, J.M. Heijdra, M..H.J. Goedhart, W. Al Arashi, Erasmus MC, University Medical Center Rotterdam, Rotterdam; T. Preijers, N.C.B. de Jager, L.H. Bukkems, M.E. Cloesmeijer, A. Janssen, Amsterdam University Medical Centers, Amsterdam. Principal investigators and local collaborators in the United Kingdom—P.W. Collins, Arthur Bloom Haemophilia Centre, Institute of Infection and Immunity, School of Medicine, Cardiff University, Cardiff; R. Liesner, Great Ormond Street Haemophilia Centre, Great Ormond Street Hospital for Children NHS Trust, London; P. Chowdary, Katharine Dormandy Haemophilia Centre and Thrombosis Unit, Royal Free London NHS Foundation Trust, London; C.M. Millar, Hammersmith Hospital-Imperial College Healthcare NHS Trust, London; D. Hart,

Department of Haematology, The Royal London Hospital Barts Health NHS Trust, London; and D. Keeling, Oxford Haemophilia and Thrombosis Centre, Oxford University Hospitals, Churchill Hospital, Oxford.

Authorship

Contribution: T.M.H.J.G. and L.H.B. performed the literature search and collected data; T.M.H.J.G. analyzed the data and wrote the manuscript; R.A.A.M. and M.H.C. supervised the study and provided critical guidance during the analyses; L.H.B., C.M.Z., R.A.A.M., and M.H.C. critically reviewed the paper and provided comments; and all authors approved the final version of the manuscript.

Conflict-of-interest disclosure: M.H.C. has received grants from governmental and societal research institutes such as NWO, ZonMW, Innovation fund, from private funding institutions, institutional grants and unrestricted investigator research grants/ educational and travel funding from the following companies over

the years: Pfizer, Baxter/Baxalta/Shire, Bayer Schering Pharma, CSL Behring, Sobi Biogen, Novo Nordisk, Novartis, and Nordic Pharma, and has served as a member on steering boards of Roche and Bayer. All grants, awards, and fees go to the Erasmus MC as an institution. R.A.A.M. has received governmental and societal research institutes such as NWO, ZonMW, and Innovation Fund and unrestricted investigator research grants from Baxter/Baxalta/Shire/Takeda, Bayer, CSL Behring, and Sobi. He has served as an advisor for Bayer, CSL Behring, Merck Sharp & Dohme, and Baxter/Baxalta/Shire/Takeda. All grants and fees were paid to the institution.

ORCID profiles: T.M.H.J.G., 0000-0002-8919-8118; L.H.B., 0000-0001-7967-1023; R.A.A.M., 0000-0001-8401-0879; M.H.C., 0000-0003-1557-2995.

Correspondence: Marjon H. Cnossen, Erasmus MC Sophia Children's Hospital, University Medical Center Rotterdam, PO Box 2060, 3000 CB Rotterdam, The Netherlands; e-mail: m.cnossen@erasmusmc.nl.

References

1. Peyvandi F, Garagiola I, Young G. The past and future of haemophilia: diagnosis, treatments, and its complications. *Lancet*. 2016;388(10040):187-197.
2. Ahlberg A. Haemophilia in Sweden. VII. Incidence, treatment and prophylaxis of arthropathy and other musculo-skeletal manifestations of haemophilia A and B. *Acta Orthop Scand Suppl*. 1965;(suppl. 77):3-132.
3. Björkman S, Blanchette VS, Fischer K, et al; Advate Clinical Program Group. Comparative pharmacokinetics of plasma- and albumin-free recombinant factor VIII in children and adults: the influence of blood sampling schedule on observed age-related differences and implications for dose tailoring. *J Thromb Haemost*. 2010;8(4):730-736.
4. Graham A, Jaworski K. Pharmacokinetic analysis of anti-hemophilic factor in the obese patient. *Haemophilia*. 2014;20(2):226-229.
5. Ragni MV, Croteau SE, Morfini M, Cnossen MH, Iorio A; Subcommittee on Factor VIII, Factor IX, and Rare Bleeding Disorders. Pharmacokinetics and the transition to extended half-life factor concentrates: communication from the SSC of the ISTH. *J Thromb Haemost*. 2018;16(7):1437-1441.
6. Dargaud Y, Delavenne X, Hart DP, Meunier S, Mismetti P. Individualized PK-based prophylaxis in severe haemophilia. *Haemophilia*. 2018;24(Suppl 2):3-17.
7. Sherwin CM, Kiang TK, Spigarelli MG, Ensom MH. Fundamentals of population pharmacokinetic modelling: validation methods. *Clin Pharmacokinet*. 2012;51(9):573-590.
8. Santoro C, Iorio A, Ferrante F, et al. Performance of recalibrated ReFacto laboratory standard in the measurement of FVIII plasma concentration via the chromogenic and one-stage assays after infusion of recalibrated ReFacto (B-domain deleted recombinant factor VIII). *Haemophilia*. 2009;15(3):779-787.
9. Duncan EM, Duncan BM, Tunbridge LJ, Lloyd JV. Familial discrepancy between the one-stage and two-stage factor VIII methods in a subgroup of patients with haemophilia A. *Br J Haematol*. 1994;87(4):846-848.
10. Kihlberg K, Strandberg K, Rosén S, Ljung R, Astermark J. Discrepancies between the one-stage clotting assay and the chromogenic assay in haemophilia B. *Haemophilia*. 2017;23(4):620-627.
11. Abrantes JA, Nielsen EI, Korth-Bradley J, Harnisch L, Jönsson S. Elucidation of factor VIII Activity pharmacokinetics: a pooled population analysis in patients with hemophilia A treated with moroctocog alfa. *Clin Pharmacol Ther*. 2017;102(6):977-988.
12. Verbruggen B, Meijer P, Novákova I, Van Heerde W. Diagnosis of factor VIII deficiency. *Haemophilia*. 2008;14(s3 Suppl 3):76-82.
13. Kitchen S, Tiefenbacher S, Gosselin R. Factor activity assays for monitoring extended half-life FVIII and factor IX replacement therapies. *Semin Thromb Hemost*. 2017;43(3):331-337.
14. Preijers T, Schutte LM, Kruij M, et al. Population pharmacokinetics of clotting factor concentrates and desmopressin in hemophilia. *Clin Pharmacokinet*. 2020;60(1):1-16.
15. U.S. Department of Health and Human Services, Food and Drug Administrations. Population Pharmacokinetics Guidance for Industry. *Clin Pharmacol*. 2019.
16. Gray E, Kitchen S, Bowyer A, et al. Laboratory measurement of factor replacement therapies in the treatment of congenital haemophilia: a United Kingdom Haemophilia Centre Doctors' Organisation guideline. *Haemophilia*. 2020;26(1):6-16.
17. Stass H. Determination of minimal sampling time points for reliable pharmacokinetic evaluation of recombinant factor VIII - an exploratory population pharmacokinetic analysis in paediatric patients suffering from severe haemophilia. *Haemophilia*. 2006;12(s4):50-55.

18. Allard Q, Djerada Z, Pouplard C, et al. Real life population pharmacokinetics modelling of eight factors VIII in patients with severe haemophilia A: is it always relevant to switch to an extended half-life? *Pharmaceutics*. 2020;12(4):E380.
19. Björkman S, Folkesson A, Jönsson S. Pharmacokinetics and dose requirements of factor VIII over the age range 3-74 years: a population analysis based on 50 patients with long-term prophylactic treatment for haemophilia A. *Eur J Clin Pharmacol*. 2009;65(10):989-998.
20. McEneny-King A, Chelle P, Foster G, Keepanasseril A, Iorio A, Edginton AN. Development and evaluation of a generic population pharmacokinetic model for standard half-life factor VIII for use in dose individualization. *J Pharmacokinetic Pharmacodyn*. 2019;46(5):411-426.
21. Shah A, Solms A, Wiegmann S, et al. Direct comparison of two extended-half-life recombinant FVIII products: a randomized, crossover pharmacokinetic study in patients with severe hemophilia A. *Ann Hematol*. 2019;98(9):2035-2044.
22. Solms A, Shah A, Berntorp E, et al. Direct comparison of two extended half-life PEGylated recombinant FVIII products: a randomized, crossover pharmacokinetic study in patients with severe hemophilia A. *Ann Hematol*. 2020;99(11):2689-2698.
23. Nestorov I, Neelakantan S, Ludden TM, Li S, Jiang H, Rogge M. Population pharmacokinetics of recombinant factor VIII Fc fusion protein. *Clin Pharmacol Drug Dev*. 2015;4(3):163-174.
24. Bukkems LH, Heijdra JM, Mathias M, et al; for UK-EHL Outcomes Registry OPTI-CLOT Collaboration. A novel, enriched population pharmacokinetic model for recombinant factor VIII-Fc fusion protein concentrate in hemophilia A patients. *Thromb Haemost*. 2020;120(5):747-757.
25. Collins PW, Møss J, Knobe K, Groth A, Colberg T, Watson E. Population pharmacokinetic modeling for dose setting of nonacog beta pegol (N9-GP), a glycoPEGylated recombinant factor IX. *J Thromb Haemost*. 2012;10(11):2305-2312.
26. Zhang Y, Roberts J, Bensen-Kennedy D, et al. Population pharmacokinetics of a new long-acting recombinant coagulation factor IX albumin fusion protein for patients with severe hemophilia B. *J Thromb Haemost*. 2016;14(11):2132-2140.
27. Diao L, Li S, Ludden T, Gobburu J, Nestorov I, Jiang H. Population pharmacokinetic modelling of recombinant factor IX Fc fusion protein (rFIXFc) in patients with haemophilia B. *Clin Pharmacokinetic*. 2014;53(5):467-477.
28. Suzuki A, Tomono Y, Korth-Bradley JM. Population pharmacokinetic modelling of factor IX activity after administration of recombinant factor IX in patients with haemophilia B. *Haemophilia*. 2016;22(5):e359-e366.
29. Björkman S, Oh M, Spotts G, et al. Population pharmacokinetics of recombinant factor VIII: the relationships of pharmacokinetics to age and body weight. *Blood*. 2012;119(2):612-618.
30. Björkman S, Ahlén V. Population pharmacokinetics of plasma-derived factor IX in adult patients with haemophilia B: implications for dosing in prophylaxis. *Eur J Clin Pharmacol*. 2012;68(6):969-977.
31. Garmann D, McLeay S, Shah A, Vis P, Maas Enriquez M, Ploeger BA. Population pharmacokinetic characterization of BAY 81-8973, a full-length recombinant factor VIII: lessons learned - importance of including samples with factor VIII levels below the quantitation limit. *Haemophilia*. 2017;23(4):528-537.
32. Wilding J, Zourikian N, Di Minno M, et al. Obesity in the global haemophilia population: prevalence, implications and expert opinions for weight management. *Obes Rev*. 2018;19(11):1569-1584.
33. Henrard S, Hermans C. Impact of being overweight on factor VIII dosing in children with haemophilia A. *Haemophilia*. 2016;22(3):361-367.
34. Henrard S, Speybroeck N, Hermans C. Impact of being underweight or overweight on factor VIII dosing in hemophilia A patients. *Haematologica*. 2013;98(9):1481-1486.
35. McLeay SC, Morrish GA, Kirkpatrick CM, Green B. Encouraging the move towards predictive population models for the obese using propofol as a motivating example. *Pharm Res*. 2009;26(7):1626-1634.
36. van Moort I, Preijers T, Hazendonk H, et al. Dosing of factor VIII concentrate by ideal body weight is more accurate in overweight and obese hemophilia A patients. *Br J Clin Pharmacol*. 2021;87(6):2602-2613.
37. Solms A, Iorio A, Ahsman MJ, et al. Favorable pharmacokinetic characteristics of extended-half-life recombinant factor VIII BAY 94-9027 enable robust individual profiling using a population pharmacokinetic approach. *Clin Pharmacokinetic*. 2020;59(5):605-616.
38. Chelle P, Yeung CHT, Croteau SE, et al. Development and validation of a population-pharmacokinetic model for rurioctacog alfa pegol (Adynovate®): a report on behalf of the WAPPS-Hemo Investigators Ad Hoc Subgroup. *Clin Pharmacokinetic*. 2020;59(2):245-256.
39. Bolon-Larger M, Chamouard V, Bressolle F, Bouliou R. A limited sampling strategy for estimating individual pharmacokinetic parameters of coagulation factor VIII in patients with hemophilia A. *Ther Drug Monit*. 2007;29(1):20-26.
40. Nederlof A, Kitchen S, Meijer P, et al. Performance of factor IX extended half-life product measurements in external quality control assessment programs. *J Thromb Haemost*. 2020;18(8):1874-1883.
41. Preijers T, van Moort I, Fijnvandraat K, Leebeek FWG, Cnossen MH, Mathôt RAA; 'OPTI-CLOT' Study Group. Cross-evaluation of pharmacokinetic-guided dosing tools for factor VIII. *Thromb Haemost*. 2018;118(3):514-525.
42. Karafoulidou A, Suarez E, Anastasopoulou I, et al. Population pharmacokinetics of recombinant factor VIII:C (ReFacto) in adult HIV-negative and HIV-positive haemophilia patients. *Eur J Clin Pharmacol*. 2009;65(11):1121-1130.
43. Jiménez-Yuste V, Lejniece S, Klamroth R, et al. The pharmacokinetics of a B-domain truncated recombinant factor VIII, turoctocog alfa (NovoEight®), in patients with hemophilia A. *J Thromb Haemost*. 2015;13(3):370-379.
44. Tiede A, Abdul Karim F, Jiménez-Yuste V, et al. Factor VIII activity and bleeding risk during prophylaxis for severe hemophilia A: a population pharmacokinetic model [published online ahead of print 23 April 2020]. *Haematologica*.
45. Shah A, Solms A, Garmann D, et al. Improved pharmacokinetics with BAY 81-8973 versus antihemophilic factor (recombinant) plasma/albumin-free method: a randomized pharmacokinetic study in patients with severe hemophilia A. *Clin Pharmacokinetic*. 2017;56(9):1045-1055.

46. Zhang Y, Roberts J, Tortorici M, et al. Population pharmacokinetics of recombinant coagulation factor VIII-SingleChain in patients with severe hemophilia A. *J Thromb Haemost.* 2017;15(6):1106-1114.
47. Delavenne X, Dargaud Y, Ollier E, Négrier C. Dose tailoring of human cell line-derived recombinant factor VIII simoctocog alfa: Using a limited sampling strategy in patients with severe haemophilia A. *Br J Clin Pharmacol.* 2019;85(4):771-781.
48. Brekkan A, Berntorp E, Jensen K, Nielsen EI, Jönsson S. Population pharmacokinetics of plasma-derived factor IX: procedures for dose individualization. *J Thromb Haemost.* 2016;14(4):724-732.
49. Björkman S. Population pharmacokinetics of recombinant factor IX: implications for dose tailoring. *Haemophilia.* 2013;19(5):753-757.