

Single-cell DNA sequencing reveals complex mechanisms of resistance to quizartinib

Cheryl A. C. Peretz,¹⁻³ Lisa H. F. McGary,¹ Tanya Kumar,¹ Hunter Jackson,⁴ Jose Jacob,⁵ Robert Durruthy-Durruthy,⁵ Mark J. Levis,⁶ Alexander Perl,⁷ Benjamin J. Huang,^{3,8} and Catherine C. Smith^{1,8}

¹Department of Medicine, University of California, San Francisco, San Francisco, CA; ²Children's Hospital and Research Center Oakland, Oakland, CA; ³Department of Pediatrics and ⁴School of Medicine, University of California, San Francisco, San Francisco, CA; ⁵Mission Bio, Inc, San Francisco, CA; ⁶Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD; ⁷Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA; and ⁸Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, San Francisco, CA

Key Points

- Single-cell sequencing exposes previously unmeasurable complexity of tumor heterogeneity and clonal evolution on quizartinib.
- Single-cell sequencing reveals on- and off-target mechanisms of resistance to quizartinib, which can preexist therapy.

Introduction

Acute myeloid leukemia (AML) with activated Fms-like tyrosine kinase receptor III (*FLT3*) causes significant mortality secondary to relapsed/refractory (R/R) disease.¹⁻⁵ Drug resistance limits the duration of response of *FLT3* inhibitors (*FLT3i*). Genetic evolution leading to R/R disease is incompletely understood, and leukemia's heterogeneity is incompletely described, even with next-generation sequencing (NGS). A deeper understanding of cellular heterogeneity is essential because intratumoral heterogeneity is ubiquitous, has prognostic relevance, affects response to therapy, and drives therapeutic resistance.⁶⁻⁹ We characterized, via single-cell sequencing (SCS), the genetic evolution of R/R AML on the *FLT3i* quizartinib.

Case description

We analyzed serial samples from patients with R/R *FLT3*-positive internal tandem duplication (*ITD*⁺) AML on phase 2 or 3 quizartinib monotherapy trials.^{10,11} After treatment, 3 of 8 patients had complete response with incomplete hematologic recovery, 2 patients obtained a partial response, and 3 patients had no response; no patients obtained complete response.

Methods

We analyzed sixteen samples from bone marrow or peripheral blood from 8 patients (Table 1; supplemental Table 1). Included patients relapsed after initial response and had serial samples available for analysis.

We performed single-cell DNA sequencing on unsorted mononuclear cells using the Tapestry platform (Mission Bio, Inc).¹² This consisted of targeted sequencing of mutational hotspots using 40 amplicons from 19 AML-specific genes as previously described.¹³ Data from 2 × 150 bp paired-end FASTQ files generated by Illumina HiSeq4000 was processed. For analysis, the Tapestry Insights software, a custom GATK-based variant-calling workflow,¹⁴ the Integrative Genomics Viewer,¹⁵ and the Maftools R scripts were used. Single-cell phylogenies and population hierarchies were reconstructed. Further details are provided in supplemental Methods.

Results and discussion

SCS provides a sensitive description of cellular populations compared with bulk NGS

We analyzed 103 031 cells from 16 timepoints from 8 patients. Patient characteristics are provided in Table 1; additional clinical information is provided in supplemental Table 1. Targeted SCS identified pathogenic variants not detected by clinical bulk NGS (supplemental Table 2). SCS revealed details of

Submitted 16 September 2020; accepted 28 January 2021; published online 5 March 2021. DOI 10.1182/bloodadvances.2020003398.

All genetic data for the dbGaP study can be accessed at the NCBI Sequence Read Archive (SRA) under accession no. phs002320.v1.p1 (https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs002320.v1.p1). All code for custom sequencing analysis is uploaded to GitHub and/or published in open source

publications. For original data, please contact the corresponding author at catherine.smith@ucsf.edu.

The full-text version of this article contains a data supplement.

© 2021 by The American Society of Hematology

Table 1. Patient characteristics

Clinical patient ID	ELN risk* at diagnosis	Cytogenetics at study entry	Best response	Duration of response, wk	Sample timepoint	No. of single cells analyzed	No. of clones
1	Adverse	46,XY,del(5)(q23q33)[5]/46,XY[4]	PR	5	Pre-quizartinib	13 538	6
					Relapse	4 734	6
2	Adverse	47,XY,+11[15]	PR	6	Pre-quizartinib	637	3
					Relapse	3 245	6
					Second relapse†	1 056	6
3	Adverse	46, XX	CRi	6	Pre-quizartinib	11 518	6
					Relapse	4 113	3
4	Adverse	46, XX	CRi	3	Pre-quizartinib	5 557	3
					Relapse	4 741	4
5	Favorable	Unavailable	CRi	10	Pre-quizartinib	9 774	4
					Relapse	6 867	8
6	Adverse	47, XX, +8, t(x;10)	None	NA	Pre-quizartinib	6 537	4
					Relapse	7 961	7
7	Unknown	47,XX,+8[1]/47,idem,del(16)(q13)[19]	None	NA	Pre-quizartinib	5 285	5
					Relapse	10 347	13
8	Unknown	47,XX,+8[3]	None	NA	Relapse	7 121	4

CRi, complete response with incomplete hematologic recovery; NA, not available; PR, partial response.

*Per European LeukemiaNet 2017 Guidelines.²¹

†Patient received quizartinib + chemotherapy between first and second relapse.

ITD number (3 patients had >1 ITD) and zygosity (supplemental Table 3). Samples were notably polyclonal: a mean of 6 distinct clones per sample (range, 3-13) with a mean of 5 clones before treatment and 7 clones at the time of relapse (Figure 1A; supplemental Table 2).

Quizartinib drives clonal selection for preexisting off-target mutations

SCS reveals selection of previously undetectable preexisting subclones containing off-target (non-FLT3) mutations. Two patients (Figure 1B-C) relapsed with off-target Ras pathway mutations, which have been previously associated with clinical gilteritinib and crenolanib resistance,^{13,16} but not with quizartinib resistance. These data suggest that *RAS*-mutant cell populations expand on quizartinib. However, we acknowledge that secondary polyclonal *RAS* mutations which cooccur in the *FLT3* mutant clone¹⁷ may behave differently than those in separate subpopulations. One of these patients (Figure 1B) had a predominant *NRAS*-mutant population at relapse in the setting of polyclonal off-target resistance clones: 2 *NRAS* mutations (at the G13 and Q61 loci) in separate cell populations as well as an additional *KIT* mutant population. All of these populations expanded on quizartinib monotherapy. *KIT* mutations have not been previously associated with quizartinib resistance. The second patient (Figure 1C) had expansion of both preexisting off-target, *KRAS* G13D, and on-target, *FLT3* N841, mutations, confirming our previous observation that on- and off-target mechanisms of resistance can coexist within the same leukemia.¹⁸ At relapse, the dominant population contained a previously undetected *FLT3* D835Y, demonstrating that the preexisting subclone will not always drive resistance.

Quizartinib drives clonal selection for on-target FLT3 KD mutations

Seven of 8 patients developed at least 1 additional FLT3 kinase domain (KD) mutation, most commonly at the D835 locus (Figure 1C-H).^{19,20} SCS reveals how KD mutations segregate with driver ITD and other mutations. FLT3 KD mutations can occur on the native (FLT3-ITD negative) allele (Figure 1C-D), in *cis* with the FLT3-ITD allele (Figure 1E-G), or in a combination of both (Figure 1H).¹⁸ Quizartinib may select for preexisting KD mutant populations in native FLT3 alleles (Figure 1C) or KD mutant cells may appear de novo to dominate at the time of relapse (Figure 1D).

In addition to clonal selection, quizartinib drives clonal evolution of on-target FLT3-ITD KD mutations.


Within a FLT3-ITD population that dominates at relapse, SCS shows gain of FLT3 KD mutations in *cis* with ITD (Figure 1E-G; supplemental Figure 1). In addition to co-mutation with ITD, multiple KD mutations can coexist with each other within a single cell (Figure 1H), suggesting previously unmeasurable complexity, even at a single tumor-relevant locus.

Quizartinib drives complex evolution, which may lead to duplicative paths to resistance

Leukemias in 4 patients had polyclonal KD mutations. In all cases, the same KD mutations were present in multiple clones at same timepoint (Figure 1C,F-H). Though we cannot definitively rule out loss of heterozygosity, there is apparent parallel acquisition of these mutations in multiple patients (Figure 1F-H), emphasizing the importance of KD mutations in resistance as well as the leukemia's persistent dependence on FLT3 signaling. This is further highlighted by the fact that KD mutations may occur as heterozygous or

Figure 1. The genetic landscape of relapse on quizartinib monotherapy.

Evolution represents 1 possibility based on detectable mutational data. (A) Overview of the mutational landscape on quizartinib monotherapy with somatic mutations shown across 70 unique clones (1 clone per column) from 16 patient samples. (B) Clonal selection of off-target mutations. Mutations in *NRAS* G13, Q61, and *KIT* D816 are detectable with SCS before initiation of therapy, each in <0.5% of the population, before expanding on quizartinib. *RAS* and *KIT* mutations have not previously been documented as clinical resistance mutations. (C) Complex on- and off-target resistance. This leukemia had a KD mutation at N841 and a *KRAS* mutation at G13, which were both detectable before therapy and expanded on quizartinib. At the time of relapse, this leukemia has gained 2 different KD mutations at the D835 locus, one of which is present in heterozygous and homozygous populations. (D) Clonal selection of *FLT3* KD mutation. This patient relapsed with a single dominant clone with an on-target KD mutation at D835 in a non-ITD-containing population. This leukemia also has 2 different ITD mutations. (E) Clonal evolution with gain of *FLT3* KD mutation. This patient relapsed with a single, dominant clone with a D835Y mutation in cis with *FLT3*-ITD. (F) Complex *FLT3* evolution. Three different KD mutations emerge at relapse, including 2 at the D835 locus in different populations. There are also mutations with hetero- and homozygous *FLT3* ITD populations. (G) Complex *FLT3* evolution. This patient relapsed with 2 different KD mutations, including a D835H mutation that arises in 2 different populations. There are also mutations with hetero- and homozygous *FLT3* ITD populations. (H) Complex clonal evolution. At time of relapse, this patient had 13 clones, including 2 ITDs and 4 KD mutations; both SNV and MNV KD mutations may or may not be in *FLT3*-ITD⁺ clones.



homozygous alleles within the same leukemia (Figure 1C; supplemental Figure 2). The complexity of these KD mutations reveals more intricate tumor heterogeneity than previously appreciated, and also raises the question of whether certain leukemias are predisposed to mutate specific sites. One leukemia in particular highlights an extreme of mutational heterogeneity (Figure 1H) with 6 FLT3 aberrations in the same cell: 2 distinct ITDs (different insertion sites and lengths) and 4 different KD mutations including S838 and 3 multiallelic mutations at the D835 locus. Further, the D835V mutation occurred as both single- and multinucleotide variants (SNV and MNV) (ATC→AAC or CAC) (supplemental Figure 2). This patient exhibited additional MNVs creating the D835F and D835I alleles (supplemental Figure 2), which presumably arose as secondary mutations from the original D835V SNV (D835F: ATC→AAC→AAC; D835I: ATC→AAC→AAT). The striking duplicative polyclonality within this single locus implies that some AML patients may experience particularly high mutational burden including a predilection to mutations at specific resistance loci.

Serial SCS offers a novel look at the genetic evolution to relapse, with the ability to accurately determine zygosity, comutations, and clonal composition. Quizartinib binds FLT3 in its inactive conformation, leaving it vulnerable to resistance mutations that keep FLT3 in its active confirmation, like those in the TKD. This series corroborates FLT3 KD mutations as the most common mechanism of resistance to quizartinib; SCS further reveals that mutations may occur alone or in *cis* with ITD alleles. The ability to identify multiple KD mutations within the same leukemia, the same timepoint, the same cell, the same gene, and even the same locus, reveals striking biologic redundancy even within single cells. Additionally, we demonstrate for the first time that Ras pathway mutations are a mechanism of clinical resistance to quizartinib. Both off- and on-target mutations may exist before therapy. The ability to detect resistance mutations before overt clinical relapse and to visualize their expansion over time supports the need for effective combination therapy to suppress outgrowth of resistant clones. However, novel resistant subclones can arise to outcompete clones detectable before drug treatment; therefore, attempts to address only preexisting resistant clones may be insufficient to prevent relapse. In the future, SCS analysis may be used to prospectively monitor clonal populations on FLT3i treatment and facilitate dynamic adjustment of therapy.

Acknowledgments

The authors are grateful to the patients and their families for participation in clinical trials and providing samples to be used in this research.

This work was funded by the National Institutes of Health, National Center for Advancing Translational Sciences (award TR001871).

References

1. Abu-Duhier FM, Goodeve AC, Wilson GA, et al. FLT3 internal tandem duplication mutations in adult acute myeloid leukaemia define a high-risk group. *Br J Haematol*. 2000;111(1):190-195.
2. Ley TJ, Miller C, Ding L, et al; Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368(22):2059-2074.
3. Kaspers GJ, Zimmermann M, Reinhardt D, et al. Improved outcome in pediatric relapsed acute myeloid leukemia: results of a randomized trial on liposomal daunorubicin by the International BFM Study Group. *J Clin Oncol*. 2013;31(5):599-607.

This work was also supported by a Research Scholar Grant from the American Cancer Society (132032-RSG-18-063-01-TBG). C.A.C.P. is funded by a Young Investigator Award from the Alex's Lemonade Stand Foundation and a Conquer Cancer Foundation of the American Society of Clinical Oncology (ASCO) Young Investigator Award, supported by Association of Northern California Oncologists. B.J.H. is the recipient of a Rally Foundation Career Development Award and a St Baldrick's Foundation Scholar Award. C.C.S. is a Damon Runyon-Richard Lumsden Foundation Clinical Investigator supported (in part) by the Damon Runyon Cancer Research Foundation (CI-99-18).

The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health. Any opinions, findings, and conclusions expressed in this material are those of the author(s) and do not necessarily reflect those of the American Society of Clinical Oncology, Conquer Cancer, or Association of Northern California Oncologists.

Authorship

Contribution: C.A.C.P., L.H.F.M., T.K., and H.J. performed processing of patient samples and sequencing as well as data analysis; B.J.H., R.D.-D., and J.J. performed additional bioinformatic analyses; M.J.L. and A.P. provided clinical correlation data and patient samples; C.C.S. guided all experiments and preparation of the manuscript; and all authors reviewed and approved the final manuscript.

Conflict-of-interest disclosure: R.D.-D. and J.J. are full-time employees of Mission Bio, Inc. M.J.L. reports consultancy and research funding from FujiFilm, consultancy and honoraria from Daiichi Sankyo Inc, consultancy and honoraria from Agios and Menarini, consultancy and research funding from Novartis and Astellas, and consultancy and honoraria from Amgen. A.P. reports honoraria, advisory board, research funding from Novartis; consultancy, honoraria, and research funding from Astellas and Daiichi Sankyo; consultancy and honoraria from NewLink Genetics; consultancy from Arog; research funding from FujiFilm; consultancy and honoraria from AbbVie; consultancy from Forma; and advisory board and research funding from Loxo and Syndax. C.C.S. reports research funding from Astellas Pharma, Abbvie, Revolution Medicines, and FujiFilm. The remaining authors declare no competing financial interests.

ORCID profiles: C.A.C.P., 0000-0002-0855-6770; M.J.L., 0000-0003-0473-6982; A.P., 0000-0002-1463-2231; B.J.H., 0000-0001-6996-0833; C.C.S., 0000-0003-0160-7026.

Correspondence: Catherine C. Smith, University of California, San Francisco, 505 Parnassus Ave, Suite M1286, Box 1270, San Francisco, CA 94143; e-mail: catherine.smith@ucsf.edu.

4. Ward E, DeSantis C, Robbins A, Kohler B, Jemal A. Childhood and adolescent cancer statistics, 2014. *CA Cancer J Clin.* 2014;64(2):83-103.
5. Papaemmanuil E, Gerstung M, Bullinger L, et al. Genomic classification and prognosis in acute myeloid leukemia. *N Engl J Med.* 2016;374(23):2209-2221.
6. Burger JA, Landau DA, Taylor-Weiner A, et al. Clonal evolution in patients with chronic lymphocytic leukaemia developing resistance to BTK inhibition. *Nat Commun.* 2016;7(1):11589.
7. Navin N, Kendall J, Troge J, et al. Tumour evolution inferred by single-cell sequencing. *Nature.* 2011;472(7341):90-94.
8. Kim C, Gao R, Sei E, et al. Chemoresistance evolution in triple-negative breast cancer delineated by single-cell sequencing. *Cell.* 2018;173(4):879-893.e13.
9. Roerink SF, Sasaki N, Lee-Six H, et al. Intra-tumour diversification in colorectal cancer at the single-cell level. *Nature.* 2018;556(7702):457-462.
10. Cortes JE, Tallman MS, Schiller GJ, et al. Phase 2b study of 2 dosing regimens of quizartinib monotherapy in FLT3-ITD-mutated, relapsed or refractory AML. *Blood.* 2018;132(6):598-607.
11. Cortes J, Perl AE, Döhner H, et al. Quizartinib, an FLT3 inhibitor, as monotherapy in patients with relapsed or refractory acute myeloid leukaemia: an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol.* 2018;19(7):889-903.
12. Pellegrino M, Sciambi A, Treusch S, et al. High-throughput single-cell DNA sequencing of acute myeloid leukemia tumors with droplet microfluidics. *Genome Res.* 2018;28(9):1345-1352.
13. McMahon CM, Ferng T, Canaani J, et al. Clonal selection with Ras pathway activation mediates secondary clinical resistance to selective FLT3 inhibition in acute myeloid leukemia. *Cancer Discov.* 2019;9(8):1050-1063.
14. Huang BJ, Wandler AM, Meyer LK, et al. Convergent genetic aberrations in murine and human T lineage acute lymphoblastic leukemias. *PLoS Genet.* 2019;15(6):e1008168.
15. Robinson JT, Thorvaldsdóttir H, Winckler W, et al. Integrative genomics viewer. *Nat Biotechnol.* 2011;29:24-26.
16. Zhang H, Savage S, Schultz AR, et al. Clinical resistance to crenolanib in acute myeloid leukemia due to diverse molecular mechanisms. *Nat Commun.* 2019;10(1):244.
17. Aikawa T, Togashi N, Iwanaga K, et al. Quizartinib, a selective FLT3 inhibitor, maintains antileukemic activity in preclinical models of RAS-mediated midostaurin-resistant acute myeloid leukemia cells. *Oncotarget.* 2020;11(11):943-955.
18. Smith CC, Paguirigan A, Jeschke GR, et al. Heterogeneous resistance to quizartinib in acute myeloid leukemia revealed by single-cell analysis. *Blood.* 2017;130(1):48-58.
19. Baker SD, Zimmerman EI, Wang YD, et al. Emergence of polyclonal FLT3 tyrosine kinase domain mutations during sequential therapy with sorafenib and sunitinib in FLT3-ITD-positive acute myeloid leukemia. *Clin Cancer Res.* 2013;19(20):5758-5768.
20. Smith CC, Zhang C, Lin KC, et al. Characterizing and overriding the structural mechanism of the quizartinib-resistant FLT3 "gatekeeper" F691L mutation with PLX3397. *Cancer Discov.* 2015;5(6):668-679.
21. Döhner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood.* 2017;129(4):424-447.