

Genomic analysis of primary and secondary myelofibrosis redefines the prognostic impact of *ASXL1* mutations: a FIM study

Damien Luque Paz,¹⁻³ Jérémie Riou,⁴ Emmanuelle Verger,^{5,6} Bruno Cassinat,^{5,6} Aurélie Chauveau,⁷ Jean-Christophe Ianotto,⁸ Brigitte Dupriez,⁹ Françoise Boyer,¹⁰ Maxime Renard,¹⁻³ Olivier Mansier,^{11,12} Anne Murati,¹³ Jérôme Rey,¹⁴ Gabriel Etienne,¹⁵ Véronique Mansat-De Mas,¹⁶ Suzanne Tavitian,¹⁷ Olivier Nibourel,^{18,19} Stéphane Girault,²⁰ Yannick Le Bris,^{21,22} François Girodon,²³ Dana Ranta,²⁴ Jean-Claude Chomel,²⁵ Pascale Cony-Makhoul,²⁶ Pierre Sujobert,²⁷ Margot Robles,²⁸ Raouf Ben Abdelali,²⁹ Olivier Kosmider,³⁰ Laurane Cottin,¹⁻³ Lydia Roy,^{31,32} Ivan Sloma,^{33,34} Fabienne Vacheret,³⁵ Mathieu Wemeau,³⁶ Pascal Mossuz,³⁷ Borhane Slama,³⁸ Vincent Cussac,³⁹ Guillaume Denis,⁴⁰ Anouk Walter-Petrich,⁴¹ Barbara Burroni,⁴² Nathalie Jézéquel,⁷ Stéphane Giraudier,^{5,6} Eric Lippert,⁷ Gérard Socié,⁴³ Jean-Jacques Kiladjan,^{6,44} and Valérie Ugo,¹⁻³ on behalf of the French Intergroup of Myeloproliferative Neoplasms

¹Univ Angers, INSERM, CRCINA, Angers, France; ²Laboratoire d'Hématologie, Centre Hospitalier Universitaire (CHU) Angers, Angers, France; ³Univ Angers, UFR Santé, Angers, France; ⁴Univ Angers, INSERM, Unit 1066 miNT, Angers, France; ⁵Laboratoire de Biologie Cellulaire, Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Paris, France; ⁶Université de Paris, U1131 INSERM, IRSL, Paris, France; ⁷Laboratoire d'Hématologie and ⁸Service d'Hématologie Clinique, CHRU Brest, Brest, France; ⁹Hématologie Clinique, CH Lens, Lens, France; ¹⁰Service des Maladies du Sang, CHU Angers, Angers, France; ¹¹Laboratoire d'Hématologie, CHU Bordeaux, Bordeaux, France; ¹²Université de Bordeaux, INSERM U1034, Bordeaux, France; ¹³Département de Biopathologie et Département d'Oncologie Prédictive and ¹⁴Département d'Hématologie, Centre de Recherche en Cancérologie de Marseille, Institut Paoli-Calmettes, INSERM, Marseille, France; ¹⁵Département d'Hématologie, Institut Bergonié, Bordeaux, France; ¹⁶Laboratoire d'Hématologie and ¹⁷Service d'Hématologie, CHU Toulouse, Institut Universitaire du Cancer de Toulouse Oncopole, Toulouse, France; ¹⁸Laboratoire d'Hématologie Cellulaire and ¹⁹UMR 9020-UMR-S 1277-Canther-Cancer Heterogeneity, Plasticity and Resistance to Therapies, Institut de Recherche contre le Cancer de Lille, University Lille, CNRS, INSERM, CHU Lille, Lille, France; ²⁰Service d'Hématologie, CHU Limoges, Limoges, France; ²¹Laboratoire d'Hématologie, CHU Nantes, Nantes, France; ²²Université de Nantes, INSERM, CRCINA, Nantes, France; ²³Service d'Hématologie Biologique, CHU Dijon, Dijon, France; ²⁴Hématologie Clinique, CHU Nancy, Nancy, France; ²⁵Service de Cancérologie Biologique, CHU Poitiers, Poitiers, France; ²⁶Hématologie, CH Annecy, Annecy, France; ²⁷Service d'Hématologie Biologique, Hospices Civils de Lyon, Hôpital Lyon Sud, Pierre-Bénite, France; ²⁸Hématologie Clinique, CH Périgueux, Périgueux, France; ²⁹Pôle Hématologie et Oncologie, Laboratoire Cerba, Saint-Ouen L'Aumône, France; ³⁰Laboratoire d'Hématologie, Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Paris, France; ³¹Service d'Hématologie, Assistance Publique-Hôpitaux de Paris, Hôpital Henri Mondor, Créteil, France; ³²Faculté de Santé, Université Paris Est Créteil (UPEC), Créteil, France; ³³Département d'Hématologie et Immunologie, Assistance Publique-Hôpitaux de Paris, Hôpital Henri Mondor, Créteil, France; ³⁴Université Paris Est Créteil, INSERM, IMRB, Créteil, France; ³⁵Service d'Hématologie, CH Perpignan, Perpignan, France; ³⁶Hématologie, CH Roubaix, Roubaix, France; ³⁷Laboratoire d'Hématologie, CHU Grenoble, Grenoble, France; ³⁸Service d'Onco-Hématologie, CH Avignon, Avignon, France; ³⁹Laboratoire d'Hématologie, CH Le Mans, Le Mans, France; ⁴⁰Service d'Hématologie, CH Rochefort, Rochefort, France; ⁴¹ECSTRRA Team U1153, INSERM, Université de Paris, Paris, France; ⁴²Département d'Anatomo-Pathologie, Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Paris, France; ⁴³Hématologie-Transplantation, Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Paris, France; and ⁴⁴Centre d'Investigations Cliniques (INSERM CIC1427), Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Paris, France

Key Points

- Mutations of *TP53* and high-risk genes (*EZH2*, *CBL*, *U2AF1*, *SRSF2*, *IDH1*, *IDH2*, *NRAS* or *KRAS*) are adverse prognostic factors in myelofibrosis.
- *ASXL1* isolated mutations (ie, without *TP53* or high-risk mutations) have no prognostic impact in myelofibrosis.

We aimed to study the prognostic impact of the mutational landscape in primary and secondary myelofibrosis. The study included 479 patients with myelofibrosis recruited from 24 French Intergroup of Myeloproliferative Neoplasms (FIM) centers. The molecular landscape was studied by high-throughput sequencing of 77 genes. A Bayesian network allowed the identification of genomic groups whose prognostic impact was studied in a multistate model considering transitions from the 3 conditions: myelofibrosis, acute leukemia, and death. Results were validated using an independent, previously published cohort (n = 276). Four genomic groups were identified: patients with *TP53* mutation; patients with ≥ 1 mutation in *EZH2*, *CBL*, *U2AF1*, *SRSF2*, *IDH1*, *IDH2*, *NRAS*, or *KRAS* (high-risk group); patients with *ASXL1*-only mutation (ie, no associated mutation in *TP53* or high-risk genes); and other patients. A multistate model found that both *TP53* and high-risk groups were associated with leukemic transformation (hazard ratios [HRs] [95% confidence interval], 8.68 [3.32-22.73] and 3.24 [1.58-6.64], respectively) and death from myelofibrosis

Submitted 18 September 2020; accepted 30 December 2020; published online 5 March 2021. DOI 10.1182/bloodadvances.2020003444.

All data folders and code for figure generation are freely available (<https://github.com/Hematology-Lab-of-Angers-Hospital/FIM-Myelofibrosis-NGS>).

The full-text version of this article contains a data supplement.

© 2021 by The American Society of Hematology

(HRs, 3.03 [1.66-5.56] and 1.77 [1.18-2.67], respectively). *ASXL1*-only mutations had no prognostic value that was confirmed in the validation cohort. However, *ASXL1* mutations conferred a worse prognosis when associated with a mutation in *TP53* or high-risk genes. This study provides a new definition of adverse mutations in myelofibrosis with the addition of *TP53*, *CBL*, *NRAS*, *KRAS*, and *U2AF1* to previously described genes. Furthermore, our results argue that *ASXL1* mutations alone cannot be considered detrimental.

Introduction

The Philadelphia chromosome–negative myeloproliferative neoplasms (MPNs), which include polycythemia vera (PV), essential thrombocythemia (ET), and primary myelofibrosis (PMF), are acquired clonal hematopoietic stem cell disorders characterized by the abnormal proliferation and accumulation of mature blood cells. Myelofibrosis is characterized by dysregulation of the bone marrow stroma with the development of a reticulin fibrosis. PMF differs from post-PV or post-ET secondary myelofibrosis (SMF), which has an evolution rate of ~10% after 10 years of follow-up.¹ The course of myelofibrosis is associated with progressive constitutional symptoms (eg, fatigue, night sweats, and fever), increasing splenomegaly, worsening cytopenia, and a risk of transformation to acute myeloid leukemia (AML). Among MPN, myelofibrosis is associated with the worst prognosis, and its evolution is extremely variable according to prognostic features and treatment.² Risk-adapted therapies range from no treatment to allogeneic hematopoietic stem cell transplantation.²

More than 90% of myelofibrosis cases harbor somatic mutations in the driver genes *JAK2*, *CALR*, or *MPL* that lead to a constitutive activation of the JAK-STAT5 pathway.³ Other somatic nondriver mutations (so-called additional mutations) have been increasingly detected in MPN with the use of high-throughput sequencing.⁴ These additional mutations involve genes with various functions, such as epigenetics, splicing, signalization, and transcription factors, and are also mutated in other myeloid neoplasms like myelodysplastic syndromes and AML. Since the seminal study by Vannucchi et al,⁵ the prognostic role of additional mutations in PMF have been confirmed, and new scoring systems have incorporated “high-risk” genes.^{6,7} Conversely, few data are currently available for SMF. In 2018, Grinfeld et al studied 2035 patients and proposed a prognostic classification for all MPN subtypes, incorporating the mutational landscape to allow personalized prognostic assessment.⁸ Overall, these results pave the way for the use of genomics in the clinical management of MPN, even if additional data are still needed to specify which genes and types of mutation are definitively associated with adverse prognosis.

In this study, we aimed to characterize the molecular landscape of PMF and SMF in a large multicentric cohort to identify homogeneous genomic groups. The prognosis of these genomic groups was then studied, and the results were validated using an external previously published cohort.

Methods

Design and method of the study were performed according the REMARK guidelines (see supplemental Data).⁹

Patients and samples

From 2005, patients with a diagnosis of PMF or SMF (ie, post-ET or PV) were registered in the French Intergroup of Myeloproliferative Neoplasms (FIM) observatory of myelofibrosis, according to the World Health Organization 2008 or 2016 classifications.^{10,11} DNAs from 497 unselected patients with PMF or SMF were centralized in the Angers Hospital for high-throughput molecular analysis. Finally, 479 patients were included in the analysis after exclusion of 10 patients who did not fulfill myelofibrosis diagnostic criteria after central review and 8 patients because of next-generation sequencing (NGS) failure (supplemental Figure 1). All samples were collected at the time of myelofibrosis diagnosis and consisted of whole blood (83%), purified blood granulocytes (14%), or whole bone marrow (3%). The median time between diagnosis and sampling was 17 days (range, 0-88 months). In detail, 77% of samples were collected within 3 months, and 95% of patients were sampled within the first 2 years following diagnosis. As a validation cohort, we used the results of Grinfeld et al on 276 cases of PMF or SMF.⁸ Details regarding patients, diagnostic review, samples, and the validation cohort are provided in supplemental Data.

NGS

A custom RNA-baits panel was designed to cover all exons of 77 genes involved in myeloid malignancies or previously described in MPN. Bioinformatics tools were used to call and annotate variants (details in supplemental Data). Only exonic or splicing mutations (donor and acceptor sites) with a variant allele frequency $\geq 2\%$ and not described as common polymorphisms (ie, $\geq 1\%$ in general population) were retained. Variants were finally classified according to their putative pathogenic effect as pathogenic, likely pathogenic, or variant of unknown significance according to standard guidelines^{12,13} (details in supplemental Data). Furthermore, mutations of unknown significance with a frequency $\geq 0.01\%$ in the general population were considered as rare polymorphisms and were removed.

Statistics

A Bayesian network analysis combined with hierarchical clustering analysis was performed to characterize homogeneous clusters of genes. To describe the natural history of myelofibrosis a multistate model was used, allowing variables associated with each transition to be identified. Correction of *P* values was performed for multiple testing using a Benjamini-Hochberg procedure. Only pathogenic and likely pathogenic mutations were taking into account for statistical analysis of prognosis. Detailed methodology is described in supplemental Data (pages 8-12).

Results

The mutational landscape of myelofibrosis

The whole cohort included 305 PMF and 174 SMF (70 post-PV and 104 post-ET). The characteristics of patients are summarized in supplemental Table 1. Driver mutations were distributed as follows: 309 *JAK2V617F* (65%), 110 *CALR* (23%), 25 *MPLW515* (5%), and 8 patients with a double mutation (4 *JAK2V617F/MPLW515*, 2 *CALR/MPLW515*, 1 *JAK2V617F/CALR*, and 1 *JAK2V617F/JAK2*exon12). For patients with 2 driver mutations, the mutation with the highest allele burden was considered the driver for further analyses. *JAK2V617F* allele burden was higher in SMF than in PMF (supplemental Figure 2). Of note, 27 patients had no driver mutation detected (ie, triple negative).

A total of 1385 additional variants were detected with the following distribution: 861 pathogenic variants, 179 likely pathogenic variants, and 345 variants of unknown significance. We found a median (range) of 3 (0-11) additional variants per patient and 2 (0-10) additional mutations when considering only pathogenic and likely pathogenic variants. The most frequently mutated genes were *ASXL1*, *TET2*, *SRSF2*, *U2AF1* and *EZH2* (Figure 1A; supplemental Figure 3). Mutations in the *SRSF2* and *U2AF1* genes were significantly more frequent in PMF, whereas mutations in *NFE2* were more frequent in SMF (Figure 1B). *SRSF2* and *U2AF1* mutations were more frequently encountered in *JAK2*-mutated than in *CALR*-mutated patients (Figure 1C). Mutations in *ATM*, *KMT2C*, *KMT2D*, and *NOTCH1* were mainly variants of unknown significance. The allele burden distribution of additional mutations harbored different profiles depending on the genes: (1) an allele burden centered on 40% to 50% (eg, *SRSF2* and *U2AF1*); (2) an allele burden with values ranging from 0% to 100% (eg, *EZH2* and *TP53*); and (3) an allele burden with a bimodal distribution, with some values at a low allele burden and others centered on 40% to 50% (eg, *ASXL1* and *TET2*) (Figure 1D). In further analyses, we considered only pathogenic and likely pathogenic mutations. Indeed, the function of “variants of unknown significance” is unclear, and most of these are likely rare polymorphisms. No difference in the number of additional mutations according to the sex was found ($P = .08$), but older patients more frequently had ≥ 1 additional mutation ($P = .0009$; supplemental Figure 4).

We visualized the molecular structure of the cohort using a Bayesian network (Figure 2). Based on this network and on a hierarchical clustering analysis, we defined 7 genomic groups. Since some of these 7 genomic groups harbored similar outcomes, we then merged some of the genomic groups to generate 4 prognostic groups according to their potent respective impact on overall survival (OS) (supplemental Figure 5; Figure 3A). The 4 groups were created in the following sequence: (1) *TP53*-mutated patients ($n = 27$); (2) patients with ≥ 1 mutation in *EZH2*, *CBL*, *U2AF1*, *SRSF2*, *IDH1*, *IDH2*, *NRAS*, or *KRAS* (high-risk group, $n = 150$); (3) *ASXL1*-only mutation (ie, no associated mutation in *TP53* or in high-risk genes, $n = 74$); and (4) “other” patients, with the main mutations being in *NFE2*, *DNMT3A*, *TET2*, and *SF3B1* ($n = 228$) (Figure 2).

Prognostic impact of genomic alterations

The prognostic impact of the 4 molecular groups on OS and cumulative incidence of leukemic transformation highlighted a worst prognostic for *TP53* and high-risk groups (median OS, 20 and 49 months, respectively; median time to leukemic transformation not reached for *TP53* and 110 months for high risk) than in patients with

ASXL1-only and the “other patients” group (median OS, 89 and 116 months, respectively; time to leukemic transformation not reached) (Figure 3; $P < .0001$ for OS and leukemic transformation). This prognostic impact was also found on leukemia-free survival (supplemental Figure 6). Then, we studied the prognostic value of these 4 groups using multistate modeling, which included the following variables: sex; age at diagnosis; hemoglobin, platelet, and leukocyte counts; type of driver mutation; type of myelofibrosis; and the 4-tier genomic classification. Univariate analyses of each transition are summarized in supplemental Table 2, and the final model is summarized in Table 1. *TP53* and high-risk mutations groups were associated with an independent and significantly higher risk of both leukemic transformation (hazard ratio [HR] [95% confidence interval], 8.68 [3.32-22.73], $P < .0001$ and HR, 3.24 [1.58-6.64], $P = .0013$, respectively) and risk of death without AML (HR, 3.03 [1.66-5.56], $P = .0003$ and HR, 1.77 [1.18-2.67], $P = .006$, respectively). Patients with *ASXL1*-only mutation did not have an adverse prognosis either in terms of leukemic transformation (HR, 2.45 [0.95-6.29], $P = .063$) or OS (HR, 1.17 [0.68-2.01], $P = .579$). Moreover, we confirm that well-known factors like age at diagnosis, anemia, and leukocytosis were associated with a higher risk of death.^{2,14} Thrombocytopenia was associated with a higher risk of leukemic transformation, whereas *CALR*-mutated patients presented with a lower risk of AML transformation. Two sensitivity analyses were performed considering either (1) a censor at the time of bone marrow transplantation for the 50 allografted patients and (2) a time 0 as the time of DNA sampling, and both analyses confirmed our findings (supplemental Tables 3 and 4).

Finally, we applied the multistate model to the validation cohort of 276 cases of myelofibrosis published by Grinfeld et al⁸ and found very similar results (supplemental Table 5). In particular the prognostic impact of *TP53* mutations on leukemic transformation and high-risk mutations on leukemic transformation and death were confirmed with HRs close to those found in our cohort. Of note, 2 results cannot be reproduced in the validation cohort: the protective effect of *CALR* mutations on leukemic transformation and reduced OS for *TP53* mutations. However, the absence of prognostic value of the *ASXL1*-only group was also supported by results obtained in the validation cohort.

Deciphering the spectrum of *ASXL1* mutations

We then examined in detail the spectrum of *ASXL1* mutations and found that they were frequently associated with *TP53* or high-risk genes mutations (Figure 1E). Indeed, an *ASXL1* mutation was found in 60% of cases in these 2 genomic groups and was associated with an increased risk of death (HR, 1.5 [1.01-2.29], $P = .044$; supplemental Figure 7).

The allele burden of *ASXL1* mutations was higher when associated with high-risk mutations (supplemental Figure 8). In the *ASXL1*-only group, the allele burden of *ASXL1* mutation was not associated with acute leukemia or death ($P = .293$ and $P = .763$, respectively, with a threshold of 20%). A sensitivity analysis of the multistate model considering only additional mutations with $> 15\%$ allele burden was performed and previous results were confirmed, showing the lack of prognostic significance for *ASXL1*-only mutations (supplemental Table 6).

Comparison of prognostic performances

Finally, we aimed to evaluate the additional value of the genomic groups for prognosis assessment as compared with other prognostic




Figure 1. Mutational landscape of the entire cohort. Proportion of patients with ≥ 1 mutation per gene according to the classification of mutations (A), the type of myelofibrosis (B), and the driver mutation (*JAK2* or *CALR*) (C). The distribution of allelic burden of additional mutations is represented by violin plots (D), and the association between mutations on the genes of the 4-tier classification is shown in a circos plot (E). * $P < .05$; ** $P < .01$; *** $P < .001$. Only pathogenic and likely pathogenic mutations were retained for the circos plot. VAF, variant allele frequency.

classifications. For this purpose, the performance (C-index and time-dependent area under the curve [AUC])^{15,16} and the accuracy (Brier score)¹⁷ to predict death and leukemic transformation were evaluated for standard prognostic scoring systems (ie, International Prognostic Scoring System [IPSS]² for PMF and Myelofibrosis Secondary to PV and ET–Prognostic Model [MYSEC-PM]¹⁸ for SMF), the present 4-tier genomic classification alone, and the combination of our 4-tier genomic with previous scores. The more recent prognostic scores MIPSS70⁶ and MIPSS70+v2⁷ were also evaluated. Results are summarized in Table 2 and supplemental Table 7. The 4-tier genomic classification alone harbored equal or slightly inferior C-index performance as compared with IPSS or MYSEC-PM for death and was superior to IPSS but inferior to MYSEC-PM for leukemic transformation. It is worth noting that the association of genomic groups with standard prognostic scoring systems improved the prediction and accuracy of prognosis for both short-term and long-term events, as illustrated in Figure 4 for death and in supplemental Figure 9 for leukemic transformation. MIPSS70 and MIPSS70+v2 had good predictive values in particular for death in PMF and SMF and for leukemic transformation in PMF. However, the combination

of our 4-tier genomic classification combined with IPSS or MYSEC-PM remained superior to MIPSS70 and MIPSS70+v2.

Discussion

Herein, we analyzed the molecular landscape of a large cohort of patients with PMF and SMF. Gene variants were classified according to their suspected pathogenicity and only pathogenic or likely pathogenic mutations were kept in the statistical analysis for prognosis. The Bayesian network enabled the characterization of the molecular structure of our cohort and it was indeed close to that of the network described by Grinfeld et al.⁸ Both genomic networks reveal *ASXL1* as a central node of the MPN molecular landscape, since most additional mutations arise from this first hit. *ASXL1* mutations were found in approximately one-third of myelofibrosis patients as previously described.^{4,6,19-22} We defined 4 genomic groups, 2 of them, the “*TP53*” and high-risk (ie, ≥ 1 mutation in groups *EZH2*, *CBL*, *U2AF1*, *SRSF2*, *IDH1*, *IDH2*, *NRAS*, or *KRAS*) groups, were associated with an adverse outcome (transition from myelofibrosis to acute leukemia and from myelofibrosis to death). Furthermore, we investigated the




Figure 2. Bayesian network of the molecular landscape. Driver genes are in blue. A green link represents a positive association, and a red one represents 2 mutually exclusive genes. Red, yellow, and blue genes represent the genomic groups identified in the following order: *TP53* mutations, high-risk mutation (without *TP53* mutation), and *ASXL1*-only mutation (ie, without *TP53* or high-risk mutation).

prognostic impact of each gene within the high-risk group by analyzing the performance for the prediction of OS by leaving out the genes one by one. This confirmed that all genes were associated and improved the prognostic evaluation (supplemental Table 8).

Conversely, the *ASXL1*-only group (ie, without mutation in *TP53* or high-risk genes) had no prognostic impact on the risk of death and

had only a moderate impact for leukemic transformation. The multivariate analysis also found that age at diagnosis, leukocytosis, and anemia were associated with a higher risk of death, while thrombocytopenia was associated with a risk of leukemic transformation, in agreement with risk factors included in previous prognosis scoring systems, from the Lille score to IPSS and Dynamic International Prognostic Scoring System-Plus (DIPSS-plus).^{2,14,23}




Figure 3. Prognosis according to the 4-tier genomic classification. Kaplan-Meier curve representing OS (A) or cumulative incidence of leukemic transformation (B). Follow-up was right censored at 72 months for Kaplan-Meier.

Table 1. Multistage model for AML and death events

| Variable | Transitions | | | | | |
|--|----------------------|--------|------------------------|--------|------------------|------|
| | Myelofibrosis to AML | | Myelofibrosis to death | | AML to death | |
| | HR [95% CI] | P | HR [95% CI] | P | HR [95% CI] | P |
| Age at diagnosis | — | — | 1.05 [1.03-1.07] | <.0001 | — | — |
| Sex (male) | — | — | 1.41 [0.96-2.06] | .077 | — | — |
| Hemoglobin | — | — | 0.81 [0.74-0.89] | <.0001 | — | — |
| Leukocyte count | — | — | 1.04 [1.02-1.05] | <.0001 | — | — |
| Platelet count | 0.99 [0.99-0.99] | .012 | — | — | — | — |
| Driver mutation (<i>JAK2</i> as reference): <i>CALR</i> | 0.21 [0.06-0.70] | .011 | — | — | — | — |
| Genomic groups (others as reference) | | | | | | |
| <i>TP53</i> | 8.68 [3.32-22.73] | <.0001 | 3.03 [1.66-5.56] | .0003 | 1.21 [0.31-4.72] | .784 |
| High risk | 3.24 [1.58-6.64] | .0013 | 1.77 [1.18-2.67] | .006 | 0.88 [0.33-2.36] | .801 |
| <i>ASXL1</i> only | 2.45 [0.95-6.29] | .063 | 1.17 [0.68-2.01] | .579 | 1.22 [0.33-4.47] | .767 |

CI, confidence interval.

We also found that *CALR*-mutated patients had a lower risk of leukemic transformation, as previously described.²⁴ Of note, this prognostic advantage was reported as being limited to type 1 and type 1-like mutations of *CALR*.^{25,26} In our cohort, 79% of *CALR* mutations were type 1/type 1-like, and no distinction was made in multivariate analysis, because no difference was found in

univariate analysis between type 1/type 1-like and other *CALR* mutations ($P = .77$ and $.6$ for leukemia-free survival and OS, respectively). The 2 major differences of the multistage model in the validation cohort concerned the absence of significance of *CALR* mutations for leukemic transformation and the *TP53* genomic group for OS and may be explained by the limited number of patients with

Table 2. Prognostic performance comparison

| | C-index | Events at 24 mo | | Events at 48 mo | | Events at 72 mo | |
|--------------------------------|-------------|-----------------|-----------|-----------------|-----------|-----------------|-----------|
| | | Brier score | AUC | Brier score | AUC | Brier score | AUC |
| OS | | | | | | | |
| PMF (n = 305) | | | | | | | |
| NGS | 0.66 | 0.052 | 66 | 0.101 | 72 | 0.13 | 73 |
| IPSS | 0.68 | 0.051 | 71 | 0.098 | 72 | 0.129 | 74 |
| MIPSS70 | 0.67 | 0.05 | 70 | 0.095 | 73 | 0.122 | 71 |
| NGS-IPSS | 0.73 | 0.05 | 74 | 0.093 | 79 | 0.118 | 82 |
| SMF (n = 174) | | | | | | | |
| NGS | 0.66 | 0.057 | 68 | 0.117 | 61 | 0.147 | 68 |
| MYSEC-PM | 0.71 | 0.061 | 74 | 0.11 | 76 | 0.133 | 80 |
| MIPSS70* | 0.68 | 0.06 | 73 | 0.111 | 71 | 0.135 | 72 |
| NGS-MYSEC-PM | 0.77 | 0.054 | 82 | 0.102 | 79 | 0.124 | 86 |
| Leukemic transformation | | | | | | | |
| PMF (n = 305) | | | | | | | |
| NGS | 0.71 | 0.017 | 76 | 0.041 | 74 | 0.058 | 79 |
| IPSS | 0.59 | 0.017 | 60 | 0.04 | 61 | 0.062 | 61 |
| MIPSS70 | 0.72 | 0.017 | 76 | 0.038 | 78 | 0.056 | 73 |
| NGS-IPSS | 0.73 | 0.016 | 78 | 0.039 | 77 | 0.056 | 79 |
| SMF (n = 174) | | | | | | | |
| NGS | 0.67 | 0.021 | 70 | 0.047 | 65 | 0.062 | 72 |
| MYSEC-PM | 0.75 | 0.022 | 76 | 0.046 | 82 | 0.059 | 85 |
| MIPSS70* | 0.66 | 0.022 | 64 | 0.048 | 71 | 0.062 | 73 |
| NGS-MYSEC-PM | 0.76 | 0.02 | 81 | 0.042 | 83 | 0.054 | 87 |

Comparisons were performed between classical prognosis score systems and the 4-tier genomic classification (NGS) alone or in combination with a clinical score. C-index and AUC evaluated the performance of the model, and Brier score reflected the accuracy of the prediction (ie, the rate of error). Bold indicates the best values.

*MIPSS70 was developed for PMF.




Figure 4. Prognosis performance comparison. The performance (time-AUC; A and C) and accuracy (Brier score; B and D) for death prediction were measured over the time for usual prognosis scoring systems, the 4-tier genomic classification (NGS) alone, or in combination for primary (A-B) and secondary (C-D) myelofibrosis patients.

either *CALR* mutation (n = 46) and *TP53* mutations, with 11 mutated cases, of which 5 progressed to acute leukemia and are therefore not considered for myelofibrosis to death transition in the multistate model.

In the present study, we identified mutations in *TP53*, *EZH2*, *SRSF2*, *IDH1*, *IDH2*, *U2AF1*, *CBL*, *NRAS*, and *KRAS* genes as adverse prognostic factors in myelofibrosis. The detrimental role of *TP53* mutations has been previously described in cohorts that included all subtypes of MPN, and the role of *TP53* in leukemic transformation has been shown in mice models.^{4,8,27,28} Among the other genes, we found a worse prognosis associated with *EZH2*, *SRSF2*, *IDH1*, and *IDH2*, as previously reported by others for high-risk profiles.⁵ *U2AF1* and *N/KRAS* have also been described as adverse prognostic factors.^{7,29,30} *CBL* protein is a RAS pathway regulator, and previous studies have reported a reduced OS in MPN patients with *CBL* mutations.^{22,30,31}

Our results about the prognostic value of *ASXL1* mutations seem, at a first glance, to contrast those of previous studies in PMF, where they were classified as high molecular risk (HMR).^{5,6,22} Nevertheless, some other studies found no prognostic impact of *ASXL1* mutations in SMF,³² and in ET and in PV in MIPSS-ET and MIPSS-PV did not include *ASXL1* mutations.³³ Santos et al have recently studied a cohort of 723 patients with myelofibrosis and found a prognostic value of *N/KRAS* mutations but no impact of HMR mutations using a multivariate model considering sex, platelet count, transfusion dependency, DIPSS classification, karyotype, driver mutation, HMR (ie, *ASXL1*, *EZH2*, and *IDH1/2*), and *N/KRAS*

mutations.³⁰ In this later study, there was a positive association between *N/KRAS* mutations and *ASXL1* mutations. Similarly, we found that 59% of patients with *ASXL1* mutations also had either a mutation in *TP53* or in a gene from the high-risk genomic group (*EZH2*, *CBL*, *U2AF1*, *SRSF2*, *IDH1*, *IDH2*, *NRAS*, or *KRAS*). The selection of genomic groups based on the Bayesian network allowed us to study the prognostic role of isolated *ASXL1* mutations. Thus, we hypothesize that the apparent prognostic impact of *ASXL1* mutations found in previous studies could be due to either (1) other mutations of adverse prognostic not detected or not included in the NGS panels or in statistical models and frequently associated with *ASXL1* or (2) the additional impact of *ASXL1* mutations in patients harboring other high-risk mutations. Our results have been validated on an independent cohort. Thus, it is unlikely that recruitment bias explains differences about the impact of *ASXL1* mutations.

Our sequencing technology allowed the detection of very low allelic burdens (down to 2%). Such *ASXL1* mutations with low allelic burden might simply reveal age-related small clones not linked to the MPN (ie, clonal hematopoiesis of indeterminate potential),³⁴ but a sensitivity analysis showed similar results when considering only mutations with an allelic burden >15%. The type of *ASXL1* mutation may also be important. Indeed, Tefferi et al demonstrated a similar prognosis for frameshift, nonsense, and missense mutations, whereas Kuykendall et al found no apparent impact of missense mutations.^{35,36} In our cohort, all *ASXL1* mutations harbored a truncated C-terminal domain.

Finally, in this study, we have considered the c.1934dupG mutation of *ASXL1*, whose status as a true mutation or artifact has previously been debated, but it is now considered a bona fide mutation (with 472 occurrences in COSMIC).^{37,38} A previous study found a worse prognosis for this particular mutation in PMF.³⁹ This led us to conduct 2 sensitivity analyses, one excluding the *ASXL1* c.1934dupG mutation and the second dissociating this mutation from the other *ASXL1* mutations, and we found no prognosis impact of *ASXL1* mutations for either analysis (supplemental Tables 9 and 10).

The role of *ASXL1* mutations in myeloid neoplasms development is still unclear. *ASXL1* mutations result in a loss of PRC2-mediated H3K27 trimethylation.⁴⁰ Deletion of *Asx1* in mice induced a myelodysplasia phenotype, and a knockin expression of an *Asx1* mutation with protein truncation led to an age-dependent myeloid skewing with anemia, thrombocytosis, and dysplasia.^{41,42} Furthermore, mice models have shown a cooperation between *Asx1* and other mutations (eg, *Nras* and *Runx1*) in promoting leukemic transformation.^{40,42} These findings fit with our results showing that *ASXL1* mutations had no prognostic impact alone but seemed to confer a worse prognosis when associated with high-risk mutations.

For the evaluation of the prognostic performance, we compiled (1) C-index reflecting the concordance probability of the prognostic model, (2) AUC for different times of follow-up (time-AUC) reflecting the performance of prediction, and (3) Brier score for different times of follow-up reflecting the accuracy of the prediction. All these indicators showed that our 4 genomic group classification allowed a good prognostic performance equivalent to most previous scores.^{2,6,7,18} Furthermore, the prognostic prediction for both death and leukemic transformation were highly improved by combination of the 4 genomic groups with standard scores.

In conclusion, our results argue for a revision of the prognostic classification of somatic mutations in myelofibrosis with the exclusion of *ASXL1* and the inclusion of the *TP53*, *U2AF1*, *CBL*, *NRAS*, and *KRAS* genes to the already-recognized *SRSF2*, *EZH2*, *IDH1*, and *IDH2* HMR mutations. These results provide a basis for a new molecular classification that will need to be validated in large multicentric cohorts. Indeed, the accurate definition of a high-risk molecular signature is crucial, as it could influence patient management and therapy.^{43,44}

Acknowledgments

The authors are indebted to Rébecca Jouanneau-Courville, who performed sequencing experiments, and Annaëlle Beucher (Angers) and Arnaud Guille (Marseille), who helped with the bioinformatics. They thank the patients, biologists, pathologists, hematologists, and data managers who participated in the FIMBANK project. They especially thank Jean-Marie Chrétien (Délégation à la Recherche Clinique et Innovation, CHU Angers) and Matthieu Resche-Rigon (Université de Paris, ECSTRRA Team U1153, INSERM) for electronic case report form and data coordination and Célia Belkhodja, Kahina Oukherfellah, Eléonore Ourouda-Mbaya, Oryanne Wojtowicz, Alicia Delisle, and Mathilde Favreau for many years of biological and clinical data collection in FIM and FIMBANK electronic case report forms. The authors also thank all participating biobanks (Centres de Ressources Biologiques) for providing high-quality DNA samples, the Plateau de Biologie et de Médecine Moléculaire and the Génomique Onco-Hématologique platform (CHU Angers) for sequencing facilities, and the team of Yves Delneste (Université d'Angers, INSERM, CRCINA, Angers) for scientific discussions.

They thank the French Innovative Leukemia Organization, especially Valérie Rolland-Neyret, for continuous support to the FIM. Finally, they thank Moritz Gerstung and Peter Campbell's groups for sharing their cohort's data via the MPN-multistage tool.

This study was supported by grants from the Force Hemato foundation (Fonds De Recherche Clinique en Hématologie) (AAP-FH-2016) and the Association Laurette Fugain (ALF 2018/12). This work is also a deliverable of the FIMBANK network, which was founded by the French Institut National du Cancer (INCa BCB 2013).

Authorship

Contribution: D.L.P. and V.U. designed the study; E.V., B.C., A.C., O.M., V.M.-D.M., Y.L.B., F.G., J.-C.C., A.M., P.S., R.B.A., O.K., I.S., O.N., P.M., V.C., and E.L. provided DNA samples; J.-C.J., B.D., F.B., J. Rey, G.E., S.T., F.V., S. Girault, F.G., D.R., P.C.-M., M. Robles, L.R., M.W., B.S., G.D., S. Giraudier, E.L., and J.-J.K. cared for patients; M. Renard developed the bioinformatics tools; D.L.P., E.V., B.C., A.C., O.M., A.M., L.C., I.S., and E.L. analyzed the NGS; B.B. coordinated the histological evaluation; A.W.-P. coordinated the maintenance of the clinical database; N.J. and V.U. coordinated the FIMBANK network; D.L.P. and J. Riou performed statistical analysis; and D.L.P., J. Riou, V.U., G.S., and J.-J.K. wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: D.L.P., 0000-0002-4523-675X; J. Riou, 0000-0002-7056-9257; E.V., 0000-0002-2773-2784; B.C., 0000-0002-6514-3905; O.M., 0000-0002-7943-8800; G.E., 0000-0001-7600-4954; F.G., 0000-0003-3151-1068; J.-C.C., 0000-0001-9227-1104; O.K., 0000-0002-6021-4057; L.C., 0000-0001-7933-8626; G.D., 0000-0003-0434-265X; S. Giraudier, 0000-0002-1817-2236; E.L., 0000-0002-2312-6584; V.U., 0000-0002-4678-2138.

Correspondence: Valérie Ugo, Laboratoire d'Hématologie, Institut de Biologie en Santé, CHU Angers, 4 Rue Larrey, 49933 Angers Cedex 9, France; e-mail: valerie.ugo@chu-angers.fr; and Damien Luque Paz, Laboratoire d'Hématologie, Institut de Biologie en Santé, CHU Angers, 4 Rue Larrey, 49933 Angers Cedex 9, France; e-mail: damien.luquepaz@chu-angers.fr.

A list of the members of the FIM appears in "Appendix."

Appendix: study group members

Additional members of the FIM include: Fiorenza Barraco (Lyon, France), Yan Beauverd (Geneva, Switzerland), Annaëlle Beucher (Angers, France), Audrey Bidet (Bordeaux, France), Chrystèle Bilhou-Nabera (Paris, France), Odile Blanchet (Angers, France), Anne Bouvier (Angers, France), Marie-Madeleine Coudé (Saint-Ouen L'Aumône, France), Alexandre De Brevern (Paris, France), François Delhommeau (Paris, France), Bernard Drénou (Mulhouse, France), Franck Geneviève (Angers, France), Alban Godon (Angers, France), Alexandre Guy (Bordeaux, France), Pierre Hirsh (Paris, France), Mathilde Hunault (Angers, France), Pascal Hutin (Quimper, France), Norbert Ifrah (Angers, France), Chloé James (Bordeaux, France), Marie-Caroline Le Bousse Kerdiles (Villejuif, France), Benjamin Lebecque (Clermont-Ferrand, France), Laurence Legros (Villejuif, France), Nabih Maslah (Paris, France), Mathieu Meunier (Grenoble, France), Dina Naguib (Caen, France), Corentin Orvain (Angers, France), Florence Pasquier (Villejuif, France), Isabelle

Plo (Villejuif, France), Aline Praire (Lyon, France), Emmanuel Raffoux (Paris, France), Bénédicte Ribourtout (Angers, France), Alberto Santagostino (Troyes, France), Isabelle Sudaka-Sammarcelli (Nice,

France), Marie-Hélène Schlagetter (Paris, France), Aline Schmidt (Angers, France), Sylvain Thépot (Angers, France), Marie Tuffigo (Angers, France), and Laura Velazquez (Villejuif, France).

References

1. Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood*. 2014;124(16):2507-2513, quiz 2615.
2. Cervantes F, Dupriez B, Pereira A, et al. New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood*. 2009;113(13):2895-2901.
3. Zoi K, Cross NCP. Genomics of myeloproliferative neoplasms. *J Clin Oncol*. 2017;35(9):947-954.
4. Lundberg P, Karow A, Nienhold R, et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood*. 2014;123(14):2220-2228.
5. Vannucchi AM, Lasho TL, Guglielmelli P, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia*. 2013;27(9):1861-1869.
6. Guglielmelli P, Lasho TL, Rotunno G, et al. MIPSS70: mutation-enhanced international prognostic score system for transplantation-age patients with primary myelofibrosis. *J Clin Oncol*. 2018;36(4):310-318.
7. Tefferi A, Guglielmelli P, Lasho TL, et al. MIPSS70+ version 2.0: mutation and karyotype-enhanced international prognostic scoring system for primary myelofibrosis. *J Clin Oncol*. 2018;36(17):1769-1770.
8. Grinfeld J, Nangalia J, Baxter EJ, et al. Classification and personalized prognosis in myeloproliferative neoplasms. *N Engl J Med*. 2018;379(15):1416-1430.
9. McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM; Statistics Subcommittee of the NCI-EORTC Working Group on Cancer Diagnostics. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol*. 2005;23(36):9067-9072.
10. Vardiman JW, Thiele J, Arber DA, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood*. 2009;114(5):937-951.
11. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia [published correction appears in *Blood*. 2016;128(3):462-463]. *Blood*. 2016;127(20):2391-2405.
12. Richards S, Aziz N, Bale S, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
13. Li MM, Datto M, Duncavage EJ, et al. Standards and guidelines for the interpretation and reporting of sequence variants in cancer: a joint consensus recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. *J Mol Diagn*. 2017;19(1):4-23.
14. Dupriez B, Morel P, Demory JL, et al. Prognostic factors in agnogenic myeloid metaplasia: a report on 195 cases with a new scoring system. *Blood*. 1996;88(3):1013-1018.
15. Harrel FE Jr., Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med*. 1996;15(4):361-387.
16. Blanche P, Dartigues J-F, Jacqmin-Gadda H. Estimating and comparing time-dependent areas under receiver operating characteristic curves for censored event times with competing risks. *Stat Med*. 2013;32(30):5381-5397.
17. van de Wiel MA, Berkhof J, van Wieringen WN. Testing the prediction error difference between 2 predictors. *Biostatistics*. 2009;10(3):550-560.
18. Passamonti F, Giorgino T, Mora B, et al. A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. *Leukemia*. 2017;31(12):2726-2731.
19. Brecqueville M, Rey J, Devillier R, et al. Array comparative genomic hybridization and sequencing of 23 genes in 80 patients with myelofibrosis at chronic or acute phase. *Haematologica*. 2014;99(1):37-45.
20. Delic S, Rose D, Kern W, et al. Application of an NGS-based 28-gene panel in myeloproliferative neoplasms reveals distinct mutation patterns in essential thrombocythaemia, primary myelofibrosis and polycythaemia vera. *Br J Haematol*. 2016;175(3):419-426.
21. Gill H, Ip H-W, Yim R, et al. Next-generation sequencing with a 54-gene panel identified unique mutational profile and prognostic markers in Chinese patients with myelofibrosis. *Ann Hematol*. 2019;98(4):869-879.
22. Tefferi A, Lasho TL, Finke CM, et al. Targeted deep sequencing in primary myelofibrosis. *Blood Adv*. 2016;1(2):105-111.
23. Gangat N, Caramazza D, Vaidya R, et al. DIPSS plus: a refined Dynamic International Prognostic Scoring System for primary myelofibrosis that incorporates prognostic information from karyotype, platelet count, and transfusion status. *J Clin Oncol*. 2011;29(4):392-397.
24. Alvarez-Larrán A, Senin A, Fernández-Rodríguez C, et al. Impact of genotype on leukaemic transformation in polycythaemia vera and essential thrombocythaemia. *Br J Haematol*. 2017;178(5):764-771.
25. Tefferi A, Lasho TL, Finke C, et al. Type 1 vs type 2 calreticulin mutations in primary myelofibrosis: differences in phenotype and prognostic impact. *Leukemia*. 2014;28(7):1568-1570.

26. Tefferi A, Lasho TL, Tischer A, et al. The prognostic advantage of calreticulin mutations in myelofibrosis might be confined to type 1 or type 1-like CALR variants. *Blood*. 2014;124(15):2465-2466.
27. Harutyunyan A, Klampfl T, Cazzola M, Kralovics R. p53 lesions in leukemic transformation. *N Engl J Med*. 2011;364(5):488-490.
28. Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic transformation of myeloproliferative neoplasms. *Proc Natl Acad Sci USA*. 2014;111(50):E5401-E5410.
29. Tefferi A, Finke CM, Lasho TL, et al. U2AF1 mutation types in primary myelofibrosis: phenotypic and prognostic distinctions. *Leukemia*. 2018;32(10):2274-2278.
30. Santos FPS, Getta B, Masarova L, et al. Prognostic impact of RAS-pathway mutations in patients with myelofibrosis. *Leukemia*. 2020;34(3):799-810.
31. Grand FH, Hidalgo-Curtis CE, Ernst T, et al. Frequent CBL mutations associated with 11q acquired uniparental disomy in myeloproliferative neoplasms. *Blood*. 2009;113(24):6182-6192.
32. Rotunno G, Pacilli A, Artusi V, et al. Epidemiology and clinical relevance of mutations in postpolycythemia vera and postessential thrombocythemia myelofibrosis: a study on 359 patients of the AGIMM group. *Am J Hematol*. 2016;91(7):681-686.
33. Tefferi A, Guglielmelli P, Lasho TL, et al. Mutation-enhanced international prognostic systems for essential thrombocythemia and polycythemia vera. *Br J Haematol*. 2020;189(2):291-302.
34. Watson CJ, Papula AL, Poon GYP, et al. The evolutionary dynamics and fitness landscape of clonal hematopoiesis. *Science*. 2020;367(6485):1449-1454.
35. Tefferi A, Lasho TL, Finke C, et al. Prognostic significance of ASXL1 mutation types and allele burden in myelofibrosis. *Leukemia*. 2018;32(3):837-839.
36. Kuykendall A, Kaplan J, Talati C, et al. Gaining a better "sense" for ASXL1 mutations in myelofibrosis: assessing the variable impact of frameshift, nonsense, and missense mutations. *Blood*. 2017;130(suppl 1):2939.
37. Gelsi-Boyer V, Trouplin V, Adélaïde J, et al. Mutations of polycomb-associated gene *ASXL1* in myelodysplastic syndromes and chronic myelomonocytic leukaemia. *Br J Haematol*. 2009;145(6):788-800.
38. Abdel-Wahab O, Kilpivaara O, Patel J, Busque L, Levine RL. The most commonly reported variant in *ASXL1* (c.1934dupG;p.Gly646TrpfsX12) is not a somatic alteration. *Leukemia*. 2010;24(9):1656-1657.
39. Rotunno G, Mannarelli C, Brogi G, et al. Spectrum of *ASXL1* mutations in primary myelofibrosis: prognostic impact of the *ASXL1* p.G646Wfs*12 mutation. *Blood*. 2019;133(26):2802-2808.
40. Abdel-Wahab O, Adli M, LaFave LM, et al. *ASXL1* mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell*. 2012;22(2):180-193.
41. Abdel-Wahab O, Gao J, Adli M, et al. Deletion of *Asxl1* results in myelodysplasia and severe developmental defects in vivo. *J Exp Med*. 2013;210(12):2641-2659.
42. Nagase R, Inoue D, Pastore A, et al. Expression of mutant *Asxl1* perturbs hematopoiesis and promotes susceptibility to leukemic transformation. *J Exp Med*. 2018;215(6):1729-1747.
43. Tefferi A, Guglielmelli P, Nicolosi M, et al. GIPSS: genetically inspired prognostic scoring system for primary myelofibrosis. *Leukemia*. 2018;32(7):1631-1642.
44. Barbui T, Tefferi A, Vannucchi AM, et al. Philadelphia chromosome-negative classical myeloproliferative neoplasms: revised management recommendations from European LeukemiaNet. *Leukemia*. 2018;32(5):1057-1069.