

TO THE EDITOR:

Low rate of subsequent malignant neoplasms after CD19 CAR T-cell therapy

Emily M. Hsieh,¹ Regina M. Myers,² Bonnie Yates,³ Colleen Annesley,⁴ Samuel John,⁵ Agne Taraseviciute,¹ Seth M. Steinberg,⁵ Jennifer Sheppard,⁶ Perry Chung,² Lee Chen,¹ Daniel W. Lee,^{3,7} Amanda DiNofia,² Stephan A. Grupp,² Michael R. Verneris,⁸ Theodore W. Laetsch,^{2,6} Deepa Bhojwani,¹ Patrick A. Brown,⁹ Michael A. Pulsipher,^{1,10} Susan R. Rheingold,² Rebecca A. Gardner,⁴ Lia Gore,⁸ Nirali N. Shah,³ and Adam J. Lambie⁴

¹Division of Hematology/Oncology/Transplantation and Cellular Therapy, Children's Hospital Los Angeles Cancer and Blood Disease Institute, Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ²Division of Oncology, Cell Therapy and Transplant Section, Children's Hospital of Philadelphia, Philadelphia, PA; ³Pediatric Oncology Branch, National Cancer Institute/Center for Cancer Research, National Institutes of Health, Bethesda, MD; ⁴Division of Hematology/Oncology, University of Washington, Seattle Children's Hospital, Seattle, WA; ⁵Biostatistics and Data Management Section, National Cancer Institute, National Institutes of Health, Bethesda, MD; ⁶Division of Pediatric Hematology/Oncology, Simmons Comprehensive Cancer Center, UT Southwestern Medical Center, Dallas, TX; ⁷Division of Pediatric Hematology/Oncology, Department of Pediatrics, University of Virginia, Charlottesville, VA; ⁸Pediatric Hematology/Oncology/BMT-CT, University of Colorado, Children's Hospital Colorado, Aurora, CO; ⁹Division of Pediatric Oncology, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD; and ¹⁰Division of Hematology and Oncology, Intermountain Primary Children's Hospital, Huntsman Cancer Institute, Spencer Fox Eccles School of Medicine, University of Utah, Salt Lake City, UT

Subsequent malignant neoplasms (SMNs) are well described after chemotherapy, radiation therapy, and hematopoietic stem cell transplantation (HSCT).¹⁻³ Cancer survivors, particularly those with cancer predisposition syndromes, are also at risk of second primary malignancies unrelated to their treatment.^{4,5} The cumulative incidence of SMNs continues to increase as more childhood cancer survivors reach adulthood.² Among survivors of childhood acute lymphoblastic leukemia (ALL), those with relapsed disease and/or those requiring HSCT have the highest 20-year cumulative incidence of SMNs at ~4.2%.⁶ There is a paucity of literature regarding how cellular immune-based therapies influence the risk of SMNs.

Chimeric antigen receptor (CAR) T-cell therapy is associated with well-described unique inflammatory toxicities in the early postinfusion period, but less is known about potential long-term toxicities. To monitor for delayed adverse events related to lentiviral/retroviral-based technologies used in CAR T-cell manufacturing, the US Food and Drug Administration requires 15 years of long-term follow-up after administration of any investigational human gene therapy product.⁷ This surveillance includes monitoring for risks associated with vector integration, which may disrupt tumor suppressor genes or activate proto-oncogenes near integration sites, thereby theoretically increasing the risk of SMNs. Although malignant transformation directly related to traditional CAR T-cell vector transduction has not been reported,^{8,9} long-term follow-up remains limited. The issue of delayed adverse events is particularly relevant in children and young adults treated with CAR T cells, given their longer potential lifespans and thus higher cumulative risk of SMNs. After a decade since the first child with B-cell ALL (B-ALL) was treated with CD19 CAR T cells, and with longitudinal data increasingly available, we sought to identify the incidence of SMNs in a large cohort of children and young adults after treatment with CD19 CAR T cells.

Based on a multicenter retrospective analysis of children and young adults with B-ALL who received CD19 CAR T cells between 2012 and 2019,¹⁰ we identified an SMN in 7 (1.7%) of 420 patients. SMN diagnoses included cholangiocarcinoma,¹¹ synovial sarcoma, malignant melanoma, papillary thyroid carcinoma, and 3 myeloid neoplasms (Table 1). Median times to SMN diagnosis after initial B-ALL diagnosis and after CD19 CAR T-cell infusion were 8.2 years (range, 1.1-12.5 years) and 3.2 years (range, 0.6-8.2 years), respectively, leading to a 5-year cumulative incidence of SMNs of 1.5% (95% confidence interval, 0.4% to 4%). Six had undergone allogeneic HSCT before diagnosis of the SMN, and 6 had received

Submitted 13 May 2022; accepted 5 July 2022; prepublished online on *Blood Advances* First Edition 14 July 2022; final version published online 7 September 2022. DOI 10.1182/bloodadvances.2022008093.

Contact the corresponding author for data sharing: adam.lambie@seattlechildrens.org.

The current affiliation for A.T. is Janssen Research & Development, LLC, Raritan, NJ. The current affiliation for P.A.B. is Bristol Myers Squibb, New York, NY.

Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

Table 1. Overview of subsequent malignant neoplasms observed post-CD19 CAR T-cell therapy

	Hematologic malignancy (n = 3)	Solid tumor (n = 4)
Diagnosis	AML (n = 2) MDS (n = 1)	Cholangiocarcinoma Synovial sarcoma Melanoma Thyroid carcinoma
Median (range) time from CAR T-cell infusion to diagnosis of SMN, y	1.2 (0.6-1.6)	5.9 (3.2-8.2)
Median (range) time from initial diagnosis to diagnosis of SMN, y	3.7 (1.1-4.2)	9.6 (8.2-12.5)
History of alloHSCT before SMN	Yes (n = 2)	Yes (n = 4)
Current status	2 patients remain alive in remission after alloHSCT; 1 died of progressive disease	3 patients remain alive in remission; 1 died of surgical complications

AML, acute myeloid leukemia; alloHSCT, allogeneic HSCT; MDS, myelodysplastic syndrome.

prior radiation therapy (total-body irradiation [TBI], n = 5; central nervous system irradiation, n = 1). Additionally, 1 patient received ongoing ALL-directed therapy for relapse after CD19 CAR T-cell therapy, before SMN diagnosis (synovial sarcoma), and 1 patient with a myeloid SMN had a germ line *TP53* mutation, a well-described cancer predisposition syndrome. Among those with SMNs, 2 patients received a CD19/28 ζ -based CD19 CAR T-cell therapy; 4 received either commercial tisagenlecleucel or the investigational CD19/41BB predecessor, and 1 received an alternative CD19/BB CAR T-cell therapy. No patient with testing had evidence of replication-competent retrovirus/lentivirus, but assessment of vector integration into SMN sites was limited. With a median follow-up of 1177 days (range, 220-2981 days) after CD19 CAR T-cell therapy, 5 (71%) of 7 patients remain alive and in remission from their SMNs, including 2 of 3 patients with myeloid SMNs. Additional details on the presentation, diagnosis, and management are listed in Tables 2 and 3. Institutional review board approval was obtained at all participating institutions before retrospective analysis. The study was conducted according to the Declaration of Helsinki.

In the limited literature available, the incidence of SMNs in patients who received CD19 CAR T cells ranges from 2.2% to 16.3% and varies between pediatric and adult reports.¹²⁻¹⁴ To that end, the incidence of SMNs in our cohort was similar to that in the pooled safety analysis of tisagenlecleucel in children and young adults with B-ALL.¹² Three (2.2%) of 137 patients included in the tisagenlecleucel safety analysis developed an SMN, with a median follow-up of 24 months, including 2 patients who developed myelodysplastic syndrome and 1 patient with a germ line *TP53* mutation who developed glioblastoma. In our cohort, a trend toward a longer latency period for developing a solid tumor was apparent when compared with developing a hematologic malignancy. This is consistent with historical reports of SMNs after HSCT, where the risk of hematologic malignancies is highest in the first 10 years, but the risk continues to increase, including long latency periods (≥ 10 years) before development

of solid tumors.^{3,15} Notably, 6 (85.7%) of 7 patients in our cohort had received prior radiation therapy, and 6 had undergone prior allogeneic HSCT, including all 4 patients who developed a solid tumor, with a majority receiving a TBI-based myeloablative regimen, a standard approach in B-ALL conditioning.^{16,17}

A rare but important phenomenon after CD19 CAR T-cell therapy for B-ALL is a lineage switch (LS), in which there is an immunophenotypic transition from lymphoid- to myeloid-defining cell surface markers. Based on a recent effort from our group, LSs comprised 7.2% of all relapse phenotypes after CD19 CAR T-cell therapy.¹⁸ It is important to distinguish an LS from a therapy-related myeloid neoplasm, because the prognosis of patients who develop an LS after CAR T-cell therapy is dismal, as recently reported.¹⁸ In contrast, 2 of the 3 patients who developed myeloid SMNs in this report received successful salvage therapy. Immunophenotype alone is insufficient to distinguish between the 2 entities, and cytogenetic changes observed at the time of the event are critical, because the original cytogenetic drivers are likely to be preserved in an LS. All 3 myeloid SMNs in our report were differentiated from LSs based on the absence of the cytogenetic abnormalities noted in the B-ALL clone, including 1 SMN emerging from the prior sex-disparate stem cell source. Given the clinical relevance of separating LSs from SMNs, use of next-generation sequencing of B-cell receptor rearrangements for minimal residual disease tracking in B-ALL^{19,20} and genomic analyses with either DNA- or RNA-based deep sequencing²¹ should be considered when distinguishing between LSs and myeloid SMNs.

With the low incidence of SMNs in our series and in the context of extensive prior ALL-directed therapy, including chemotherapy, radiation therapy, and/or HSCT, direct attribution of SMNs to CAR T-cell therapy is challenging, particularly in those undergoing HSCT after CD19 CAR T-cell therapy. However, given our findings and the absence of reported events related directly to CD19 CAR T-cell vector integration, current surveillance for SMNs, which incorporates both the US Food and Drug Administration long-term follow-up guidelines and routine postcancer long-term monitoring (including heightened monitoring in those with cancer predisposition syndromes),⁵ is likely sufficient at this time.

Although the rate of SMNs in this cohort is low, ongoing surveillance over decades will be required to understand the full risk of subsequent neoplasms. Additionally, the risk of SMNs beyond CD19 CAR T-cell targeting, within other diseases, and with next-generation constructs is unknown and will be informed by additional studies. Indeed, 2 recently published reports detail the development of T-cell lymphomas derived from CD19 CAR T cells that were generated using the *piggyBac* transposon system,^{22,23} a delivery system developed to decrease the cost, logistics, and theoretic risk of insertional mutagenesis associated with viral vectors.

Demonstrating a low incidence of SMNs, this report adds to the limited literature surrounding delayed adverse events after CD19 CAR T-cell therapy and helps inform patient counseling and surveillance approaches. Monitoring for delayed toxicities is particularly relevant in children as CAR T-cell therapy continues to advance and moves earlier into frontline treatment for high-risk ALL and potentially spares the need for more intensive therapies that increase SMN risk. Although the extensive prior therapy received in our cohort,

Table 2. Presentation, diagnosis, and management of solid tumor SMNs

	Cholangiocarcinoma ¹¹	Synovial sarcoma	Melanoma	Thyroid carcinoma
Oncologic history	History of multiply relapsed B-ALL, initially diagnosed at age 18 y; received CD19 CAR T cells with subsequent consolidative HSCT from 10/10 matched sibling donor, with TBI -based myeloablative conditioning regimen (TBI and CTX)	History of multiply relapsed B-ALL, initially diagnosed at age 20 y; underwent haploidentical HSCT from parent, with TBI-based myeloablative conditioning regimen (TBI and CTX) in CR1 and CD19 CAR T-cell therapy in CR2	History of multiply relapsed B-ALL, initially diagnosed at age 21 y; underwent double cord blood HSCT, with TBI-based myeloablative conditioning regimen (fludarabine, CTX, and TBI) in first relapse. Received CD19 CAR T cells in second relapse	History of multiply relapsed/refractory B-ALL, initially diagnosed at age 8 y; received CD19 CAR T cells with subsequent consolidative HSCT from 9/10 matched unrelated donor, with TBI-based myeloablative conditioning regimen (TBI, CTX, and thiotepa)
Total radiation exposure (TBI), cGy	3600	1200 (plus 400 testicular boost)	1320	1200
Brief CAR T-cell therapy course	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR and proceeded directly to consolidative HSCT	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR; developed CD19 ⁻ relapse 6 mo post-CD19 CAR T-cell therapy, and received ALL-directed therapy before SMN diagnosis	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR; continued B-cell aplasia (>7 y)	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR and proceeded directly to consolidative HSCT
CAR T-cell construct	CD19/28z	CD19/BB	CD19/BB (tisagenlecleucel/CTL019)	CD19/28z
HSCT (before/after CD19 CAR T-cell therapy)	Yes, after	Yes, before	Yes, before	Yes, after
SMN presenting symptoms	Presented at routine follow-up 3 y post-CD19 CAR T-cell therapy and post-HSCT at age 26 y with asymptomatic mild elevation of transaminases	Dental pain and swelling noted after wisdom teeth removal 5 y post-CD19 CAR T-cell therapy at age 29 y	Growing mole on right buttock 6 y post-CD19 CAR T-cell therapy	Presented for follow-up 8 y post-CD19 CAR T-cell therapy and post-HSCT with suspicious thyroid nodule
Diagnosis	<u>MRI with MRCP</u> : hilar biliary obstruction extending into left hepatic duct with no obstructing mass/stone identified <u>ERCP</u> : focal biliary stricturing in common and left hepatic ducts <u>Diagnostic laparoscopy with exploration and resection</u> : frozen section of left hepatic duct positive for carcinoma, with final surgical pathology results showing moderately differentiated cholangiocarcinoma	<u>Biopsy</u> : synovial sarcoma, biphasic type (FNCLCC grade 2)	<u>Biopsy</u> : malignant melanoma, superficial spreading type, in radial growth phase	<u>Thyroidectomy</u> : 1.5-cm nodule that contained papillary carcinoma of thyroid with extrathyroidal extension and microscopic strap muscle invasion
Management	Underwent left hepatectomy with extrahepatic bile duct resection and portal lymphadenectomy	Underwent partial left maxillary teeth extraction, left mandibular reconstruction, soft tissue flap, bilateral level 1 and left level 2-4 lymph node dissection as well as 660-cGy proton radiation therapy	Local resection with negative margins	Thyroidectomy and ¹³¹ I radioactive iodine ablation
Outcome	Died on postoperative day 3 from fulminant hepatic failure	Continued remission from synovial sarcoma with active B-ALL	Continued remission from both B-ALL and malignant melanoma	Continued remission from both B-ALL and thyroid carcinoma

CR1, first complete remission; CR2, second complete remission; CTX, cyclophosphamide; ERCP, endoscopic retrograde cholangiopancreatography; FNCLCC, Fédération Nationale des Centres de Lutte Contre Le Cancer; MRCP, magnetic resonance cholangiopancreatography; MRD, minimal residual disease; MRI, magnetic resonance imaging.

including TBI, is likely causative for SMNs, additional long-term follow-up studies after CAR T-cell therapy are needed to evaluate the cumulative incidence and type of SMNs after CAR T-cell therapy as childhood cancer survivors age into adulthood.

Acknowledgments: The authors thank the treating and referring centers, care providers, supporting staff, and referring physicians who cared for the patients included in this study.

This work was supported in part by the Center for Cancer Research and the Warren Grant Magnuson Clinical Center, Intramural Research Program, National Cancer Institute, National Institutes of Health (grant ZIA BC 011823; N.N.S.). L.G. is supported by grant CA046934.

Contribution: E.M.H., N.N.S., and A.J.L. wrote the first draft of the manuscript; and all authors contributed to data collection and provided relevant details, substantially contributed to the final version of the manuscript, and approved the submission.

The content of this publication does not necessarily reflect the views or policies of the US Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the US Government.

Conflict-of-interest disclosure: D.W.L. has consulted for Harpoon Therapeutics, has advised for Amgen and Bristol Myers Squibb, and has received research funding from Kite Pharma and Gilead. S.A.G. has received research funding from Novartis, Kite

Table 3. Presentation, diagnosis, and management of hematologic SMNs

	AML/MDS	AML	MDS
Oncologic history	History of Li Fraumeni syndrome with very high-risk B-ALL diagnosed at age 18 y; attempt at curative frontline therapy with CD19 CAR T cells in hope of lesser toxicity compared with myeloablative regimen because of Li Fraumeni-associated risks; because of early loss of B-cell aplasia after CD19 CAR T-cell therapy at 4 mo, received CD19 CAR T-cell reinfusion, with subsequent development of MDS (chromosome 5 changes) 2 mo later; underwent unrelated cord blood HSCT for MDS using busulfan, fludarabine, and thiotepa-based conditioning regimen	History of disseminated B-lymphoblastic lymphoma (TCF-PBX1 translocation) diagnosed at age 10 y, including involvement of multiple cranial nerves; required emergency cranial XRT (2350 cGy) and boost (1500 cGy); isolated CNS relapse of B-lymphoblasts 2.5 y after initial diagnosis; received triple IT therapy and VCR/MTX as bridging therapy; received CD19 CAR T cells in first relapse	History of with B-ALL (unknown cytogenetics) diagnosed at age 10 y; first relapse was very early; received salvage chemotherapy followed by unrelated cord blood HSCT with TBI-based myeloablative conditioning regimen (TBI, etoposide, and CTX); second relapse was 2.5 y post-HSCT; received bridging chemotherapy while awaiting CD19 CAR T-cell infusion
Total radiation exposure (TBI), cGy	None	2850 (CNS only)	1200
Brief CAR T-cell therapy course	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells, without significant complications; good count recovery but developed early loss of B-cell aplasia and received CD19 CAR T-cell reinfusion	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR; continued B-cell aplasia	Received lymphodepleting chemotherapy with CTX and fludarabine before CD19 CAR T cells; achieved MRD ⁻ CR; continued B-cell aplasia until time of SMN diagnosis
CAR T-cell construct	CD19/BB	CD19/BB	CD19/BB
HSCT (before/after CD19 CAR T-cell therapy)	Yes, after	NA	Yes, before
Presenting symptoms	Progressive cytopenias (trilineage) after full count recovery after first CD19 CAR T-cell treatment and early post-HSCT; fatigue	Developed worsening pancytopenia 18 mo post-CD19 CAR T-cell therapy	Progressive thrombocytopenia and new fevers 13 mo post-CD19 CAR T-cell therapy
Diagnosis	<u>Bone marrow evaluation:</u> MDS/AML with loss of original clone and new cytogenetics consisting of translocation(5;17)/AML	<u>Bone marrow evaluation:</u> AML NPM1 ⁺ and CEBPA ⁺ ; negative for TCF3-PBX1 translocation	<u>Bone marrow evaluation:</u> MDS (RAEB-2), trisomy 8 and 9; MDS was XY and thought to be derived from cord blood
Management	Therapy with venetoclax/azacitidine to control disease; not eligible for second HSCT because of pulmonary GVHD/IPS complications from first HSCT	Therapy with CPX-351 (liposomal daunorubicin and cytarabine) and FLAG, then proceeded to 10/10 HLA-matched unrelated donor HSCT; myeloablative conditioning regimen with busulfan, fludarabine, and thiotepa	Therapy with azacitidine, then mismatched sibling HSCT; myeloablative conditioning regimen with busulfan and melphalan
Outcome	Patient died of refractory MDS/AML with minimal response to therapy and progressive IPS post-HSCT at ~3 mo post-HSCT	Alive without disease 14 mo post-HSCT	Alive without disease 18 mo post-second HSCT

AML, acute myeloid leukemia; CNS, central nervous system; CR, complete remission; FLAG, fludarabine, cytarabine, and filgrastim; GVHD, graft-versus-host disease; IPS, idiopathic pneumonia syndrome; IT, intrathecal; MDS, myelodysplastic syndrome; MRD, minimal residual disease; MTX, methotrexate; NA, not applicable; RAEB-2, refractory anemia with excess blasts type 2; VCR, vincristine; XRT, radiation therapy.

Pharma, Vertex, and Servier; has consulted for Novartis, Roche, GlaxoSmithKline, Humanigen, Cellular Biomedicine Group, Eureka, Janssen/Johnson & Johnson, and Jazz Pharmaceuticals; and has advised for Novartis, Adaptimmune, TCR2, Cellectis, Juno, Vertex, Allogene, Jazz Pharmaceuticals, and Cabaletta. M.R.V. has advised for Novartis, Equillium, Mederus, and Takeda. L.G. has consulted for Amgen, Novartis, and Roche; has advised for Amgen, Novartis, and Celgene; and holds equity in Amgen, Anchiano, Blueprint Medicines, Celgene, Clovis, Mirati, and Sanofi Paris. P.A.B. has advised for Novartis, Takeda, Amgen, Kura, and Kite. S.R.R. has received research funding from Pfizer. M.A.P. has advised for Mesoblast, Novartis, Equillium, Medexus, and Vertex; has received research funding from Adaptive and Miltenyi; and has received honoraria from Novartis, Miltenyi, and Bellicum. R.A.G. has consulted for Novartis and received patents

and royalties from Bristol Myers Squibb. T.W.L. has advised for Bayer, Cellectis, Novartis, Deciphera, Juno, and Y-mAbs Therapeutics; has received honoraria from Bayer, Cellectis, Novartis, Deciphera, Juno, and Y-mAbs Therapeutics; and has received research support from Pfizer and Bayer. The remaining authors declare no competing financial interests.

ORCID profiles: E.M.H., 0000-0001-5088-7621; R.M.M., 0000-0002-2542-2061; D.W.L., 0000-0002-3249-9796; P.A.B., 0000-0002-8653-1069; M.A.P., 0000-0003-3030-8420; S.R.R., 0000-0001-8025-6767; L.G., 0000-0002-2546-4616; N.N.S., 0000-0002-8474-9080; A.J.L., 0000-0001-5088-7621.

Correspondence: Adam J. Lamble, Division of Hematology/Oncology, University of Washington, Seattle Children's Hospital, Seattle, WA; e-mail: adam.lamble@seattlechildrens.org.

References

1. Friedman DL, Whitton J, Leisenring W, et al. Subsequent neoplasms in 5-year survivors of childhood cancer: the Childhood Cancer Survivor Study. *J Natl Cancer Inst*. 2010;102(14):1083-1095.
2. Meadows AT, Friedman DL, Neglia JP, et al. Second neoplasms in survivors of childhood cancer: findings from the Childhood Cancer Survivor Study cohort. *J Clin Oncol*. 2009;27(14):2356-2362.
3. Inamoto Y, Shah NN, Savani BN, et al. Secondary solid cancer screening following hematopoietic cell transplantation. *Bone Marrow Transplant*. 2015;50(8):1013-1023.
4. Cullinan N, Schiller I, Di Giuseppe G, et al. Utility of a cancer predisposition screening tool for predicting subsequent malignant neoplasms in childhood cancer survivors. *J Clin Oncol*. 2021;39(29):3207-3216.
5. Turcotte LM, Neglia JP, Reulen RC, et al. Risk, risk factors, and surveillance of subsequent malignant neoplasms in survivors of childhood cancer: a review. *J Clin Oncol*. 2018;36(21):2145-2152.
6. Dixon SB, Chen Y, Yasui Y, et al. Reduced morbidity and mortality in survivors of childhood acutelymphoblastic leukemia: a report from the childhood cancer survivor study. *J Clin Oncol*. 2020;38(29):3418-3429.
7. Center for Biologics Evaluation and Research, US Food and Drug Administration. Long Term Follow-Up After Administration of Human Gene Therapy Products; Guidance for Industry. Available at: <https://www.fda.gov/media/113768/download>. Accessed 1 April 2022.
8. Cornetta K, Duffy L, Turtle CJ, et al. Absence of replication-competent lentivirus in the clinic: analysis of infused T cell products. *Mol Ther*. 2018;26(1):280-288.
9. Cornetta K, Duffy L, Feldman SA, et al. Screening clinical cell products for replication competent retrovirus: the National Gene Vector Biorepository experience. *Mol Ther Methods Clin Dev*. 2018;10:371-378.
10. Myers RM, Taraseviciute A, Steinberg SM, et al. Blinatumomab nonresponse and high-disease burden are associated with inferior outcomes after CD19-CAR for B-ALL. *J Clin Oncol*. 2022;40(9):932-944.
11. Good ML, Malekzadeh P, Kriley IR, et al. Intrahepatic cholangiocarcinoma as a rare secondary malignancy after allogeneic hematopoietic stem cell transplantation for childhood acute lymphoblastic leukemia: a case report. *Pediatr Transplant*. 2020;24(2):e13653.
12. Levine JE, Grupp SA, Pulsipher MA, et al. Pooled safety analysis of tisagenlecleucel in children and young adults with B cell acute lymphoblastic leukemia. *J Immunother Cancer*. 2021;9(8):e002287.
13. Cappell KM, Sherry RM, Yang JC, et al. Long-term follow-up of anti-CD19 chimeric antigen receptor T-cell therapy. *J Clin Oncol*. 2020;38(32):3805-3815.
14. Cordeiro A, Bezerra ED, Hirayama AV, et al. Late events after treatment with CD19-targeted chimeric antigen receptor modified T cells. *Biol Blood Marrow Transplant*. 2020;26(1):26-33.
15. Morton LM, Saber W, Baker KS, et al. National Institutes of Health hematopoietic cell transplantation late effects initiative: the Subsequent Neoplasms Working Group report. *Biol Blood Marrow Transplant*. 2017;23(3):367-378.
16. Peters C, Dalle JH, Locatelli F, et al; EBMT Paediatric Diseases Working Party. Total body irradiation or chemotherapy conditioning in childhood ALL: a multinational, randomized, noninferiority phase III study. *J Clin Oncol*. 2021;39(4):295-307.
17. Willasch AM, Peters C, Sedláček P, et al; EBMT Paediatric Diseases Working Party. Myeloablative conditioning for allo-HSCT in pediatric ALL: FTBI or chemotherapy?-A multicenter EBMT-PDWP study [published correction appears in *Bone Marrow Transplant*. 2021;56(10):2615]. *Bone Marrow Transplant*. 2020;55(8):1540-1551.
18. Lamble A, Myers RM, Taraseviciute A, et al. Preinfusion factors impacting relapse immunophenotype following CD19 CAR T cells [published online ahead of print 28 April 2022]. *Blood Adv*.
19. Du J, Zhang Y. Sequential anti-CD19, 22, and 20 autologous chimeric antigen receptor T-cell (CAR-T) treatments of a child with relapsed refractory Burkitt lymphoma: a case report and literature review [published correction appears in *J Cancer Res Clin Oncol*. 2020;146(8):2177]. *J Cancer Res Clin Oncol*. 2020;146(6):1575-1582.
20. Pulsipher MA, Han X, Maude SL, et al. Next-generation sequencing of minimal residual disease for predicting relapse after tisagenlecleucel in children and young adults with acute lymphoblastic leukemia. *Blood Cancer Discov*. 2022;3(1):66-81.
21. Mo G, Wang HW, Talleur AC, et al. Diagnostic approach to the evaluation of myeloid malignancies following CAR T-cell therapy in B-cell acute lymphoblastic leukemia. *J Immunother Cancer*. 2020;8(2):e001563.
22. Bishop MR, Maziarz RT, Waller EK, et al. Tisagenlecleucel in relapsed/refractory diffuse large B-cell lymphoma patients without measurable disease at infusion. *Blood Adv*. 2019;3(14):2230-2236.
23. Micklethwaite KP, Gowrishankar K, Gloss BS, et al. Investigation of product-derived lymphoma following infusion of piggyBac-modified CD19 chimeric antigen receptor T cells. *Blood*. 2021;138(16):1391-1405.