

Integrated flow cytometry and sequencing to reconstruct evolutionary patterns from dysplasia to acute myeloid leukemia

Catia Simoes,¹ Maria-Carmen Chillon,² David Martínez-Cuadrón,³ Maria-José Calasanz,^{1,4} Maria-Belén Vridiales,² Iria Vazquez,^{1,4} Montserrat Hernández-Ruano,² Beñat Ariceta,¹ Paula Aguirre-Ruiz,¹ Leire Burgos,¹ Diego Alignani,¹ Sarai Sarvide,¹ Sara Villar,¹ Ana Alfonso Pierola,¹ Felipe Prosper,¹ Rosa Ayala,⁵ Joaquín Martínez-López,⁵ Juan Miguel Bergua Burgues,⁶ Susana Vives,⁷ Jose A. Perez-Simon,⁸ Maria Garcia-Fortes,⁹ Teresa Bernal del Castillo,¹⁰ Mercedes Colorado,¹¹ Mayte Olave,¹² Juan I. Rodríguez-Gutiérrez,¹³ Jorge Labrador,¹⁴ Marcos González,² Jesús F. San-Miguel,¹ Miguel Ángel Sanz,³ Pau Montesinos,³ Bruno Paiva,¹ on behalf of the Programa para Tratamientos en Hematología (PETHEMA) Cooperative Study Group

¹Departamento de Hemato-Oncology, Clínica Universidad de Navarra, Centro de Investigación Médica Aplicada (CIMA), Instituto de Investigación Sanitaria de Navarra (IDISNA), Pamplona, Spain; ²Centro de Investigación del Cáncer de Salamanca-IBMCC (USAL-CSIC), Hospital Universitario de Salamanca, IBSAL, Salamanca, Spain; ³Hospital Universitario y Politécnico La Fe, Valencia, Spain; ⁴Cima Lab Diagnostics, Universidad de Navarra, IDISNA, Pamplona, Spain; ⁵Hospital Universitario 12 de Octubre, Universidad Complutense, Madrid, Spain; ⁶Hospital San Pedro de Alcántara, Cáceres, Spain; ⁷Hospital Germans Trias i Pujol, Badalona, Spain; ⁸Hospital Universitario Virgen del Rocío, Seville, Spain; ⁹Hospital Universitario Virgen de la Victoria, Málaga, Spain; ¹⁰Hospital Central de Asturias, SPA, IUOPA, Oviedo, Spain; ¹¹Hospital Universitario Marqués de Valdecilla, Cantabria, Spain; ¹²Hospital Clínico Universitario Lozano Blesa, Zaragoza, Spain; ¹³Hospital Universitario de Basurto, Bilbao, Spain; and ¹⁴Hospital Universitario de Burgos, Burgos, Spain

Key Points

- Targeted sequencing of dysplastic cells and blasts at diagnosis identifies evolutionary patterns from dysplasia to the onset of AML.
- Patients with AML may present different clonal involvement in dysplastic myeloid-erythropoiesis, leukemic transformation, and chemoresistance.

Clonal evolution in acute myeloid leukemia (AML) originates long before diagnosis and is a dynamic process that may affect survival. However, it remains uninvestigated during routine diagnostic workups. We hypothesized that the mutational status of bone marrow dysplastic cells and leukemic blasts, analyzed at the onset of AML using integrated multidimensional flow cytometry (MFC) immunophenotyping and fluorescence-activated cell sorting (FACS) with next-generation sequencing (NGS), could reconstruct leukemogenesis. Dysplastic cells were detected by MFC in 285 of 348 (82%) newly diagnosed patients with AML. Presence of dysplasia according to MFC and World Health Organization criteria had no prognostic value in older adults. NGS of dysplastic cells and blasts isolated at diagnosis identified 3 evolutionary patterns: stable (n = 12 of 21), branching (n = 4 of 21), and clonal evolution (n = 5 of 21). In patients achieving complete response (CR), integrated MFC and FACS with NGS showed persistent measurable residual disease (MRD) in phenotypically normal cell types, as well as the acquisition of genetic traits associated with treatment resistance. Furthermore, whole-exome sequencing of dysplastic and leukemic cells at diagnosis and of MRD uncovered different clonal involvement in dysplastic myeloid-erythropoiesis, leukemic transformation, and chemoresistance. Altogether, we showed that it is possible to reconstruct leukemogenesis in ~80% of patients with newly diagnosed AML, using techniques other than single-cell multiomics.

Submitted 19 May 2022; accepted 12 September 2022; prepublished online on *Blood Advances* First Edition 14 October 2022. <https://doi.org/10.1182/bloodadvances.2022008141>.

The sequencing data reported in this article have been deposited in the European Genome-phenome Archive database (accession number EGAS00001006318).

Data are available on request from the corresponding author, Bruno Paiva (bpaiva@unav.es).

The full-text version of this article contains a data supplement.

© 2022 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

Introduction

Clonal evolution in acute myeloid leukemia (AML) originates long before diagnosis and is a highly dynamic process that may affect survival.¹ Thus, having the possibility of defining in each patient the genetic events causing AML would be a benefit to routine diagnostic workups. However, dissecting leukemic transformation at the onset of AML is challenging without single-cell sequencing,^{2,3} and most clinical laboratories do not have the infrastructure to perform single-cell studies routinely.

Patients with newly diagnosed AML may present with dysplasia regardless of their clinical ontogeny (ie, secondary AML [sAML] vs therapy-related vs de novo AML).⁴⁻⁶ If these residual, mature dysplastic cells were generated before the differentiation blockage of blasts preceding leukemic transformation, it could be hypothesized that studying the mutational profile of dysplastic cells and blasts could uncover the evolutionary process from dysplasia to AML. This hypothesis has never been investigated.

Methods

We aimed at reconstructing clonal evolution from dysplasia to AML based on the analysis of bone marrow (BM) dysplastic cells and leukemic blasts using integrated multidimensional flow cytometry (MFC) immunophenotyping and fluorescence-activated cell sorting (FACS) with next-generation sequencing (NGS) (Figure 1A).

In total, 358 patients with AML were studied after signing informed consent forms approved by the ethics committees, in accordance with the Declaration of Helsinki. The presence of dysplasia according to aberrant phenotypic differentiation of the neutrophil, monocytic, and erythroid lineages was investigated using EuroFlow protocols^{7,8} in 348 patients with newly diagnosed AML; 103 were eligible for intensive treatment, whereas 245 were older adults and enrolled in the phase 3 PETHEMA/FLUGAZA trial.⁹ Up to 168 cell types from a total of 38 patients with AML were isolated at diagnosis and in complete response for deep-targeted (supplemental Table 1) or exome sequencing. T cells were used as control. A description of methods is available in the supplemental Data.

Results and Discussion

We first assessed the applicability of our hypothesis by investigating how many patients show dysplasia by MFC at the onset of AML (supplemental Figure 1). Dysplastic cells were observed in 285 of 348 (82%) cases (supplemental Figure 2A). Phenotypic abnormalities were more frequent in neutrophils (45%), followed by monocytic (32%) and erythroid cells (23%) (supplemental Figure 2B). Lack of data on aberrant megakaryocytic differentiation is a limitation of the study. Multilineage dysplasia was detected in 186 (53%) patients. Only 35 (10%) cases had no signs of dysplasia, whereas the remaining 28 (8%) had nearly undetectable hematopoiesis due to massive blast infiltration. The presence of dysplasia was less frequent in patients eligible for intensive chemotherapy vs those ineligible (75% vs 96%, $P < .001$). Thus, the median age of the series (69 years) may help in explaining presence of dysplastic cells in most patients.

In the cohort of older patients enrolled in the PETHEMA/FLUGAZA trial, there were no differences in overall survival between those

with or without dysplasia according to MFC (supplemental Figure 2C). The presence of dysplasia assessed by MFC and diagnosis of AML with morphological myelodysplasia-related changes were not significantly associated (supplemental Table 2). Diagnostic classification of AML with myelodysplasia-related changes vs other subtypes had no impact on overall survival (supplemental Figure 2D). These results in older patients treated with semi-intensive chemotherapy or hypomethylating agents support previous observations in other treatment scenarios of the limited prognostic value of assessing dysplasia in AML.^{10,11}

Targeted sequencing of dysplastic cells and blasts isolated by FACS in 21 patients with newly diagnosed AML uncovered 3 evolutionary patterns (Figure 1B). Stable evolution in those displaying identical mutational profiles in blasts and dysplastic cells ($n = 12$ of 21), branching evolution in patients where blasts may have originated from leukemic stem cells other than the ones driving dysplasia due to mutations absent in blasts and present in dysplastic cells ($n = 4$ of 21), and clonal evolution in cases showing new mutations in blasts onto mutations shared between these and dysplastic cells ($n = 5$ of 21). These results are not novel but rather consistent with previous observations in whole BM samples sequenced before and after leukemic transformation^{4,12} and in FACS stem cell compartments in paired samples from 7 patients progressing from myelodysplastic syndrome to sAML.¹³ Furthermore, using single-cell DNA sequencing, Morita et al³ showed how single-cell data reveal cell-level mutation co-occurrence and enable reconstruction of mutational histories characterized by linear and branching patterns of clonal evolution. Accordingly, targeted sequencing of dysplastic cells and blasts at the onset of AML may help to recreate the history of the disease despite the notably lower resolution of this approach when compared with single-cell DNA sequencing.

The unique experimental design of this study yielded interesting observations (Figure 1C). Mutations in the cohesin complex and in transcription factors were never exclusive of dysplastic cells, whereas spliceosome mutations were never exclusive of blasts, which is consistent with their role in sAML.^{14,15} Mutations in genes involved in DNA methylation or tumor suppressors were systematically present in dysplastic cells and blasts. At the individual patient level, mutations in *TP53* appeared to drive leukemogenesis in patients 1, 2, and 3 (Figure 1B). Conversely, leukemic transformation in patient 18 was associated with a new mutation in *STAG2* onto a background of mutated *DNMT3A* and *TP53* in dysplastic and leukemic cells. Other mutations observed exclusively in blasts were found in *BCORL1*, *FLT3-ITD*, and *PTPN11*; all 3 genes are associated with poor prognosis in AML,¹⁶⁻¹⁸ *FLT3-ITD* is a therapeutic target,¹⁹ and *PTPN11* may drive resistance to newer targeted therapies.²⁰ Of note, subclonal mutations may be missed if whole BM samples are sequenced instead of isolated cell types (supplemental Figure 3).

The increased sensitivity in detecting mutations granted by prior FACS urged us to use the same experimental approach in 10 patients achieving CR after induction, stratified according to undetectable vs persistent MRD by MFC. Phenotypically normal CD34⁺ hematopoietic progenitor cells (HPCs) were isolated in the former, whereas MRD cells were sorted in the latter patients; cells of the neutrophil, monocyte, and erythroid lineages were isolated in both.




Figure 1. Reconstructing clonal evolution from dysplasia to AML. (A) Study design. BM aspirates collected from 358 patients with newly diagnosed AML were analyzed using MFC. In 21 patients, leukemic cells were isolated using FACS according to patient-specific aberrant phenotypes and so were cells of the neutrophil (Neutro), monocytic (Mono), and erythroid (Erythro) lineages whenever dysplastic phenotypes were observed in 1 or more lineages. T cells were systematically isolated for germline DNA. NGS of genes frequently mutated in myeloid neoplasms was performed in all cell types available in each patient. (B) Mutational status of 33 genes in T cells (T), neutrophils (N), monocytes (M), erythroblasts (E), and blasts (B) isolated from 21 patients at the onset of AML; cell types not available for NGS are represented with gray lines. Mutated genes were colored in a gradient of red according to the variant allele frequency (VAF). Three models were identified: (1) stable transition according to identical mutational landscapes in blasts and dysplastic cells; (2) branching evolution with blasts originating from leukemic stem cells other than the ones driving dysplasia, due to mutations absent in blasts and present in dysplastic cells; and (3) clonal evolution with new mutations in blasts onto mutations shared between these and dysplastic cells. (C) Percentage of mutations grouped according to functional categories that were simultaneously mutated in mature dysplastic cells and blasts (gray), or that were private in the former (blue) and the latter (purple). Functional categories with significantly different distributions were highlighted in bold.




Figure 1 (continued)

Among patients with undetectable MRD, most mutations present at diagnosis became undetectable after induction (Figure 2A). One exception was mutated *DNMT3A* in CD34⁺ HPCs from patient 23, which could represent postremission clonal hematopoiesis,²¹ and in erythroid cells from patient 24, which could represent the expansion of hematopoietic populations unrelated to the AML founding clone after induction therapy.²² Although the clinical implications of postremission clonal hematopoiesis remain poorly understood,²¹ by performing NGS in isolated cell types, we showed that it is possible to monitor mutations with low variant allele frequency (VAF) present in single-cell compartments. Patient 27 unexpectedly showed mutated *IDH2* and *JAK2* in CD34⁺ HPCs and cells of the neutrophil, monocyte, and erythroid lineages, with VAF similar to that found at diagnosis. These results may reflect the inability of MFC to detect leukemic cells and their differentiated progeny in the selected patients with AML (Figure 2A). Accordingly, NGS performed in isolated cell types could reduce the rate of false-negative results in patients with undetectable MRD by MFC.^{23,24}

Among patients with persistent MRD, most mutations present at diagnosis were detected after induction, both in MRD and cells of the neutrophil, monocyte, and erythroid lineages (Figure 2B). These results may help to explain the presence of dysplasia detectable by MFC during MRD assessment, particularly in patients with persistent MRD.²⁵ Interestingly, new mutations emerged in MRD cells and their differentiated progeny in patients 30 and 31 (ie, *PHF6* and *TET2*, respectively), which could either represent the acquisition of genetic traits enabling treatment resistance or the selection and expansion of minor clones hidden at diagnosis.^{3,26}

To investigate if our experimental approach could provide insights into the overlap between genetic alterations associated with leukemic transformation and chemoresistance, we relied on the broader coverage of exome sequencing to compare the genomic profile of dysplastic and leukemic cells at diagnosis and of MRD cells after induction in 6 patients with AML. In 5 of the 6 cases, there were mutations in dysplastic cells undetected in blasts at diagnosis and MRD (eg, *MUC16* and *FAM155B* in patients 34 and 37, respectively), mutations shared by dysplastic cells and blasts at diagnosis undetected in MRD (eg, *ZNF717* and *MUC12* in

patients 36 and 37, respectively), and mutations shared by dysplastic cells and MRD but undetected in blasts at diagnosis (eg, *MUC2* and *KIR2DL3* in patients 33 and 35, respectively) (supplemental Figure 4; Figure 2C). These results uncover different clonal involvement in dysplastic myelo-erythropoiesis, leukemic transformation, and chemoresistance, in accordance with recent single-cell analysis suggesting that clones emerging during AML transformation and those selected by treatment may differ.^{2,3}

In conclusion, we showed for the first time that it is possible to reconstruct leukemogenesis in ~80% of patients with newly diagnosed AML, using techniques other than single-cell multiomics. The possibility of identifying the genetic drivers of leukemic transformation and chemoresistance could be clinically meaningful to develop tailored treatment strategies aiming at the eradication of genetically diverse leukemic clones.

Acknowledgments

The authors acknowledge the patients, caregivers, and the biobank of the University of Navarra.

This work was supported by grants from the Área de Oncología del Instituto de Salud Carlos III, Centro de Investigación Biomédica en Red (CIBER-ONC) (CB16/12/00369, CB16/12/00233, CB16/12/00489, and CB16/12/00284), Instituto de Salud Carlos III/Subdirección General de Investigación Sanitaria (FIS numbers PI16/01661, PI16/00517, and PI19/01518), and the Plan de Investigación de la Universidad de Navarra (PIUNA 2014-18). This work was supported internationally by the Cancer Research UK, FCAECC, and AIRC under the Accelerator Award Program (EDITOR).

Authorship

Contribution: C.S., J.F.S.-M., P.M., and B.P. conceived and designed the study; C.S., M.-C.C., I.V., B.A., L.B., and B.P. analyzed and interpreted the data; C.S. and B.P. performed the statistical analysis; M.C.-C., D.M.-C., M.-J.C., M.B.V., M.H.-R., P.A.-R., D.A., S.S., S. Villar, A.A.P., F.P., R.A., J.M.-L., J.M.B.B., S. Vives, J.A.P.-S., M.G.-F., T.B.d.C., M.C., M.O., J.I.R.-G., J.L., M.G., M.A.S., and P.M. provided the study material and/or patients; C.S. and B.P. drafted the manuscript; and all authors reviewed and approved the manuscript.




Figure 2. Different clonal involvement in dysplastic myelo-erythropoiesis, leukemic transformation, and chemoresistance. (A-B) Patients (Pat.) with undetectable (n = 5) and persistent (n = 5) MRD had phenotypically normal CD34⁺ HPCs and MRD cells, respectively isolated, whereas cells of the neutrophil, monocyte, and erythroid lineages were isolated in all patients. NGS of genes frequently mutated in myeloid neoplasms was performed in all cell types available in each patient. The VAF of mutated genes was colored in a gradient of gray. If 2 mutations in the same gene were detected, the 2 VAFs are indicated. (C) Representative patient with exome sequencing of dysplastic cell types, blasts at diagnosis, and persistent MRD. The fish plot illustrates different clonal compositions at different stages of disease progression. The blue arrow is pointing to a mutation present in dysplastic cells though absent in blasts at diagnosis and MRD, the olive arrow is pointing to mutations present in dysplastic cells and blasts at diagnosis though not at MRD, and the black arrow points to mutations present in MRD and dysplastic cells though not in blasts at diagnosis. The bar widths indicate the respective VAFs.




Figure 2 (continued)

Conflict-of-interest disclosure: F.P. received honoraria and research funding from Oryzon, Janssen, Bristol-Myers Squibb (BMS)/Celgene. R.A. serves as a member of the board of directors advisory committees for Incyte Corporation and Astellas and received honoraria from Novartis, Celgene, and Incyte. J.A.P.-S. received honoraria and funding budget for research projects and is active on the advisory board and learning activities or conferences of Janssen, Takeda, Pfizer, Jazz, BMS, Amgen, and Gilead. J.F.S.-M. does consultancy and is a member of the board of directors advisory committees for AbbVie, Amgen, BMS, Celgene, GlaxoSmithKline (GSK), Janssen, Karyopharm, Merck Sharpe & Dohme, Novartis, Regeneron, Roche, Sanofi, SecuraBio, and Takeda. P.M. provides consultancy, is a member of the board of directors advisory committees and speaker's bureau, and received research funding from Celgene, Sanofi, Incyte, Karyopharm, Novartis, Stemline/Menarini, Agios, Astellas Pharma, and Daiichi Sankyo; is also a member of the board of directors advisory committees for Pfizer, Teva, and AbbVie; received research funding from and is a member of speaker's bureau for Janssen; and provides consultancy for Tolero Pharmaceutical, Forma Therapeutics, and Glycomimetics. B.P. served as a consultant and received honoraria from Adaptive, Amgen, BD Biosciences, BMS/Celgene, GSK,

Janssen, Roche, Sanofi, and Takeda and received research support from BMS/Celgene, GSK, Roche, Sanofi, and Takeda. The remaining authors declare no competing financial interests.

ORCID profiles: C.S., [0000-0002-5713-8428](https://orcid.org/0000-0002-5713-8428); M.-C.C., [0000-0003-1624-6059](https://orcid.org/0000-0003-1624-6059); D.M.-C., [0000-0001-7540-4091](https://orcid.org/0000-0001-7540-4091); M.-J.C., [0000-0002-0374-3008](https://orcid.org/0000-0002-0374-3008); I.V., [0000-0003-0027-6384](https://orcid.org/0000-0003-0027-6384); B.A., [0000-0001-8819-402X](https://orcid.org/0000-0001-8819-402X); P.A.-R., [0000-0002-5871-372X](https://orcid.org/0000-0002-5871-372X); L.B., [0000-0003-0998-2496](https://orcid.org/0000-0003-0998-2496); D.A., [0000-0003-0241-1375](https://orcid.org/0000-0003-0241-1375); S.V., [0000-0002-4761-6692](https://orcid.org/0000-0002-4761-6692); A.A.P., [0000-0002-2478-5354](https://orcid.org/0000-0002-2478-5354); F.P., [0000-0001-6115-8790](https://orcid.org/0000-0001-6115-8790); R.A., [0000-0002-2699-8353](https://orcid.org/0000-0002-2699-8353); J.M.-L., [0000-0001-7908-0063](https://orcid.org/0000-0001-7908-0063); J.M.B.B., [0000-0001-8229-1208](https://orcid.org/0000-0001-8229-1208); M.G.-F., [0000-0002-0959-9449](https://orcid.org/0000-0002-0959-9449); T.B.d.C., [0000-0002-2338-513X](https://orcid.org/0000-0002-2338-513X); M.C., [0000-0002-4036-5050](https://orcid.org/0000-0002-4036-5050); M.O., [0000-0002-6326-5992](https://orcid.org/0000-0002-6326-5992); M.G., [0000-0001-6637-1072](https://orcid.org/0000-0001-6637-1072); J.F.S.-M., [0000-0002-9183-4857](https://orcid.org/0000-0002-9183-4857); M.Á.S., [0000-0003-1489-1177](https://orcid.org/0000-0003-1489-1177); P.M., [0000-0002-3275-5593](https://orcid.org/0000-0002-3275-5593); B.P., [0000-0003-1977-3815](https://orcid.org/0000-0003-1977-3815).

Correspondence: Bruno Paiva, Clínica Universidad de Navarra, Centro de Investigación Médica Aplicada (CIMA), Instituto de Investigación Sanitaria de Navarra (IDISNA), Avenida Pio XII 55, 31008 Pamplona, Spain; email: bpaiva@unav.es.

References

1. Benard BA, Leak LB, Azizi A, Thomas D, Gentles AJ, Majeti R. Clonal architecture predicts clinical outcomes and drug sensitivity in acute myeloid leukemia. *Nat Commun.* 2021;12(1):1-13.
2. Miles LA, Bowman RL, Merlinsky TR, et al. Single-cell mutation analysis of clonal evolution in myeloid malignancies. *Nature.* 2020;587(7834):477-482.
3. Morita K, Wang F, Jahn K, et al. Clonal evolution of acute myeloid leukemia revealed by high-throughput single-cell genomics. *Nat Commun.* 2020; 11(1):1-17.
4. Lindsley RC, Mar BG, Mazzola E, et al. Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood.* 2015;125(9):1367-1376.
5. Weinberg OK, Pozdnyakova O, Campigotto F, et al. Reproducibility and prognostic significance of morphologic dysplasia in de novo acute myeloid leukemia. *Mod Pathol.* 2015;28(7):965-976.
6. Miesner M, Haferlach C, Bacher U, et al. Multilineage dysplasia (MLD) in acute myeloid leukemia (AML) correlates with MDS-related cytogenetic abnormalities and a prior history of MDS or MDS/MPN but has no independent prognostic relevance: a comparison of 408 cases classified as "AML not otherwise specified" (AML-NOS) or "AML with myelodysplasia-related changes" (AML-MRC). *Blood.* 2010;116(15):2742-2751.

7. Kalina T, Flores-Montero J, Van Der Velden VHJ, et al. EuroFlow standardization of flow cytometer instrument settings and immunophenotyping protocols. *Leukemia*. 2012;26(9):1986-2010.
8. van Dongen JJM, Lhermitte L, Böttcher S, et al. EuroFlow antibody panels for standardized n-dimensional flow cytometric immunophenotyping of normal, reactive and malignant leukocytes. *Leukemia*. 2012;26(9):1908-1975.
9. Vives S, Martínez-Cuadrón D, Bergua Burgues J, et al. A phase 3 trial of azacitidine versus a semi-intensive fludarabine and cytarabine schedule in older patients with untreated acute myeloid leukemia. *Cancer*. 2021;127(12):2003-2014.
10. Duchmann M, Wagner-Ballon O, Boyer T, et al. Machine learning identifies the independent role of dysplasia in the prediction of response to chemotherapy in AML. *Leukemia*. 2021;36(3):656-663.
11. Falini B, Maciejewski K, Weiss T, et al. Multilineage dysplasia has no impact on biologic, clinicopathologic, and prognostic features of AML with mutated nucleophosmin (NPM1). *Blood*. 2010;115(18):3776-3786.
12. Shlush LI, Zandi S, Mitchell A, et al. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature*. 2014;506(7488):328-333.
13. Chen J, Kao YR, Sun D, et al. Myelodysplastic syndrome progression to acute myeloid leukemia at the stem cell level [published correction appears in *Nat Med*. 2018]. *Nat Med*. 2019;25(1):103-110.
14. Makishima H, Visconte V, Sakaguchi H, et al. Mutations in the spliceosome machinery, a novel and ubiquitous pathway in leukemogenesis. *Blood*. 2012;119(14):3203-3210.
15. Sperling AS, Gibson CJ, Ebert BL. The genetics of myelodysplastic syndrome: from clonal haematopoiesis to secondary leukaemia. *Nat Rev Cancer*. 2016;17(1):5-19.
16. Damm F, Chesnais V, Nagata Y, et al. BCOR and BCORL1 mutations in myelodysplastic syndromes and related disorders. *Blood*. 2013;122(18):3169-3177.
17. Sallmyr A, Fan J, Datta K, et al. Internal tandem duplication of FLT3 (FLT3/ITD) induces increased ROS production, DNA damage, and misrepair: implications for poor prognosis in AML. *Blood*. 2008;111(6):3173-3182.
18. Stasik S, Eckardt JN, Kramer M, et al. Impact of PTPN11 mutations on clinical outcome analyzed in 1529 patients with acute myeloid leukemia. *Blood Adv*. 2021;5(17):3279-3289.
19. Wooten DJ, Gebru M, Wang HG, Albert R. Data-driven math model of FLT3-ITD acute myeloid leukemia reveals potential therapeutic targets. *J Pers Med*. 2021;11(3):193.
20. Stevens BM, Jones CL, Winters A, et al. PTPN11 mutations confer unique metabolic properties and increase resistance to venetoclax and azacitidine in acute myelogenous leukemia. *Blood*. 2018;132(suppl 1):909.
21. Tanaka T, Morita K, Loghavi S, et al. Clonal dynamics and clinical implications of postremission clonal hematopoiesis in acute myeloid leukemia. *Blood*. 2021;138(18):1733-1739.
22. Wong TN, Miller CA, Kloco JM, et al. Rapid expansion of preexisting nonleukemic hematopoietic clones frequently follows induction therapy for de novo AML. *Blood*. 2016;127(7):893-897.
23. Jongen-Lavrencic M, Grob T, Hanekamp D, et al. Molecular minimal residual disease in acute myeloid leukemia. *N Engl J Med*. 2018;378(13):1189-1199.
24. Getta BM, Devlin SM, Levine RL, et al. Multicolor flow cytometry and multigene next-generation sequencing are complementary and highly predictive for relapse in acute myeloid leukemia after allogeneic transplantation. *Biol Blood Marrow Transplant*. 2017;23(7):1064-1071.
25. Loghavi S, DiNardo CD, Furudate K, et al. Flow cytometric immunophenotypic alterations of persistent clonal haematopoiesis in remission bone marrows of patients with NPM1-mutated acute myeloid leukaemia. *Br J Haematol*. 2021;192(6):1054-1063.
26. Ediriwickrema A, Aleshin A, Reiter JG, et al. Single-cell mutational profiling enhances the clinical evaluation of AML MRD. *Blood Adv*. 2020;4(5):943-952.