

TO THE EDITOR:

Cellular and humoral immune responses to SARS-CoV-2 vaccination in patients after CD19.CAR T-cell therapy

Hannah Reimann,^{1,*} Anita N. Kremer,^{1,*} Viktoria Blumenberg,²⁻⁴ Katja Schmidt,⁵ Michael Aigner,¹ Benedikt Jacobs,¹ Nina Eisenhauer,¹ Alina Kämpf,¹ Wolf Rösler,¹ Soraya Kharboutli,¹ Dimitrios Mougiakakos,^{1,6} Vanessa Lang,¹ Christopher Lischer,⁷ Pascal Irrgang,⁸ Moritz Leppkes,⁹ Julio Vera Gonzalez,⁷ Gerhard Krönke,⁵ Andreas E. Kremer,^{9,10} Matthias Tenbusch,⁸ Heiko Bruns,¹ Thomas Harrer,⁵ Fabian Müller,¹ Georg Schett,⁵ Andreas Mackensen,¹ Marion Subklewe,^{2-4,*} and Simon Völkl^{1,*}

¹Department of Internal Medicine 5, Hematology and Oncology, Friedrich-Alexander-Universität Erlangen-Nürnberg and University Hospital Erlangen, Erlangen, Germany; ²Department of Medicine III, University Hospital, Ludwig Maximilian University of Munich, Munich, Germany; ³German Cancer Consortium and German Cancer Research Center, Heidelberg, Germany; ⁴Laboratory for Translational Cancer Immunology, Gene Center, Ludwig Maximilian University of Munich, Munich, Germany; ⁵Department of Internal Medicine 3, Rheumatology and Immunology, Friedrich-Alexander-Universität Erlangen-Nürnberg and University Hospital Erlangen, Erlangen, Germany; ⁶Department for Hematology and Oncology, Otto-von-Guericke University, Magdeburg, Germany; ⁷Department of Dermatology, Friedrich-Alexander-Universität Erlangen-Nürnberg and University Hospital Erlangen, Erlangen, Germany; ⁸Institute of Clinical and Molecular Virology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen and University Hospital, Erlangen, Germany; ⁹Department of Internal Medicine 1, Friedrich-Alexander-Universität Erlangen-Nürnberg and University Hospital Erlangen, Erlangen, Germany; and ¹⁰Department of Gastroenterology and Hepatology, University Hospital Zürich, University of Zürich, Zürich, Switzerland

The coronavirus disease 2019 (COVID-19) pandemic led to millions of infections and deaths worldwide. Patients with cancer and particularly those with hematologic malignancies showed an increased probability of a severe outcome including death.¹ Immense efforts led to a rapid development of vaccines against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections.^{2,3}

Despite the high efficacy of the vaccination, the benefit for severely immunocompromised patients such as those undergoing anti-CD19 chimeric antigen receptor T-cell (CAR-T) therapies is still uncertain. After CAR-T therapy, patients often display a long-term B-cell depletion and presumably do not develop proper humoral immunity in response to vaccinations. In line, only 11% to 36% of patients vaccinated after CAR-T therapy developed antibodies against the SARS-CoV-2–derived spike (S) protein.^{4,5} However, it remains unknown whether these patients still benefit from vaccination by developing cellular immunity. At present, most patients treated with CAR T cells are vaccinated against SARS-CoV-2, which is why the question of whether CAR-T therapy has an impact on preexisting cellular or humoral immune responses is gaining increasing relevance.

Here, we sought to analyze whether patients vaccinated after CAR-T therapy develop cellular immunity despite lacking antibody levels and whether vaccination before CAR-T therapy allows for sustained immunity throughout and beyond lymphodepletion and CAR-T therapy. Therefore, we analyzed humoral and cellular immune responses to SARS-CoV-2 in 21 and 15 patients vaccinated before and after CAR-T therapy, respectively. [Table 1](#) displays the baseline characteristics of all patients, with none of them having a confirmed COVID-19 infection diagnosis before the vaccination. Furthermore, 15 convalescent donors and 15 vaccinated healthy controls (HCs) served as the control group.

First, we tested 15 patients who received CAR-T therapy before SARS-CoV-2 vaccination ([Figure 1A-B](#)). Of 15 patients, 13 were still completely B-cell depleted at the time of vaccination. CAR T cells were detected in 10 patients ([supplemental Table 1](#)). Consistent with recent studies,⁴⁻⁶ only 2 (13%) of the patients developed neutralizing antibodies. In contrast, all convalescent donors and vaccinated HCs developed anti-S immunoglobulin G (IgG), with most of them being virus neutralizing. To determine whether patients may develop a humoral response over time, we repeated antibody

Submitted 8 April 2022; accepted 12 August 2022; prepublished online on *Blood Advances* First Edition 7 October 2022; final version published online 11 May 2023.
<https://doi.org/10.1182/bloodadvances.2022007806>.

*H.R., A.N.K., M.S., and S.V. contributed equally to this study.

Data are available on request from the corresponding author, Simon Völkl (simon.voelkl@uk-erlangen.de).

The full-text version of this article contains a data supplement.

© 2023 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](#), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

Table 1. Demographics and clinical characteristics

Patient characteristics	n = 36
Age, median (range)	65 (19-83)
Age <65 y, n (%)	16 (44)
Disease, n (%)	
DLBCL	20 (56)
Other NHL	9 (25)
Other	7 (19)
CD19 CAR T-cell product, n (%)	
Tisa-cel	10 (28)
Axi-cel	10 (28)
Other	16 (44)
Status of disease, n (%)	
Remission	31 (86)
Refractory	5 (14)
Postinfusion complications and related treatments, n (%)	
CRS	27 (75)
ICANS	13 (36)
Corticosteroids	14 (39)
anti-IL-6R	27 (75)
Vaccination type, n (%)	
BNT162b2 (Pfizer-BioNTech)	23 (64)
mRNA-1273 (Moderna)	4 (11)
Vakzevria (AstraZeneca)	4 (11)
Vakzevria/BNT162b2	5 (14)
Circulating B cells at vaccination, n (%)	17 (46)
Days from CAR T-cell therapy to vaccination (n = 15), median (range)	257 (115-817)
Days from vaccination to CAR T-cell therapy (n = 21), median (range)	208 (70-365)

anti-IL-6R, anti-interleukin 6 receptor (tocilizumab); Axi-cel, axicabtagene ciloleucel; CRS, cytokine release syndrome; DLBCL, diffuse large B-cell lymphoma; ICANS, immune effector cell-associated neurotoxicity syndrome; mRNA, messenger RNA; NHL, non-Hodgkin lymphoma; Tisa-cel, tisagenlecleucel.

measurements in 7 patients, 3 to 7 months after vaccination, but no patient seroconverted. Furthermore, we tested 21 patients vaccinated before undergoing lymphodepletion and CAR-T treatment. Interestingly, the humoral responses in previously vaccinated patients who received CAR-T therapy remained largely unchanged with a slight decline in antibody response compared with healthy individuals. For 2 patients, an antibody increase was observed due to a reported infection. After the exclusion of these, all 8 previously seropositive patients remained positive for neutralizing antibodies, and all 10 of 11 seronegative patients remained negative. One patient seroconverted, which might be explained by an unrecognized infection. All participants gave their informed, written consent. The study has been performed according to the Declaration of Helsinki. Sample collection and analysis were approved by the ethics committee of the Medical Center, University of Erlangen, Germany (protocol 118_20B, 174_20B, 19-336_1-B).

Although only a minority of patients developed protective antibody titers, for patients vaccinated after CAR-T therapy, most (13 of 15, 87%) developed S-specific T-cell responses (Figure 1B). Again, in

patients vaccinated before CAR-T therapy, the immune response was sustained throughout the treatment with a constant 88% (14 of 16 patients) T-cell response rate. Specificity of responses was shown for HCs by the new occurrence of these anti-S-specific T cells after vaccination. The measured T-cell response consisted of CD4 and CD8 T cells with a higher proportion of CD4 T cells in both cohorts (supplemental Figure 1). As expected, most of the convalescent donors (14 of 15, 93%) showed anti-N T-cell responses (Figure 1B). Follow-up for up to 7 months showed relatively constant immune responses.

Taken together, the presence of circulating B cells and vice versa loss of CAR T cells do not appear to be the only requirements for humoral responses. In both the groups of patients with seroconversion, patients vaccinated after CAR-T therapy and in most seropositive patients (10 of 13 patients) vaccinated before CAR-T therapy, B cells were detectable. In addition, 73% of the nonresponders of the patients vaccinated before CAR-T therapy showed circulating B cells. Moreover, we observed no differences in absolute T-cell or CD4 T-helper cell numbers in responders vs nonresponders in either of the patient cohort. No other parameter, including patient characteristics, type and timing of vaccination, IgG level, and side effects due to CAR-T therapy, seemed to have an impact on the immune responses (supplemental Table 1). We cannot exclude that missing correlations are due to the limited study population.

The low humoral response rates in CAR-T recipients are consistent with recent reports.⁴⁻⁶ In contrast, the incomplete response rates in patients vaccinated before CAR-T treatment might be explained by their underlying diseases. Recent reports describe a lack of humoral immune responses in 21% of patients with diffuse large B-cell lymphoma.⁷ However, our findings indicate that once humoral responses are present, they persist over time with CAR-T therapy at least for the period considered here. The persistence of plasma cells even 25 months after CAR-T infusion in the absence of B cells⁸ serves as a possible explanation for the maintained humoral immune response in these patients who received CAR-T therapy.⁹

Interestingly, we detected a cellular response in 87% of vaccinated patients previously exposed to CAR-T therapy. This was independent from the humoral immunity, CAR-T therapy-related complications, or circulating CAR T cells. First reports of convalescent hematologic patients describe that CD8⁺ T cells in particular lead to recovery from COVID-19 in the case of B-cell aplasia¹⁰ and confirm the importance of cellular responses. New vaccination approaches mainly addressing T-cell immunity should be considered for patients with B-cell deficiency in the future.¹¹ In addition, booster vaccinations might be beneficial for this patient population. The humoral and cellular response rate after a third vaccination increased in patients after allogeneic transplantation^{12,13} and patients treated for B-cell malignancies¹⁴ with a primary series of 3 doses and 1 additional booster dose as recommended by the American Society of Hematology and the American Society of Transplantation and Cellular Therapy for the recipients of CAR-T cells.¹⁵

The cellular immune response seems to be of particular relevance as SARS-CoV-2 mutations may escape the humoral response.^{16,17} This immune escape is most prominently displayed by the enormous mutational profile in the B.1.1.529 strain with more than 30 mutations in the S protein.¹⁸ Nevertheless, recent studies depicted cross-reactivity of T cells after vaccination or convalescence by




Figure 1. Anti-SARS-CoV-2 immune responses in patients who received CAR T-cell therapy. Patients after CAR T-cell therapy (CAR⇒VAC), before (pre), and 1 to 2 months (M1/2), and 3 to 7 months (M3-7) after SARS-CoV-2 vaccination and patients after vaccination (VAC⇒CAR), before and 1 to 2 months (M1/2), and 3 to 7 months (M3-7) after CAR T-cell therapy were analyzed for anti-SARS-CoV-2 immune responses. Convalescent patients (n = 15, open square) and HCs after vaccination (HC VAC, n = 15, circles) were used as controls. (A) Graph shows the humoral response with anti-spike IgG (left) and neutralizing antibody activity (right). (B) Cellular response to anti-spike (left) and antinucleocapsid-specific T cells (right) is graphed. Individual patients were interconnected; dotted line depicts the cutoff for positivity as indicated in the "Methods." Square symbols were used for patients after a confirmed diagnosis of SARS-CoV-2 infection. Box plots depict the 75th percentile, median, and 25th percentile values, and whiskers represent maximum and minimum values. **P* < .05, ***P* < .01, ****P* < .001. AU, arbitrary units; BAU, binding antibody units; ns, not significant.

intact responses of CD4⁺ or CD8⁺ T cells against B1.1.529,¹⁹ which further underlines the relevance of cellular response.

Although the patient number is limited, our observations showed that most patients benefit from vaccination after CAR-T therapy. This is of special interest because those patients are at a higher risk of severe COVID-19 and fatal outcomes.^{20,21} The joint recommendation of the American Society of Hematology and the American Society of Transplantation and Cellular Therapy recommends SARS-CoV-2 vaccination as early as 3 months after CAR-T therapy and a vaccine regimen including 4 administrations.¹⁵ This is in line with our findings showing effective immune responses in

short distance after CAR-T therapy. Our data suggest high and persistent response rates for patients who were vaccinated before CAR-T therapy. The relatively good tolerance of vaccination, also observed for CAR-T recipients, underlines this recommendation.^{5,6} In addition, during the follow-up, 10 patients had a positive test result for COVID-19 with only mild symptoms and fully recovered.

Taken together, our data confirm strong cellular immune responses to SARS-CoV-2 vaccination in most patients with severe B-cell depletion after CAR-T therapy. Because most patients vaccinated before CAR-T therapy showed a humoral and cellular response, our study provides evidence for vaccination before CAR-T therapy

with subsequent antibody response detection. Although we could not measure protection from infection in this cohort, these data show the necessity to look beyond antibody levels and underline the importance and meaning of vaccination in these vulnerable patients.

Acknowledgments: The authors thank all of the patients and healthy donors who participated in this study. They also thank Florentine Schonath (Core Unit Cell Sorting and Immunomonitoring Erlangen), Dorothea Gebhardt, Lina Meretuk, and Alina Bauer for excellent technical assistance.

This work was supported by grants from Sander Stiftung 2020.045.1, German Research Foundation (VO1835/3-1, MA1351/12-1), Bayerische Forschungsförderung, and Thorwart-Jeska Stiftung and support from the Else Kröner Forschungskolleg (V.B.) and the German Cancer Consortium School of Oncology fellows (V.B.).

Contribution: A.N.K., H.R., V.B., F.M., A.M., M.S., and S.V. designed the research; A.N.K., H.R., K.S., M.A., N.E., A.K., V.L., T.H., and S.V. performed experiments; M.T. and P.I. performed antibody measurements; V.B., B.J., W.R., D.M., S.K., M.L., G.K., F.M., G.S., and A.E.K. repeatedly referred patients; A.N.K., H.R., M.A., C.L., J.V.G., H.B., F.M., and S.V. analyzed and interpreted data; A.N.K., H.R., V.B., M.S., and S.V. wrote the manuscript; and all the authors edited the manuscript.

Conflict-of-interest-disclosure: The authors declare no competing financial interests.

ORCID profiles: H.R., 0000-0002-7336-2008; V.B., 0000-0003-1379-0876; C.L., 0000-0001-8876-7756; J.V.G., 0000-0002-3076-5122; A.E.K., 0000-0002-9263-948X; M.T., 0000-0003-3951-9056; T.H., 0000-0001-6945-3241; F.M., 0000-0001-5487-5839; A.M., 0000-0002-0685-4483; M.S., 0000-0001-9154-9469.

Correspondence: Simon Völkl, Department of Internal Medicine 5, Hematology and Oncology, University Hospital Erlangen, Schwabachanlage 12, 91052 Erlangen, Germany; email: simon.voelkl@uk-erlangen.de.

References

1. Dai M, Liu D, Liu M, et al. Patients with cancer appear more vulnerable to SARS-CoV-2: a multicenter study during the COVID-19 outbreak. *Cancer Discov.* 2020;10(6):783-791.
2. Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med.* 2020;383(27):2603-2615.
3. Baden LR, El Sahly HM, Essink B, et al. Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N Engl J Med.* 2021;384(5):403-416.
4. Dhakal B, Abedin S, Fenske T, et al. Response to SARS-CoV-2 vaccination in patients after hematopoietic cell transplantation and CAR T-cell therapy. *Blood.* 2021;138(14):1278-1281.
5. Ram R, Hagin D, Kikozashvili N, et al. Safety and immunogenicity of the BNT162b2 mRNA COVID-19 vaccine in patients after allogeneic HCT or CD19-based CART therapy—a single-center prospective cohort study. *Transplant Cell Ther.* 2021;27(9):788-794.
6. Gastinne T, Le Bourgeois A, Coste-Burel M, et al. Safety and antibody response after one and/or two doses of BNT162b2 anti-SARS-CoV-2 mRNA vaccine in patients treated by CAR T cells therapy. *Br J Haematol.* 2022;196(2):360-362.
7. Greenberger LM, Saltzman LA, Senefeld JW, Johnson PW, DeGennaro LJ, Nichols GL. Antibody response to SARS-CoV-2 vaccines in patients with hematologic malignancies. *Cancer Cell.* 2021;39(8):1031-1033.
8. Bhoj VG, Arhontoulis D, Wertheim G, et al. Persistence of long-lived plasma cells and humoral immunity in individuals responding to CD19-directed CAR T-cell therapy. *Blood.* 2016;128(3):360-370.
9. Walti CS, Loes AN, Shuey K, et al. Humoral immunogenicity of the seasonal influenza vaccine before and after CAR-T-cell therapy: a prospective observational study. *J Immunother Cancer.* 2021;9(10).
10. Bange EM, Han NA, Wileto P, et al. CD8(+) T cells contribute to survival in patients with COVID-19 and hematologic cancer. *Nat Med.* 2021;27(7):1280-1289.
11. Heitmann JS, Bilich T, Tandler C, et al. A COVID-19 peptide vaccine for the induction of SARS-CoV-2 T cell immunity. *Nature.* 2022;601(7894):617-622.
12. Kamar N, Abravanel F, Marion O, Couat C, Izopet J, Del Bello A. Three doses of an mRNA Covid-19 vaccine in solid-organ transplant recipients. *N Engl J Med.* 2021;385(7):661-662.
13. Hall VG, Ferreira VH, Ku T, et al. Randomized trial of a third dose of mRNA-1273 vaccine in transplant recipients. *N Engl J Med.* 2021;385(13):1244-1246.
14. Greenberger LM, Saltzman LA, Senefeld JW, Johnson PW, DeGennaro LJ, Nichols GL. Anti-spike antibody response to SARS-CoV-2 booster vaccination in patients with B cell-derived hematologic malignancies. *Cancer Cell.* 2021;39(10):1297-1299.
15. ASH-ASTCT. ASH-ASTCT COVID-19 vaccination for HCT and CAR T cell recipients: frequently asked questions. Updated March 2022. Accessed 22 March 2022. <https://www.hematology.org/covid-19/ash-astct-covid-19-vaccination-for-hct-and-car-t-cell-recipients>
16. Cele S, Gazy I, Jackson L, et al. Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent plasma. *Nature.* 2021;593(7857):142-146.
17. Zhang X, Wu S, Wu B, et al. SARS-CoV-2 Omicron strain exhibits potent capabilities for immune evasion and viral entrance. *Signal Transduct Target Ther.* 2021;6(1):430.
18. Callaway E. Heavily mutated Omicron variant puts scientists on alert. *Nature.* 2021;600(7887):21.
19. Gao Y, Cai C, Grifoni A, et al. Ancestral SARS-CoV-2-specific T cells cross-recognize the Omicron variant. *Nat Med.* 2022;28:472-476.
20. Busca A, Salmanton-Garcia J, Corradini P, et al. COVID-19 and CAR-T cells: current challenges and future directions—a report from the EPICOVIDEHA survey by EHA-IDWP. *Blood Adv.* 2021;6(7):2427-2433.
21. Hensley MK, Bain WG, Jacobs J, et al. Intractable coronavirus disease 2019 (COVID-19) and prolonged severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) replication in a chimeric antigen receptor-modified T-cell therapy recipient: a case study. *Clin Infect Dis.* 2021;73(3):e815-e821.