

TO THE EDITOR:

An agenda to advance research in myelodysplastic syndromes: a TOP 10 priority list from the first international workshop in MDS

Maximilian Stahl,¹ Omar Abdel-Wahab,² Andrew H. Wei,³ Michael R. Savona,⁴ Mina L. Xu,⁵ Zhuoer Xie,⁶ Justin Taylor,⁸ Daniel Starczynowski,⁹ Guillermo F. Sanz,¹⁰⁻¹² David A. Sallman,⁶ Valeria Santini,¹³ Gail J. Roboz,¹⁴ Mrinal M. Patnaik,⁷ Eric Padron,⁶ Olatoyosi Odenike,¹⁵ Aziz Nazha,¹⁶ Stephen D. Nimer,⁸ Ravindra Majeti,¹⁷ Richard F. Little,¹⁸ Steven Gore,¹⁸ Alan F. List,¹⁹ Vijay Kutchroo,²⁰ Rami S. Komrokji,⁶ Tae Kon Kim,⁴ Nina Kim,¹⁸ Christopher S. Hourigan,²¹ Robert P. Hasserjian,²² Stephanie Halene,²³ Elizabeth A. Griffiths,²⁴ Peter L. Greenberg,¹⁷ Maria Figueroa,⁸ Pierre Fenaux,²⁵ Fabio Effiace,²⁶ Amy E. DeZern,²⁷ Matteo G. Della Porta,²⁸ Naval G. Daver,²⁹ Jane E. Churpek,³⁰ Hetty E. Carraway,³¹ Andrew M. Brunner,³² Uma Borate,³³ John M. Bennett,³⁴ Rafael Bejar,³⁵ Jacqueline Boulton,³⁶ Sanam Loghavi,³⁷ Jan Philipp Bewersdorf,² Uwe Platzbecker,³⁸ David P. Steensma,³⁹ Mikkael A. Sekeres,⁸ Rena J. Buckstein,⁴⁰ and Amer M. Zeidan²³

¹Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA; ²Department of Medicine, Leukemia Service, Memorial Sloan Kettering Cancer Center, New York, NY; ³Peter MacCallum Cancer Centre, Royal Melbourne Hospital, University of Melbourne and Walter and Eliza Hall Institute of Medical Research, Melbourne, Australia; ⁴Division of Hematology/Oncology, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN; ⁵Departments of Pathology & Laboratory Medicine, Yale University School of Medicine and Yale Cancer Center, New Haven, CT; ⁶Department of Malignant Hematology, H. Lee Moffitt Cancer Center, Tampa, FL; ⁷Division of Hematology, Department of Internal Medicine, Mayo Clinic, Rochester, MN; ⁸Sylvester Comprehensive Cancer Center, University of Miami Miller School of Medicine, Miami, FL; ⁹Division of Experimental Hematology and Cancer Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH; ¹⁰Hematology Department, Hospital Universitario y Politécnico La Fe, Valencia, Spain; ¹¹Health Research Institute La Fe, Valencia, Spain; ¹²CIBERONC, Instituto de Salud Carlos III, Madrid, Spain; ¹³MDS Unit, AOUC, University of Florence, Florence, Italy; ¹⁴Weill Cornell Medicine and The New York Presbyterian Hospital, New York, NY; ¹⁵The University of Chicago, Chicago, IL; ¹⁶Department of Medical Oncology, Sidney Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA; ¹⁷Division of Hematology, Department of Medicine, Cancer Institute, Stanford University School of Medicine, Stanford, CA; ¹⁸National Cancer Institute, National Institutes of Health, Bethesda, MD; ¹⁹Precision BioSciences, Inc, Durham, NC; ²⁰Evergrande Center for Immunologic Diseases, Harvard Medical School, Brigham and Women's Hospital, Boston, MA; ²¹Laboratory of Myeloid Malignancies, Hematology Branch, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD; ²²Department of Pathology, Massachusetts General Hospital, Boston, MA; ²³Section of Hematology, Department of Internal Medicine, Yale University School of Medicine and Yale Cancer Center, New Haven, CT; ²⁴Roswell Park Comprehensive Cancer Center, Buffalo, NY; ²⁵Hôpital Saint Louis, Paris, France; ²⁶Italian Group for Adult Hematologic Diseases (GIMEMA), Data Center and Health Outcomes Research Unit, Rome, Italy; ²⁷The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, MD; ²⁸Humanitas Clinical and Research Center & Humanitas University, Department of Biomedical Sciences, Milan, Italy; ²⁹Department of Leukemia, The University of Texas MD Anderson Cancer Center, Houston, TX; ³⁰Department of Hematology, Oncology, and Palliative Care, Carbone Cancer Center, The University of Wisconsin-Madison, Madison, WI; ³¹Leukemia Program, Hematology and Medical Oncology, Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH; ³²Center for Leukemia, Massachusetts General Hospital, Boston, MA; ³³Division of Hematology, Department of Internal Medicine, James Cancer Center, Ohio State University, Columbus, OH; ³⁴Hematopathology Division, Departments of Pathology and Medicine, University of Rochester Medical Center, Rochester, NY; ³⁵Division of Hematology and Oncology, Moores Cancer Center, UC San Diego, La Jolla, CA; ³⁶Blood Cancer UK Molecular Haematology Unit, Nuffield Division of Clinical Laboratory Sciences, Radcliffe Department of Medicine, University of Oxford, Oxford, UK; ³⁷Department of Hematopathology, The University of Texas MD Anderson Cancer Center, Houston, TX; ³⁸Department of Hematology and Cellular Therapy, Medical Clinic and Policlinic I, Leipzig University Hospital, Leipzig, Germany; ³⁹Novartis Institutes of BioMedical Research; and ⁴⁰Odette Cancer Centre, Sunnybrook Health Sciences Centre, Toronto, Canada

Myelodysplastic syndromes (MDS) are neoplasms with high molecular, biological, and clinical heterogeneity.^{1,2} Consequently, conducting basic, translational, and clinical research on MDS has historically been challenging and the field has lagged behind in terms of achieving significant therapeutic advances.

To advance research in MDS and translate those advances into clinical benefits for patients, the first international workshop for MDS (iwMDS) was conducted in Miami, Florida. Workshop participants represented a wide variety of international stakeholders in MDS across multiple disciplines. The workshop participants concluded that there is an overarching need for international and interdisciplinary collaboration and the coordination of research efforts. Here, we outline 10 critical areas that would benefit from broad collaborations (Table 1) and propose some concrete steps for future efforts (Table 2).

Submitted 17 August 2022; accepted 14 September 2022; prepublished online on *Blood Advances* First Edition 19 October 2022; final version published online 14 June 2023. <https://doi.org/10.1182/bloodadvances.2022008747>.

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Data are available on request from the corresponding author, Amer M. Zeidan (amer.zeidan@yale.edu).

Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution.

Table 1. Top 10 list of MDS collaborative priority research goals

Priority research goals
1. To establish a new standard of care for frontline higher-risk MDS
2. To develop better treatment options for DNA methyltransferase inhibitor (DNMTi)-refractory MDS
3. To develop effective strategies for <i>TP53</i> -mutated MDS
4. To advance novel treatment strategies to impact the underlying pathophysiology of lower-risk MDS
5. To conduct clinical trials in a collaborative international effort with emphasis on equal access and on PROs
6. To formulate unified diagnostic criteria and classification subgroups for MDS
7. To establish and systematically validate clinically meaningful response criteria for MDS therapy
8. To establish tools to predict, and ultimately reduce, risk of progression of CH to MDS and other hematological malignancies in clinical practice
9. To establish linked clinical databases and biobanks allowing sharing of data
10. To improve the development and dissemination of reliable preclinical models of MDS

To establish a new standard of care for frontline higher-risk MDS

DNA methyltransferase inhibitor (DNMTi) monotherapy remains the standard of care therapy for higher-risk MDS.³ Despite the established role of DNMTi therapeutics in MDS, several trials and real-world registry analyses have failed to replicate the survival benefit initially described with azacitidine in the AZA-001 trial.^{4,5} To improve outcomes, multiple clinical trials have tested azacitidine (or DNMTi backbone) combination therapy approaches, but to date, none have improved overall survival (OS) compared with

azacitidine monotherapy.⁶⁻⁸ Hence, establishing a DNMTi backbone combination therapy for the frontline treatment of MDS is the top priority that has been identified by the faculty.

To develop better treatment options for DNMTi-refractory MDS

Outcomes for patients with DNMTi resistance are unfortunately dismal, with a median survival of <6 months and a 2-year survival probability of only 15% for patients with higher-risk MDS.⁹ Outcomes for lower-risk patients with DNMTi resistance are also poor with a median survival of 17 months.¹⁰ There is currently no standard

Table 2. Potential steps to achieve MDS priority research goals

Priority research goals	Action steps
1	- Development of new frontline treatment of MDS: Several large, randomized trials, including azacitidine in combination with the BCL-2 inhibitor venetoclax, the anti CD47 antibody magrolimab and the anti TIM3 antibody sibatolimab are ongoing
2	- Development of treatment options for DNMTi-refractory MDS: Several early phase clinical trials are ongoing; however, no therapy is approved yet at time of DNMTi resistance
3	- Develop trials with broad inclusion of both <i>TP53</i> mutated MDS and AML - Focus on trials with emphasis on engaging the immune system instead of chemotherapy based clinical trials - Utilize platform-based approaches to conduct multiple randomized phase II trials to establish a promising approach for phase III testing early (eg, NCI myeloMATCH)
4	- Palliative goals of treatment in LR-MDS are somewhat antithetical, considering that for patients with CH, a primary effort is to prevent the progression of disease. The focus of drug development should be on reversing the underlying pathophysiology of the disease. - One promising area of investigation is targeting immune dysregulation in MDS clones and the BM microenvironment (eg, targeting IRAK1/4 and/or the NLRP3 inflammasome)
5	- Expand eligibility to be representative and inclusive of the population including minority and underserved populations - Partner with pharmaceutical companies, government and regulatory agencies with a shared mission and set of goals - Inclusion of HRQoL and other type of PROs should always be considered for phase III trials to generate definite data that can facilitate clinical-decisions
6	- Formulate one single set of consensus diagnostic criteria (WHO classification has historically represented the gold standard in pathological classification)
7	- Update the International Working Group 2006 response criteria for higher-risk MDS with emphasis on association with long term benefit - Systematically and prospectively validate specific blood count cut-offs in response criteria - Eliminate response criteria without clear association with improvement in overall survival (eg, mCR without hematologic improvement)
8	- Secondary prevention of progression from CHIP to MDS in germ line predisposed, those receiving cytotoxic exposures, or acquired with age are all components of the same broader goal. - Develop practical decision-making tools to guide hematologists, medical oncologists, and patients alike in balancing the risks and benefit of adjuvant chemotherapy and radiation therapy in patients with high-risk CH
9	- Develop strategies to allow data sharing between large registries in the U.S. (National MDS Natural History Study and the Connect Myeloid Disease registry) and Europe (HARMONY Alliance and the MDS-RIGHT project) - Accomplishing these goals will require to overcome several bureaucratic and regulatory hurdles and necessitates sensitivity to the distinct privacy laws in Europe and the U.S.
10	- Utilize and widely distribute the improved in vivo modeling systems for MDS preclinical studies (eg, "MISTRG" mice)

treatment approach approved for these patients and clinical trial options are urgently needed, especially for patients who are not candidates for allogeneic hematopoietic stem cell transplantation.

To develop effective strategies for TP53-mutated MDS

TP53 alterations (deletions and mutations) are associated with extremely poor prognosis, with lower response to all established therapies, higher rates of progression to AML, and significantly shortened OS.¹¹ Even with allogeneic stem cell transplant outcomes remain poor.¹² Hence, international efforts should be focused on dedicated trials for this patient population as a holistic platform to also include *TP53*-mutated AML, which shares similar features with *TP53*-mutated MDS.¹³ During the iwMDS, the NCI myeloMATCH program was presented which will include a multiarm phase II trial platform entirely dedicated to patients with *TP53* mutated MDS. Working groups developing directed algorithms for all *TP53*-mutated myeloid neoplasms should be considered given the dismal outcome for these disorders irrespective of blast percentage.¹³

To advance novel treatment strategies to impact the underlying pathophysiology of lower-risk MDS

In lower-risk MDS, the goal of treatment is to improve cytopenia, most commonly anemia. The standard of care remains supportive treatment with red blood cell transfusions, erythropoiesis-stimulating agents, luspatercept, lenalidomide for certain subtypes of LR-MDS, and DNMTis. However, “lower-risk” MDS is a misconception: while temporary benefit can be derived, these approaches do not have a significant disease-modifying effect, and therefore responses are of limited durability.¹⁰ Future efforts should focus on attempting to reverse the underlying pathophysiology of MDS.^{14,15}

To conduct clinical trials in a collaborative international effort with emphasis on equal access and on patient-reported outcomes (PROs)

Clinical trials should be conducted jointly in different countries to achieve rapid study completion. Trials should also promote health care equity for patients with MDS and focus on the inclusion of PROs. Collaboration among research groups worldwide will facilitate these research goals. In addition, the focus should be on optimizing the inclusivity of patients concerning eligibility criteria to encompass the older population of patients (eg, limited eligibility due to creatinine and or cardiac functions) that are currently frequently excluded, thereby hampering the investigation (and enrollment) of the population most afflicted with MDS.

To formulate unified diagnostic criteria for MDS

Recently the World Health Organization and the International Consensus Classification for myeloid neoplasms have published separate diagnostic criteria for AML and MDS.^{16,17} Several

important differences between the 2 classifications exist, especially in how patients with 10% to 19% blasts are classified.¹⁸ There is a significant concern that having 2 separate classification systems could have a negative impact on clinical care and research and have the potential to create confusion among patients and providers, as well as in the design, conduct, and interpretation of clinical trials. The need to establish a unified consensus classification for MDS has been emphasized.

To establish and systematically validate clinically meaningful response criteria for MDS therapy

The current response MDS criteria suffer from significant limitations.^{19,20} First, there is a clear inconsistency in how complete remission (CR) is defined for MDS based on the International Working Group 2006²⁰ compared with the definition of CR for AML in the ELN 2017²¹ and ELN 2022²² criteria. While no hemoglobin threshold is used for CR in AML, for MDS, CR requires a hemoglobin value ≥ 11 g/DL, which has not been validated. These differences in CR definition can lead to different outcomes depending on which definition is used.²³ There is a need to develop and validate new response criteria for higher-risk MDS that capture meaningful clinical benefits, similar to the introduction of CR with partial hematological recovery in ELN 2022 AML criteria.^{22,24,25} Conversely, the elimination of responses with doubtful clinical benefits (eg, marrow CR without meaningful count recovery) should be considered.²⁶

To establish tools to predict, and ultimately reduce, risk of progression of clonal hematopoiesis (CH) to MDS and other hematological malignancies in clinical practice

The advent and widespread availability of next generation sequencing has led to the discovery of CH, which, in some cases, is associated with higher risks of cardiovascular disease²⁷ and hematologic malignancies.²⁸⁻³⁰ Therefore, early intervention should be monitored and considered, especially among patients with solid tumors where high-risk CH clones can dramatically expand under the selective pressure of chemotherapy, increasing the risk of subsequent therapy-related myeloid neoplasms. We need to learn more about the genotypic and phenotypic characteristics of CH evolution to accurately predict which patients with CH will develop MDS and other blood cancers. Ultimately, the goal is to develop effective secondary preventive/therapeutic strategies that reduce the risk of aggressive hematologic malignancies and/or other CH-related diseases in patients with higher-risk CH lesions. The identification of CH before exposure to intensive chemotherapy or radiation therapy is not frequently covered by insurance companies, which is a barrier to obtaining adequate informed consent and poses risks to patients when deciding to embark on potentially toxic therapies. Lastly, how best to identify and classify patients with a germ line predisposition to develop myeloid malignancies such as MDS and AML, will be a significant future focus of genetic counseling and risk prevention efforts.³¹

To establish linked clinical databases and biobanks allowing sharing of data

Several large MDS registries have recently been developed to better understand the natural history of MDS and to establish large biorepositories. The global MDS community should establish a common minimum data set to enable data sharing across platforms. Achieving international alignment is essential for identifying and characterizing risk factors, clinical outcomes, and particularly drug responsiveness for subgroups of this rare disease, which are likely to be increasingly subdivided into more defined molecular subtypes.

To improve the development and dissemination of reliable preclinical models of MDS

Comprehensive *in vivo* preclinical studies of MDS have historically suffered from the limited ability of MDS stem cells to engraft in immunodeficient murine hosts. Recently, a patient-derived xenotransplantation model in cytokine-humanized immunodeficient “MISTRG” mice has been established, allowing disease modeling across MDS subtypes.³² A broader availability of improved preclinical models and wider dissemination of these resources to laboratories around the world is critical to maximize the efficient evaluation of novel drugs and therapeutics in MDS.

We conclude that, given that MDS are rare and biologically heterogeneous blood cancers, there is an overarching dire need for international and interdisciplinary collaboration and coordination of research efforts. What became clear from this first iwMDS meeting was the necessity for the frequent continued exchange of ideas in a smaller informal setting that is more conducive to the creation of new collaborative efforts.

Acknowledgments: A.M.Z. is a Leukemia and Lymphoma Society Scholar in Clinical Research. M.E.F. is a Leukemia and Lymphoma Society Scholar. T.K.K. received the Clinician Scientist Development Award from the American Cancer Society. The authors acknowledge Kelly Norsworthy for her critical review and comments on the manuscript.

Contribution: M.S. and A.M.Z. wrote the initial draft of the manuscript; M.S., A.M.Z., R.B., M.A.S., D.P.S., U.P., S.L., A.H.W., and V.S. were involved with the conception and design of the study; and all authors were involved in writing, reviewing, and editing the manuscript, and approved the final version for submission.

Conflict-of-interest disclosure: The first international workshop on MDS was planned by an independent scientific steering committee. The meeting was supported by unrestricted educational grants from Gilead, Novartis, Bristol Myers Squibb, Syros, Geron, and Karyopharm to Magdalen Medical/VJHemOnc. The participants met in Miami, FL, in June 2022, and deliberated on the content of this manuscript, which was subsequently finalized after the meeting. The meeting sponsors were not involved in the development of this manuscript. A.N., D.P.S., R.B., A.F.L., and N.K. authored this manuscript in their personal capacity, and their views do not necessarily represent/reflect that of their employers. The content is the sole responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. M.M.P., E.P., U.P., R.B., and S.L. were authors of the WHO

classification. R.P.H. was a cochair of the Pathology ICC and co-leader of the ICC Premalignant Clonal Cytopenias and MDS working group. O.O., M.R.S., and A.H.W. were the authors of the ICC classification. A.M.Z. received research funding (institutional) from Celgene/Bristol Myers Squibb, AbbVie, Astex, Pfizer, Medimmune/AstraZeneca, Boehringer-Ingelheim, Cardiff Oncology, Incyte, Takeda, Novartis, Aprea, and ADC Therapeutics. A.M.Z. participated in advisory boards, and/or had a consultancy with and received honoraria from AbbVie, Otsuka, Pfizer, Celgene/Bristol Myers Squibb, Jazz, Incyte, Agios, Boehringer-Ingelheim, Novartis, Acceleron, Astellas, Daiichi Sankyo, Cardinal Health, Taiho, Seattle Genetics, BeyondSpring, Cardiff Oncology, Takeda, Ionis, Amgen, Janssen, Epizyme, Syndax, Gilead, Kura, Chiesi, ALX Oncology, BioCryst, Notable, Orum, and Tyme, and served on the clinical trial committees for Novartis, AbbVie, Gilead, BioCryst, ALX Oncology, Geron, and Celgene/Bristol Myers Squibb. D.P.S. is an employee of Novartis. U.P. received honoraria from Bristol Myers Squibb, Jazz Pharmaceuticals, AbbVie, and Geron. The National Heart, Lung, and Blood Institute received research funding for the laboratory of C.S.K. from Sellas and the Foundation of the NIH AML M.R.D. Biomarkers Consortium. F.E. had a consultancy or advisory role for Amgen, AbbVie, Janssen, and Novartis, and received research funding (institutional) from AbbVie, Amgen, and Novartis, outside the submitted work. O.O. has served on the advisory board of Bristol Myers Squibb, Celgene, Novartis, Taiho, and Kymera Therapeutics; serves on a DSMB for Threadwell Therapeutics; and received research funding (institutional) from AbbVie, Agios, Aprea, Astex, AstraZeneca, Bristol Myers Squibb, Celgene, CTI, Daiichi, Incyte, Janssen, Kartos, Novartis, NS-Pharma, and Oncotherapy Sciences. P.L.G. received research funding (institutional) from Celgene/Bristol Myers Squibb and Gilead. P.F. received research funding from Bristol Myers Squibb, AbbVie, Jazz Pharmaceuticals, Novartis, and Janssen, and had consultancy with and received honoraria from Bristol Myers Squibb, AbbVie, Jazz Pharmaceuticals, and Novartis. M.S. consulted for Curis Oncology and Boston Consulting; served on the advisory boards for Novartis and Kymera; and participated in the GME activity for Novartis, Curis Oncology, Haymarket Media, and Clinical Care Options (CCO). G.F.S. received grants/research support (institutional) from Bristol Myers Squibb, Novartis, and Gamida Cell. R.M. is on the Board of Directors of CircBio; on the advisory boards of Kodikaz Therapeutic Solutions, Syros Pharmaceuticals, and TenSixteen Bio; is an inventor on a number of patents related to CD47 cancer immunotherapy licensed to Gilead Sciences; received research support from the Gilead Sciences and CircBio; and is the cofounder and equity holder of CircBio, Pheast Therapeutics, MyeloGene, and RNAC Therapeutics. G.F.S. received honoraria, advisory board membership, or consultation fees from AbbVie, Bristol Myers Squibb, Novartis, Roche, and Takeda, and participated in the sponsored speaker's bureau for Bristol Myers Squibb, Novartis, and Takeda. A.N. owns stock at Amazon and is an employee of Incyte Pharmaceuticals. D.T.S. is a consultant and received research funding from Kymera Therapeutics, Kurome Therapeutics, Captor Therapeutics, and Tolero Therapeutics. D.T.S. has equity in Kurome Therapeutics. D.A.S. served on the advisory boards of Aprea, AvenCell, Bluebird Bio, Bristol Myers Squibb, Intellia, Kite, Novartis, Shattuck Labs, Servier, and Syndax; served as a consultant for AbbVie, Magenta, Molecular Partners AG, and Takeda; served on the speakers' bureau for Bristol Myers Squibb, Incyte, and Servier; and received research funding from Aprea and

Jazz Pharmaceuticals. M.M.P. received research funding from Kura Oncology and StemLine Pharmaceuticals. A.M.B. received consulting or advisory board honoraria from Novartis, Acceleron, Agios, AbbVie, Takeda, Celgene/Bristol Myers Squibb, Keros Therapeutics, Taiho, and Gilead, and has research support from NIH SPORE in Myeloid Malignancies and the Edward P. Evans Foundation. E.P. receives research funding from Incyte and Bristol Myers Squibb, and honoraria from Stemline, Taiho, Blueprint, and Bristol Myers Squibb. T.K.K. received research funding from Nextcure and is a consultant for the Agenus. R.S.K. served on the speaker bureau of Jazz Pharmaceuticals, Servio, CTI, and PharmaEssentia, and served on advisory boards and received honoraria from Bristol Myers Squibb, Novartis, AbbVie, Jazz Pharmaceuticals, Servio, PharmaEssentia, Taiho, Geron, and CTI. M.S.A. has served on advisory boards for Bristol Myers Squibb, Novartis, Kurome, and Gilead. R.B. is employed by and has equity in Aptose Biosciences, and serves on independent drug monitoring committees for Gilead and Epizyme. R.P.H. has a consultancy agreement with Bluebird Bio. G.J.R. has a consultancy agreement and/or serves on the advisory board or data and safety monitoring committee for AbbVie, Amgen, Astellas, AstraZeneca, Bristol Myers Squibb, Blueprint Medicines, Bluebird Bio, Celgene, Glaxo SmithKline, Janssen, Jasper Therapeutics, Jazz Pharmaceuticals, Mesoblast, Novartis, Pfizer, Syndax, and Takeda, and received research support from Janssen. S.H. received honoraria from Forma Therapeutics. A.H.W. has served on advisory boards for Novartis, AstraZeneca, Astellas, Janssen, Amgen, Roche, Pfizer, AbbVie, Servier, Gilead, Bristol Myers Squibb, Shoreline, MacroGenics, and Agios; receives research funding (to institution) from Novartis, AbbVie, Servier, Janssen, Bristol Myers Squibb, Syndax, Astex, AstraZeneca, and Amgen; serves on speakers' bureaus for AbbVie, Novartis, Bristol Myers Squibb, Servier, and Astellas; is an employee of the Walter and Eliza Hall Institute (WEHI); and is eligible for financial benefits associated with payments that WEHI receives in relation to venetoclax. A.F.L. is employed by and has equity in Precision BioSciences, and has served as a consultant for Halia Therapeutics, CTI Biopharma, and Aileron. M.X. participated in advisory boards and/or had a consultancy agreement with and received honoraria from Seattle Genetics, Pure Marrow and Blueprint Medicines. N.G.D. has received research funding from Daiichi-Sankyo, Bristol Myers Squibb, Pfizer, Gilead, Sevier, Genentech, Astellas, Daiichi-Sankyo, AbbVie, Hanmi, Trovogene, Fate Therapeutics, Amgen, Novimmune, Glycomimetics, Trillium, and ImmunoGen, and has served in a consulting or advisory role for Daiichi-Sankyo, Bristol Myers Squibb, Arog, Pfizer, Novartis, Jazz, Celgene, AbbVie, Astellas, Genentech, ImmunoGen, Servier, Syndax, Trillium, Gilead, Amgen, Shattuck Labs, and Agios. S.L. received research support from Astellas and Amgen; owns stock from AbbVie; and has received consultancy fees/honoraria from AbbVie, Gerson Lehrman Group, QualWorld, and Guidepoint. E.A.G. received honoraria from AbbVie, Alexion Pharmaceuticals, Genentech, Novartis, CTI Biopharma, Apellis, Celgene/Bristol Myers Squibb, Takeda Oncology, Taiho Oncology, Physician Educational Resource, MediCom Worldwide, American Society of Hematology, Picnic Health, and AAMDSIF, and research support (institutional) from Astex Pharmaceuticals, Genentech, Blueprint Medicine, Alexion Pharmaceuticals, Apellis, Bristol Myers Squibb/Celgene, and Celldex Therapeutics. M.R.S. serves on a board or advisory committee for AbbVie, Bristol Myers Squibb, CTI, Forma, Geron, Karyopharm, Novartis, Ryvu, Sierra Oncology, Taiho, Takeda, and TG

Therapeutics; receives research funding from ALX Oncology, Astex, Incyte, Takeda, and TG Therapeutics; and has equity ownership in Karyopharm and Ryvu. O.A.W. has served as a consultant for H3B Biomedicine, Foundation Medicine Inc., Merck, Prelude Therapeutics, and Janssen; is on the scientific advisory board of Envisagenics Inc., AlChem, Harmonic Discovery Inc., and Pfizer Boulder; and has received prior research funding from H3B Biomedicine and LOXO Oncology, unrelated to the manuscript. The remaining authors declare no competing financial interests.

ORCID profiles: A.H.W., 0000-0002-7514-3298; M.R.S., 0000-0003-3763-5504; Z.X., 0000-0003-1266-9804; J.T., 0000-0003-4407-6325; G.F.S., 0000-0002-2767-8191; V.S., 0000-0002-5439-2172; G.J.R., 0000-0002-0384-3658; M.M.P., 0000-0001-6998-662X; R.F.L., 0000-0002-4091-1699; R.S.K., 0000-0002-1876-5269; T.K.K., 0000-0002-5182-3870; N.K., 0000-0002-1809-0417; C.S.H., 0000-0002-6189-8067; S.H., 0000-0002-2737-9810; E.A.G., 0000-0002-0288-8248; P.L.G., 0000-0002-7615-9443; N.G.D., 0000-0001-7103-373X; H.E.C., 0000-0001-5241-3614; R.B., 0000-0002-5603-4598; J.B., 0000-0002-4330-2928; S.L., 0000-0001-8980-3202; U.P., 0000-0003-1863-3239; D.P.S., 0000-0001-5130-9284.

Correspondence: Amer M. Zeidan, Hematology Section, Department of Internal Medicine, Yale School of Medicine, 333 Cedar St, PO Box 208028, New Haven, CT 06520-8028; email: amer.zeidan@yale.edu.

References

1. Cogle CR, Kurtin SE, Bentley TG, et al. The incidence and health care resource burden of the myelodysplastic syndromes in patients in whom first-line hypomethylating agents fail. *Oncologist*. 2017;22(4):379-385.
2. Papaemmanuil E, Gerstung M, Malcovati L, et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*. 2013;122(22):3616-3627; quiz 3699.
3. Fenaux P, Mufti GJ, Hellström-Lindberg E. Azacitidine prolongs overall survival compared with conventional care regimens in elderly patients with low bone marrow blast count acute myeloid leukemia. *J Clin Oncol*. 2010;28(4):562-569.
4. Zeidan AM, Stahl M, Sekeres MA, Steensma DP, Komrokji RS, Gore SD. A call for action: increasing enrollment of untreated patients with higher-risk myelodysplastic syndromes in first-line clinical trials. *Cancer*. 2017;123(19):3662-3672.
5. Zeidan AM, Salimi T, Epstein RS. Real-world use and outcomes of hypomethylating agent therapy in higher-risk myelodysplastic syndromes: why are we not achieving the promise of clinical trials? *Future Oncol*. 2021;17(36):5163-5175.
6. Sekeres MA, Othus M, List AF, et al. Randomized phase II study of azacitidine alone or in combination with lenalidomide or with vorinostat in higher-risk myelodysplastic syndromes and chronic myelomonocytic leukemia: north american intergroup study SWOG S1117. *J Clin Oncol*. 2017;35(24):2745-2753.
7. Adès L, Girshova L, Doronin VA, et al. Pevonedistat plus azacitidine vs azacitidine alone in higher-risk MDS/chronic myelomonocytic leukemia or low-blast-percentage AML. *Blood Adv*. 2022;6(17):5132-5145.
8. Therapeutics A. Aprea Therapeutics Announces Results of Primary Endpoint from Phase 3 Trial of Eprenetapopt in TP53 Mutant Myelodysplastic Syndromes (MDS). *Journal of Clinical Oncology*. 2020.

9. Prebet T, Gore SD, Esterni B, et al. Outcome of high-risk myelodysplastic syndrome after azacitidine treatment failure. *J Clin Oncol*. 2011;29(24):3322-3327.
10. Jabbour E, Garcia-Manero G, Batty N, et al. Outcome of patients with myelodysplastic syndrome after failure of decitabine therapy. *Cancer*. 2010;116(16):3830-3834.
11. Sallman DA, Komrokji R, Vaupel C, et al. Impact of TP53 mutation variant allele frequency on phenotype and outcomes in myelodysplastic syndromes. *Leukemia*. 2016;30(3):666-673.
12. Lindsley RC, Saber W, Mar BG, et al. Prognostic mutations in myelodysplastic syndrome after stem-cell transplantation. *N Engl J Med*. 2017;376(6):536-547.
13. Grob T, Al Hinai ASA, Sanders MA, et al. Molecular characterization of mutant TP53 acute myeloid leukemia and high-risk myelodysplastic syndrome. *Blood*. 2022;139(15):2347-2354.
14. Smith MA, Choudhary GS, Pellagatti A, et al. U2AF1 mutations induce oncogenic IRAK4 isoforms and activate innate immune pathways in myeloid malignancies. *Nat Cell Biol*. 2019;21(5):640-650.
15. Basiorka AA, McGraw KL, Eksioglu EA, et al. The NLRP3 inflammasome functions as a driver of the myelodysplastic syndrome phenotype. *Blood*. 2016;128(25):2960-2975.
16. Khoury JD, Solary E, Abal O, et al. The 5th edition of the World Health Organization classification of haematolymphoid tumours: myeloid and histiocytic/dendritic neoplasms. *Leukemia*. 2022;36(7):1703-1719.
17. Arber DA, Orazi A, Hasserjian RP, et al. International consensus classification of myeloid neoplasms and acute leukemia: integrating morphological, clinical, and genomic data. *Blood*. 2022;140(11):1200-1228.
18. Estey E, Hasserjian RP, Döhner H. Distinguishing AML from MDS: a fixed blast percentage may no longer be optimal. *Blood*. 2022;139(3):323-332.
19. Cheson BD, Bennett JM, Kantarjian H, et al. Report of an international working group to standardize response criteria for myelodysplastic syndromes. *Blood*. 2000;96(12):3671-3674.
20. Cheson BD, Greenberg PL, Bennett JM, et al. Clinical application and proposal for modification of the International Working Group (IWG) response criteria in myelodysplasia. *Blood*. 2006;108(2):419-425.
21. Döhner H, Estey E, Grimwade D. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017;129(4):424-447.
22. Döhner H, Wei AH, Appelbaum FR, et al. Diagnosis and management of AML in adults: 2022 ELN recommendations from an international expert panel. *Blood*. 2022;140(12):1345-1377.
23. Peterlin P, Turlure P, Chevallier P, et al. Conference procedure issue. *Blood*. 2021;138:243.
24. Shallis RM, Pollyea DA, Zeidan AM. The complete story of less than complete responses: the evolution and application of acute myeloid leukemia clinical responses. *Blood Rev*. 2021;48:100806.
25. Brunner AM, Gavralidis A, Ali NA, et al. Evaluating complete remission with partial hematologic recovery (CRh) as a response criterion in myelodysplastic syndromes (MDS). *Blood Cancer J*. 2022;12(11):153.
26. Komrokji RS, Al Ali NH, Sallman D, et al. Validation of International Working Group response criteria in higher-risk myelodysplastic syndromes: a report on behalf of the MDS Clinical Research Consortium. *Cancer Med*. 2021;10(2):447-453.
27. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26):2488-2498.
28. Malcovati L, Galli A, Travaglino E, et al. Clinical significance of somatic mutation in unexplained blood cytopenia. *Blood*. 2017;129(25):3371-3378.
29. Galli A, Todisco G, Catamo E, et al. Relationship between clone metrics and clinical outcome in clonal cytopenia. *Blood*. 2021;138(11):965-976.
30. Bolton KL, Ptashkin RN, Gao T, et al. Cancer therapy shapes the fitness landscape of clonal hematopoiesis. *Nat Genet*. 2020;52(11):1219-1226.
31. Kico JM, Mullighan CG. Advances in germline predisposition to acute leukaemias and myeloid neoplasms. *Nat Rev Cancer*. 2021;21(2):122-137.
32. Song Y, Rongvaux A, Taylor A, et al. A highly efficient and faithful MDS patient-derived xenotransplantation model for pre-clinical studies. *Nat Commun*. 2019;10(1):366.