

Gata2 noncoding genetic variation as a determinant of hematopoietic stem/progenitor cell mobilization efficiency

Alexandra A. Soukup and Emery H. Bresnick

Wisconsin Blood Cancer Research Institute, Department of Cell and Regenerative Biology, Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, Madison, WI

Key Points

- Noncoding *Gata2* genetic variation reduces efficacy of single and multiagent clinical regimens to mobilize HSPCs.
- Noncoding *Gata2* genetic variation abrogates polyvinyl alcohol-induced hematopoietic stem/progenitor cell expansion.

Germline genetic variants alter the coding and enhancer sequences of *GATA2*, which encodes a master regulator of hematopoiesis. The conserved murine *Gata2* enhancer (+9.5) promotes hematopoietic stem cell (HSC) genesis during embryogenesis. Heterozygosity for a single-nucleotide Ets motif variant in the human enhancer creates a bone marrow failure and acute myeloid leukemia predisposition termed GATA2 deficiency syndrome. The homozygous murine variant attenuates chemotherapy- and transplantation-induced hematopoietic regeneration, hematopoietic stem and progenitor cell (HSPC) response to inflammation, and HSPC mobilization with the therapeutic mobilizer granulocyte colony-stimulating factor (G-CSF). Because a *Gata2* +9.5 variant attenuated G-CSF-induced HSPC expansion and mobilization, and HSC transplantation therapies require efficacious mobilization, we tested whether variation affects mechanistically distinct mobilizers or only those operating through select pathways. In addition to affecting G-CSF activity, *Gata2* variation compromised IL-8/CXCR2- and VLA-4/VCAM1-induced mobilization. Although the variation did not disrupt HSPC mobilization mediated by plerixafor, which functions through CXCR4/CXCL12, homozygous and heterozygous variation attenuated mobilization efficacy of the clinically used plerixafor/G-CSF combination. The influence of noncoding variation on HSPC mobilization efficacy and function is important clinically because comprehensive noncoding variation is not commonly analyzed in patients. Furthermore, our mobilization-defective system offers unique utility for elucidating fundamental HSPC mechanisms.

Introduction

To obtain hematopoietic stem and progenitor cells (HSPCs) for transplantation, granulocyte colony-stimulating factor (G-CSF) is used to mobilize HSPCs into peripheral blood. Low levels of mobilization¹⁻⁴ have been attributed to aging, history of ineffective mobilization, diabetes, prior chemotherapy, and low baseline bone marrow (BM) cellularity/CD34⁺ cell numbers.⁵ Poor mobilization in diabetes is associated with hyperglycemia⁶ and sympathetic denervation in the hematopoietic system.⁷ Sympathetic agonists increase mobilization.⁸ Genetic variation in patients with Fanconi anemia⁹ is associated with poor mobilization and decreased HSPCs in the marrow,^{10,11} although the mechanisms are unresolved.

Submitted 21 June 2023; accepted 13 October 2023; republished online on *Blood Advances* First Edition 23 October 2023; final version published online 15 December 2023. <https://doi.org/10.1182/bloodadvances.2023011003>.

All data generated and analyzed during this study are included in this article and its supplemental information files. Original data and further information on resources and reagents are available upon reasonable request from the corresponding author, Emery Bresnick (ehbresni@wisc.edu).

The full-text version of this article contains a data supplement.

© 2023 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

Mechanisms mediating G-CSF-induced HSPC mobilization involve interplay among HSPCs, neutrophils, macrophages, and nonhematopoietic BM niche components.¹² Disruption of the HSPC-expressed CXCR4 and niche-secreted SDF1 (CXCL12) HSC homing mechanism with plerixafor (AMD3100 and Mozobil)^{10,13} results in HSPC egress from the niche into peripheral blood. The plerixafor/G-CSF combination is leveraged clinically to increase CD34⁺ cell numbers. Disruption of the axis involving HSPC VLA-4 and niche VCAM1 with the VLA-4 antagonist BIO5192 or other inhibitors promotes mobilization.¹⁴⁻¹⁶ Inflammatory signals, including CXCL2 (Gro-beta; Mip2) and CXCL8 (IL-8), activate neutrophils and mobilize HSPCs.¹⁷⁻¹⁹ Although mice lack IL-8, CXCR2 activation with human (rh)IL-8, Gro-beta or truncated Gro-beta induces mobilization.^{18,20,21}

We reasoned that variation in genes causing hematologic pathology predispositions may compromise mobilization. Germline variants alter coding and enhancer sequences of *GATA2*,²² which encodes a master regulator of hematopoiesis.²³ The conserved murine *Gata2* enhancer (+9.5) promotes hematopoietic stem cell (HSC) genesis during embryogenesis.²⁴⁻²⁶ Heterozygosity for a single-nucleotide Ets motif variant in the human enhancer creates a BM failure and acute myeloid leukemia predisposition termed GATA2 deficiency syndrome.²⁷ The homozygous murine variant (+9.5(Ets)^{-/-}) attenuates chemotherapy- and transplantation-induced hematopoietic regeneration and HSPC response to inflammation.²⁸ Biallelic +9.5 enhancer variation²⁵ (compound heterozygous; CH) blocks mobilization by multidose G-CSF administration.²⁹ This system involves an allele harboring the +9.5 Ets motif variant and another with an inactivating enhancer deletion (Figure 1A). This “conditionally pathogenic” configuration creates a BM failure predisposition in mice²⁹ and models humans in which epigenetic silencing of the normal allele exacerbates the impact of a germline predisposition variant.³⁰ We asked if *Gata2* variation interferes with all, or select, mobilization mechanisms and if heterozygosity, reflecting the human predisposition state, compromises mobilization.

Methods

Age- and sex-matched mice were used for all experiments. Protocols were approved by the UW-Madison Institutional Animal Care and Use Committee in accordance with Association for Assessment and Accreditation of Laboratory Animal Care International regulations. rhG-CSF (Zarxio, 125 µg/kg) was administered subcutaneously as a single dose or twice a day (8 doses) beginning in the evening of the first day. Mice were harvested 3 hours after the final dose. For alternative mobilizers, plerixafor 5 mg/kg was administered intraperitoneally (IP), BIO5192 (R&D systems, 5051/10) 1 mg/kg was administered IV, and rhIL-8 (Peprotech, 200-08M) 1 mg/kg was injected IP. Mice were harvested after 15 minutes, 1 hour, or 2 hours. Colony forming unit (CFU) assays, transcript analysis, and flow cytometry were performed as described.²⁹ 5-ethynyl-2'-deoxyuridine (EdU); (Sigma, 900584-50 mg) 500 µg was injected IP 4 hours before harvest. EdU incorporation was detected using Click-iT EdU Alexa Fluor 647 Flow Cytometry Assay Kit (ThermoFisher, C10419) with modifications.³¹ Annexin V binding was detected using Alexa Fluor 647 Annexin V (BioLegend, 640943) as described.³² Complete blood counts were performed using an Oxford Science Inc GENESIS.

Adult C57BL/6 recipient mice (CD45.1⁺, 6- to 8-week-old; Stock # 002014, Jackson Labs) were lethally irradiated using an XRAD 320 irradiator for a single dose of 8 Gy. Competitor BM cells were harvested from individual 8-week-old mice (CD45.1⁺). A total of 10⁶ BM cells were mixed with 50 mL of peripheral blood (6 total per group; 2 experiments). Transplant-recipient mice were maintained on Irradiated Uniprim Diet (Envigo; Cat# TD.06596) for 2 weeks. Blood obtained from the retro-orbital venous sinus was isolated after transplantation and analyzed using flow cytometry for donor-derived hematopoiesis.²⁹

For HSC culture, BM from 8–12-week-old age- and sex- matched mice was isolated in phosphate-buffered saline. Fifty Lin⁻ Sca1⁺ Kit⁺ (LSK) CD34⁻ CD48⁻ CD150⁺ cells were isolated using fluorescence-activated cell sorter and plated in the central wells of fibronectin-coated 96-well plates (Corning #354409) containing 0.2 mL polyvinyl alcohol (PVA) medium (Ham F-12 Nutrient Mix liquid medium (Gibco #11765-054) with 10 mM HEPES (Gibco #15630-080), 1× penicillin-streptomycin-Glutamine (Gibco #10378-016), insulin-transferrin-selenium-ethanolamine (Gibco #51500-056), 100 ng/mL recombinant animal-free murine thrombopoietin (Peprotech #AF-315-14), and 10 ng/mL recombinant animal-free murine stem cell factor (Peprotech #AF-250-03).^{33,34} PVA (1 mg/mL; 87%-90%-hydrolyzed; Sigma, #P8136) was included as indicated. Medium changes were initiated 5 days after plating and repeated every 2 to 3 days.

Group sizes required for each experiment followed the National Academies recommended calculation to detect a *P* value. The results are presented as box and whisker plots, with bounds from the 25th to the 75th percentiles, the median line, and whiskers ranging from minimum to maximum values or as individual values summarized by median and standard error. Alternatively, individual data points were plotted with median and standard error values. Multiple independent cohorts were used in each experiment. Statistical comparisons were performed using 2-tailed Student *t* tests (significance cutoff of *P* < .05), with correction of statistical overrepresentation of functions calculated using the Benjamini-Hochberg multiple tests correction procedure or 1-way analysis of variance (ANOVA) followed by Tukey multiple comparisons test. GraphPad Prism (GraphPad Software) was used to perform statistical analyses.

Results

Short-term G-CSF treatment induces HSPC metabolic gene signatures,³⁵ cell cycle entry, and expansion.^{36,37} Because CH have decreased multipotent progenitors (MPPs; LSK CD150⁻ CD48⁻) and common myeloid progenitors (Lin⁻ Sca1⁻ Kit⁺ FcγR⁻ CD34⁺),²⁹ we asked if mobilization defects reflected decreased HSPCs in BM and if *Gata2* variation affects HSPC proliferation before circulation. Wild-type (WT) BM immunophenotypic HSCs expanded 1.8- (*P* = .01) and 1.4-fold (*P* = .03) at 8 and 16 hours, respectively (supplemental Figure 1). No population frequency changes were detected at 4 hours (Figures 1B-D; supplemental Figure 1). We assessed whether CH were competent to expand under these conditions. At 16 hours, WT HSCs (LSK CD48⁻ CD150⁺; *P* = .001) and LSK CD48⁻ CD150⁻ MPPs (*P* = .01) expanded 1.8-fold relative to vehicle treatment; CH HSCs and MPPs were unchanged in response to G-CSF. As seen previously, CH MPPs were 3.5-fold lower than WT in vehicle-treated samples




Figure 1.

and 5.8-fold lower than WT after 4 hours of G-CSF. G-CSF expanded the multipotent LSK CD48⁺ population 3.5-fold in WT without affecting the heterogeneous Lin⁻Kit⁺Sca1⁻ population. CH variation abrogated expansion of immunophenotypic HSPC expansion after G-CSF treatment.

We asked if increased *Gata2* expression in HSPC populations is associated with HSPC expansion in response to G-CSF. *Gata2* expression in WT and CH long-term (LT)-HSCs was comparable in the steady state²⁹ or after G-CSF (Figure 2A). CH variation impaired *Gata2* expression in MPPs in a G-CSF-independent manner. *Gata2* expression was 2.5-fold lower ($P = .003$) in vehicle-treated CH MPP vs WT and 3.3-fold lower ($P = .037$) in CH MPP vs WT 16 hours after G-CSF treatment (Figure 2A). Because the LSK CD48⁻ CD150⁻ MPP pool contains additional MPP subtypes, including MPP5 (LSK CD48⁻ CD150⁻ CD34⁺ Flt3⁻) and MPP6 (LSK CD48⁻ CD150⁻ CD34⁻ Flt3⁻), we asked whether the decreased LSK CD48⁻ CD150⁻ population corresponded to MPP5, MPP6, or both. MPP5 and MPP6 were 1.7-fold ($P = .0001$ and $.003$) lower in untreated animals (supplemental Figure 2).

We asked if decreased immunophenotypic populations resulted in reduced functional activity in BM and peripheral blood. CH had 1.8-fold fewer CFU than WT BM in the steady state ($P = .007$) and 1.5-fold fewer 16 hours after G-CSF ($P = .017$), reflecting a paucity of functional HSPCs. No differences in colony counts from peripheral blood were detected, confirming that HSPCs have not egressed from the BM³⁸ (Figure 2B). Thus, CH mice exhibit decreased HSPCs and failed to rapidly elevate HSPCs after G-CSF administration.

We asked if defective expansion resulted from inability of CH HSPCs to enter the cell cycle. No increases were detected at 4 hours. By contrast, 16 hours after G-CSF administration, WT BM EdU⁺ (S phase) HSCs increased 2.8-fold (Figure 2C). CH HSCs had a 1.7-fold ($P = .04$) higher percentage of cells in S phase and 1.9-fold ($P = .04$) higher percentage 4 hours after G-CSF. At 16 hours, no differences were detected. WT and CH LSK CD48⁺ cell numbers in S phase did not differ at any time, although both genotypes exhibited greater than twofold increases at 16 hours (supplemental Figure 3A). No differences were detected with Lin⁻Kit⁺Sca1⁻ progenitors. Cycling CH MPPs were 2.7-fold higher than WT in the steady state, 2.2-fold higher at 4 hours, and comparable by 16 hours (Figure 2C).

Although immunophenotypic CH HSCs and MPPs exhibited increased cycling, their numbers decreased relative to WT. This did not reflect ectopic mobilization as peripheral blood CFU in the steady state or 16 hours after treatment was unaltered (Figure 2B). We asked if CH variation rendered these populations apoptotic.^{32,39} Although Annexin V⁺ CH HSCs increased 1.8-fold relative to WT ($P = .002$) 16 hours after G-CSF administration, there was no increase with vehicle alone. CH MPP apoptosis increased (1.6- to 3.1-fold vs WT MPP) ($P = .02$, $.04$, and $.02$ with

vehicle, 4 hours, and 16 hours G-CSF, respectively) regardless of treatment (Figure 2D; supplemental Figure 3B). GATA2 was therefore required for MPP population homeostasis.

We tested whether CH HSC expansion defects could be overcome by PVA, which supports expansion of HSPCs *ex vivo*.³³ Defined serum-free medium containing PVA expands purified HSPCs (LSK CD150⁺ CD34⁻) more than 100-fold over a month culture.³³ We asked if WT and CH LT-HSCs (LSK CD150⁺ CD34⁻ CD48⁻) had similar growth potential over 15 days (Figure 3A-B). Fifty cells were seeded per well. By day 15, WT cultures had a mean of 4286 cells per well, an 86-fold expansion (Figure 3C). In contrast, CH cultures had an average of 396 cells/well, a less than eightfold ($P = .0072$ vs WT) expansion. Under these conditions, the number of WT immunophenotypic LT-HSCs (LSK CD150⁺ CD34⁻ CD48⁻) increased fivefold, CH LT-HSCs decreased to an average of 10 cells per well ($P = .0007$ vs WT) (Figure 3C). Neither culture expanded in medium lacking PVA. Thus, GATA2 is required for HSPC expansion *ex vivo* and *in vivo*.

To analyze relationships between mobilization and expansion defects, we used mobilization strategies involving distinct mechanisms. We asked if *Gata2* variation attenuated mobilization via disrupting the CXCR4/CXCL12 chemokine axis with the CXCR4 antagonist plerixafor, stromal/neutrophil CXCR2 activation with rhIL-8, or VLA-4/VCAM1 binding with VLA-4 inhibitor BIO5192. Although the mobilization kinetics differed slightly, maximal mobilization by these mechanisms occurs within 1 hour, before complex BM remodeling.^{10,14,18} To ensure that the differences in mobilization did not reflect differences in mobilization kinetics between genotypes, we analyzed mobilization at 3 times.

In contrast to CH mobilization with G-CSF, plerixafor increased peripheral blood CFU in WT 3.3- to 4.1-fold at all times analyzed. CH CFU increased 3.4- and 4.1-fold at 1- and 2-hours after G-CSF administration, respectively (Figure 4A-B). CH mobilization was not decreased relative to WT at any time, demonstrating that the CXCR4/CXCL12 mobilization axis is intact in CH. Maximal rhIL-8-mediated mobilization in WT was achieved by 1 hour (3.9-fold over vehicle for WT). CH mobilization was not increased at any time analyzed and was significantly lower ($P = .0003$) than WT at the 1-hour time. Although VLA-4 inhibition with BIO5192 increased WT CFU maximally at 15 minutes (5.5-fold relative to vehicle), CH CFUs were significantly lower at 15 minutes ($P = .016$) and did not increase relative to vehicle treatment. Individual agents increased neutrophil numbers in peripheral blood in WT.^{18,40,41} Consistent with CFU analyses, CH responded to plerixafor, but not IL-8 or BIO5192 (Figure 4C). Although the CXCR4/CXCL12 pathway was insensitive to *Gata2* variation, mobilization by means of CXCR2 activation or VLA-4 inhibition was impaired.

G-CSF treatment of WT mice decreases CXCL12 levels in the BM niche, increases neutrophil activation, and decreases VCAM1/VLA-4 interaction.⁴²⁻⁴⁴ We tested whether targeting these pathways collectively with G-CSF restores mobilization in CH. Synergistic

Figure 1. *Gata2* germline variation blocks HSPC expansion after acute G-CSF treatment. (A) Configuration of *Gata2* variant alleles in WT and CH. (B) Mice were treated with 1 dose of 125 mg/kg rhG-CSF or vehicle and harvested after 4 or 16 hours. (C) Representative flow cytometric plots for BM HSPC quantification. The numbers represent the percentage of the parent cell population. N = 6 to 9 per condition, 4 experiments (D) BM HSPC quantification. HSC: Lin⁻ Sca1⁺ Kit⁺ (LSK) CD48⁻ CD150⁺; MPP: LSK CD48⁻ CD150⁻; LSK CD48⁺; Lin⁻ Sca1⁻ c-Kit⁺. * $P < .05$, ** $P < .01$, *** $P < .001$; 1-way ANOVA followed by Tukey multiple comparisons test. FSC, forward scatter; PBS, phosphate-buffered saline.




Figure 2. G-CSF-independent and -dependent CH HSPC defects. (A) Messenger RNA quantification from FACS-isolated HSC (LSK CD150⁺ CD48⁻) or MPP (LSK CD150⁻ CD48⁻). N = 3 to 6 per condition from 3 experiments. (B) CFU from 2 × 10⁵ total BM cells or 0.1 mL of peripheral blood. N = 3 to 4 per condition from 3 experiments.




Figure 3. *Gata2* germline variation prevents polyvinyl alcohol-supported HSPC expansion ex vivo. (A) Schematic for isolation and culture of LT-HSCs in defined medium.³⁴ (B) Representative flow cytometric plots for FACS-isolation of LT-HSCs (LSK CD150⁺ CD48⁻ CD34⁻). (C) Flow cytometric quantification of total live cells and LT-HSCs (LSK CD150⁺ CD48⁻ CD34⁻) per well. N = 3 replicates per condition, average of 3 experiments. * $P < .05$, ** $P < .01$, *** $P < .001$; 1-way ANOVA followed by Tukey multiple comparisons test. DAPI, 4',6-diamidino-2-phenylindole.

Figure 2 (continued) * $P < .05$, ** $P < .01$; 1-way ANOVA followed by Tukey multiple comparisons test. (C) Top: EdU incorporation to quantify cell entry into S phase within HSCs and MPPs. Bottom: representative flow cytometric plots. N = 4 to 7 per condition, 5 experiments. (C) Top: annexin V binding to quantify HSC and MPP apoptosis. Bottom: representative flow cytometric plots. N = 4 to 7 per condition, 5 experiments. * $P < .05$, ** $P < .01$, *** $P < .001$; 2-tailed unpaired Student *t* test with Benjamini-Hochberg correction. DAPI, 4',6-diamidino-2-phenylindole; FACS, fluorescence-activated cell sorter.




Figure 4. *Gata2* germline variation selectively attenuates acute HSPC mobilization. (A) Mice were treated with 5 mg/kg plerixafor (CXCR4 antagonist), 1 mg/kg rhIL-8 (CXCR2 agonist), or 1 mg/kg BIO5192 (VLA-4 antagonist) and harvested after 0.25, 1, or 2 hours. (B) CFU from 0.1 mL of peripheral blood. (C) Peripheral blood neutrophil quantification. N = 3 to 12 per condition, 9 experiments. * $P < .05$, ** $P < .01$, *** $P < .001$; 1-way ANOVA followed by Tukey multiple comparisons test. PBS, phosphate-buffered saline.

mobilization was detected in WT with dual treatments of G-CSF and plerixafor, IL-8, or BIO5192 (2.6-, 2.3-, and 5.8-fold increases in CFU; $P = .0002$, $.0007$, and $.0001$, respectively) relative to G-CSF alone (Figure 5A-B). The CH variant reduced CFU, circulating neutrophils, and splenic HSC frequencies (Figure 5A-D). We tested whether these deficits resulted in decreased repopulating activity. Competitive transplantations with peripheral blood revealed that G-CSF + plerixafor increased WT donor-derived hematopoiesis 24-, 20-, and 18-fold at 8, 12, and 16 weeks, respectively, relative to the vehicle-treated control (Figure 5E). Treatment of CH did not increase donor-derived contributions. CH mice lacked the rapid HSPC expansion after G-CSF administration, yielding fewer cells to mobilize. Normal mobilization was not achieved even with plerixafor, which was insensitive to CH variation in the single-dose paradigm. The G-CSF + plerixafor mobilizing regimen did not elevate CH repopulating activity.

Because GATA2 deficiency syndrome and somatic GATA2 variation involve heterozygous alleles, we tested whether monoallelic

Ets motif variation^{28,29} disrupts mobilization. Monoallelic variation did not decrease circulating, steady-state HSPCs. G-CSF increased CFU 28.5-fold ($P < .0001$) relative to vehicle-treated WT mice (Figure 5F-G). Mobilization of *Gata2 Ets*^{+/-} HSPCs were reduced significantly ($P = .003$). CH variation severely disrupted mobilization, with 7.3-fold fewer CFU than *Gata2 Ets*^{+/-} ($P = .02$). Synergistic G-CSF/plerixafor treatment increased CFU 89-fold in WT vs untreated ($P < .0001$), and *Gata2 Ets*^{+/-} mobilization was reduced ($P = .006$) and not improved relative to *Gata2 Ets*^{-/-} or CH. *Ets*^{+/-} mobilization with G-CSF/BIO5192 was also impaired. Thus, *Gata2* heterozygosity, modeling a human pathological state, impaired G-CSF-induced HSPC mobilization with all regimens tested.

Discussion

Considering the complex cell-intrinsic and -extrinsic mechanisms governing HSPC mobilization, genetic variation in genes encoding a plethora of components is almost certainly an important




Figure 5.




Figure 6. GATA2 genetic variation reduces HSPC activity and mobilization efficiency. Before treatment, WT quiescent HSPCs are maintained in BM through retention mechanisms including CXCR4/CXCL12, VLA-4/VCAM1, and low inflammation (IL-8). During mobilization, these mechanisms are disrupted, leading to HSPC expansion and circulation. *Gata2* variation decreases HSPC levels, reducing the available HSPCs to mobilize and compromises VLA-4/VCAM1- and IL-8–induced mobilization mechanisms.

determinant of mobilization efficiency. The many unanswered questions include: is variation in genes encoding cell-intrinsic regulators of HSPC development and function an important determinant of mobilization efficiency? Although mobilization differs among inbred mouse strains¹⁰ and genetic background influences mobilization efficiency,⁴⁵ the mechanisms are not established. In patients, failure to mobilize has been attributed to age, prior failure to mobilize, disease (eg, diabetes), prior chemotherapy, and low baseline BM cellularity and/or CD34⁺ cell numbers.⁵ Heritability estimations suggested that genetic variation plays a role in determining circulating CD34⁺ levels,⁴⁶ and a genome-wide association study analysis demonstrated that *PPM1H*, *CXCR4*, *ENO1-RERE*, *ITGA9*, *ARHGAP45*, *CEBPA*, *TERT*, and *MYC* are associated with steady-state levels of circulating CD34⁺.⁴⁷ Although *CXCR4* variation may dictate CD34⁺ levels in the steady state, the relationship between *CXCR4* polymorphism and mobilization efficiency is unclear.^{48,49}

Prior research has focused on the role of genetic variation within genes encoding the receptors involved in the mobilization mechanisms analyzed herein. Not all genetic variation alters mobilization efficiency, even when occurring in genes encoding established regulators of mobilization. In most cases, *CXCL12* single-nucleotide polymorphisms are not associated with CD34⁺ yield,^{48,50,51} although a 3' variant is associated with high yield.⁵² As with *CXCR4*, *VCAM1* polymorphisms can contribute or have no impact on mobilization efficiency.^{48,53} Given the millions of single-nucleotide polymorphisms in the human genome, there are many unknowns regarding the broader impact of genetic variation on mobilization efficiency.

It is instructive to consider variation in genes encoding cell-intrinsic regulators of HSPC development and function. Our results

established *Gata2* variation as a vital determinant of mobilization efficiency. Although patients with pathogenic *GATA2* variants display altered chemotaxis in response to CXCL12,⁵⁴ we demonstrated that the CXCR4/CXCL12 axis is intact in CH, unlike other mobilization regimens tested. Lymphoma, multiple myeloma, and Fanconi anemia are associated with inadequate mobilization.^{1,9,55,56} *Fancc*^{-/-} and *Fanca*^{-/-} HSPCs retain the capacity to mobilize in response to plerixafor and Rac inhibition,^{10,57,58} although whether the mechanism that impairs mobilization is identical to that used in CH is unclear. As with CH, these patients frequently have low HSPC levels and/or dysfunctional HSPCs. Because proinflammatory cytokine levels correlate with mobilization,⁵⁹ and *GATA2* alterations can dysregulate cytokine production,^{60,61} it seems reasonable that *GATA2* alterations generate HSPC intrinsic and extrinsic defects that compromise *GATA2*-dependent mobilization.

Our mobilization-defective system (Figure 6) offers unique utility for elucidating fundamental HSPC mechanisms. It will be instructive to evaluate noncoding variation genome-wide with patients exhibiting suboptimal mobilization and evaluate a spectrum of variants linked to BM failure to establish rules, which may unveil innovations to surmount mobilization defects and pinpoint contexts in which specific variants preclude mobilization.

Acknowledgments

The work was supported by National Institutes of Health (NIH) grant DK68634 (E.H.B.), Edward P. Evans Foundation (E.H.B.), 1U01CA257666-01A1 (E.H.B.), ASH Scholar Award (A.A.S.), Leukemia and Lymphoma Society Career Development Program (A.A.S.), Carbone Cancer Center P30CA014520, and NIH shared instrumentation grant 1S10RR025483-01.

Figure 5. Gata2 germline variation attenuates combinatorial HSPC mobilization. (A) Mice were given 8 doses of 125 mg/kg rhG-CSF over 5 days followed by vehicle, 5 mg/kg plerixafor, 1 mg/kg rhIL-8, or 1 mg/kg BIO5192 and harvested after 1 hour. (B) CFU from 0.1 mL of peripheral blood. (C) Peripheral blood neutrophil quantification. (D) Splenic LT-HSC (LSK CD150⁺ CD48⁻) quantification. N = 3 to 10 per condition, 7 experiments. **P* < .05, ***P* < .01, ****P* < .001; 2-tailed unpaired Student *t* test with Benjamini-Hochberg correction. (E) Repopulating activity in peripheral blood 16 weeks after competitive transplant of 50 μ L peripheral blood with 10⁶ BM competitor cells. N = 5 to 6 per condition from 2 experiments. **P* < .05, ****P* < .001; 1-way ANOVA followed by Tukey multiple comparisons test. (F-G) Monoallelic Ets motif variation attenuates mobilization. WT, monoallelic (Ets^{+/+}), homozygous biallelic (Ets^{-/-}), and CH variants were tested. CFU from 0.1 mL of peripheral blood. N = 3 to 6 per condition, 6 experiments. **P* < .05, ***P* < .01, ****P* < .001; 1-way ANOVA followed by Tukey multiple comparisons test. PBS, phosphate-buffered saline.

Authorship

Contribution: A.A.S. and E.H.B. conceived and designed the research, and wrote and edited the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: A.A.S., [0000-0002-4149-2393](https://orcid.org/0000-0002-4149-2393); E.H.B., [0000-0002-1151-5654](https://orcid.org/0000-0002-1151-5654).

Correspondence: Emery H. Bresnick, Wisconsin Blood Cancer Research Institute, Department of Cell and Regenerative Biology, Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, 4009 WIMR, 1111 Highland Ave, Madison, WI 53705; email: ehbresni@wisc.edu.

References

1. DiPersio JF, Stadtmauer EA, Nademanee A, et al. Plerixafor and G-CSF versus placebo and G-CSF to mobilize hematopoietic stem cells for autologous stem cell transplantation in patients with multiple myeloma. *Blood*. 2009;113(23):5720-5726.
2. Gertz MA, Wolf RC, Micallef IN, Gastineau DA. Clinical impact and resource utilization after stem cell mobilization failure in patients with multiple myeloma and lymphoma. *Bone Marrow Transplant*. 2010;45(9):1396-1403.
3. Wuchter P, Ran D, Bruckner T, et al. Poor mobilization of hematopoietic stem cells—definitions, incidence, risk factors, and impact on outcome of autologous transplantation. *Biol Blood Marrow Transplant*. 2010;16(4):490-499.
4. Lee KH, Jung SK, Kim SJ, et al. Incidence and risk factors of poor mobilization in adult autologous peripheral blood stem cell transplantation: a single-centre experience. *Vox Sang*. 2014;107(4):407-415.
5. Hopman RK, DiPersio JF. Advances in stem cell mobilization. *Blood Rev*. 2014;28(1):31-40.
6. Ferraro F, Lymeri S, Méndez-Ferrer S, et al. Diabetes impairs hematopoietic stem cell mobilization by altering niche function. *Sci Transl Med*. 2011;3(104):104ra101.
7. Katayama Y, Battista M, Kao WM, et al. Signals from the sympathetic nervous system regulate hematopoietic stem cell egress from bone marrow. *Cell*. 2006;124(2):407-421.
8. Gao X, Zhang D, Xu C, Li H, Caron KM, Frenette PS. Nociceptive nerves regulate haematopoietic stem cell mobilization. *Nature*. 2021;589(7843):591-596.
9. Croop JM, Cooper R, Fernandez C, et al. Mobilization and collection of peripheral blood CD34+ cells from patients with Fanconi anemia. *Blood*. 2001;98(10):2917-2921.
10. Broxmeyer HE, Orschell CM, Clapp DW, et al. Rapid mobilization of murine and human hematopoietic stem and progenitor cells with AMD3100, a CXCR4 antagonist. *J Exp Med*. 2005;201(8):1307-1318.
11. Broxmeyer HE, Capitano ML, Cooper S, Potchanant ES, Clapp DW. Numbers of long-term hematopoietic stem cells from bone marrow of fanca and fancd knock-out mice can be greatly enhanced by their collection and processing in physioxia conditions. *Blood Cells Mol Dis*. 2021;86:102492.
12. Tay J, Levesque JP, Winkler IG. Cellular players of hematopoietic stem cell mobilization in the bone marrow niche. *Int J Hematol*. 2017;105(2):129-140.
13. Liles WC, Broxmeyer HE, Rodger E, et al. Mobilization of hematopoietic progenitor cells in healthy volunteers by AMD3100, a CXCR4 antagonist. *Blood*. 2003;102(8):2728-2730.
14. Ramirez P, Rettig MP, Uy GL, et al. BIO5192, a small molecule inhibitor of VLA-4, mobilizes hematopoietic stem and progenitor cells. *Blood*. 2009;114(7):1340-1343.
15. Karpova D, Rettig MP, Ritchey J, et al. Targeting VLA4 integrin and CXCR2 mobilizes serially repopulating hematopoietic stem cells. *J Clin Invest*. 2019;129(7):2745-2759.
16. Papayannopoulou T, Nakamoto B. Peripheralization of hemopoietic progenitors in primates treated with anti-VLA4 integrin. *Proc Natl Acad Sci U S A*. 1993;90(20):9374-9378.
17. Fukuda S, Bian H, King AG, Pelus LM. The chemokine GRObeta mobilizes early hematopoietic stem cells characterized by enhanced homing and engraftment. *Blood*. 2007;110(3):860-869.
18. Laterveer L, Lindley IJ, Hamilton MS, Willemze R, Fibbe WE. Interleukin-8 induces rapid mobilization of hematopoietic stem cells with radioprotective capacity and long-term myelolymphoid repopulating ability. *Blood*. 1995;85(8):2269-2275.
19. Fibbe WE, Pruijt JF, Velders GA, et al. Biology of IL-8-induced stem cell mobilization. *Ann N Y Acad Sci*. 1999;872:71-82.
20. Pruijt JF, Verzaal P, van Os R, et al. Neutrophils are indispensable for hematopoietic stem cell mobilization induced by interleukin-8 in mice. *Proc Natl Acad Sci U S A*. 2002;99(9):6228-6233.
21. Hoggatt J, Singh P, Tate TA, et al. Rapid mobilization reveals a highly engraftable hematopoietic stem cell. *Cell*. 2018;172(1-2):191-204.e10.
22. Bresnick EH, Jung MM, Katsumura KR. Human GATA2 mutations and hematologic disease: how many paths to pathogenesis? *Blood Adv*. 2020;4(18):4584-4592.
23. Tsai FY, Keller G, Kuo FC, et al. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature*. 1994;371(6494):221-226.
24. Gao X, Johnson KD, Chang YI, et al. Gata2 cis-element is required for hematopoietic stem cell generation in the mammalian embryo. *J Exp Med*. 2013;210(13):2833-2842.

25. Johnson KD, Hsu AP, Ryu MJ, et al. Cis-element mutated in GATA2-dependent immunodeficiency governs hematopoiesis and vascular integrity. *J Clin Invest.* 2012;122(10):3692-3704.
26. Mehta C, Johnson KD, Gao X, et al. Integrating enhancer mechanisms to establish a hierarchical blood development program. *Cell Rep.* 2017;20(12):2966-2979.
27. Hsu AP, Johnson KD, Falcone EL, et al. GATA2 haploinsufficiency caused by mutations in a conserved intronic element leads to MonoMAC syndrome. *Blood.* 2013;121(19):3830-3837, S1-7.
28. Soukup AA, Zheng Y, Mehta C, et al. Single-nucleotide human disease mutation inactivates a blood-regenerative GATA2 enhancer. *J Clin Invest.* 2019;129(3):1180-1192.
29. Soukup AA, Matson DR, Liu P, Johnson KD, Bresnick EH. Conditionally pathogenic genetic variants of a hematopoietic disease-suppressing enhancer. *Sci Adv.* 2021;7(50):eabk3521.
30. Al Seraihi AF, Rio-Machin A, Tawana K, et al. GATA2 monoallelic expression underlies reduced penetrance in inherited GATA2-mutated MDS/AML. *Leukemia.* 2018;32(11):2502-2507.
31. Jalbert E, Pietras EM. Analysis of murine hematopoietic stem cell proliferation during inflammation. *Methods Mol Biol.* 2018;1686:183-200.
32. Mehta C, Fraga de Andrade I, Matson DR, Dewey CN, Bresnick EH. RNA-regulatory exosome complex confers cellular survival to promote erythropoiesis. *Nucleic Acids Res.* 2021;49(16):9007-9025.
33. Wilkinson AC, Ishida R, Kikuchi M, et al. Long-term ex vivo haematopoietic-stem-cell expansion allows nonconditioned transplantation. *Nature.* 2019;571(7763):117-121.
34. Wilkinson AC, Ishida R, Nakauchi H, Yamazaki S. Long-term ex vivo expansion of mouse hematopoietic stem cells. *Nat Protoc.* 2020;15(2):628-648.
35. Fast EM, Sporrij A, Manning M, et al. External signals regulate continuous transcriptional states in hematopoietic stem cells. *Elife.* 2021;10:e66512.
36. Schuettelpelz LG, Borgerding JN, Christopher MJ, et al. G-CSF regulates hematopoietic stem cell activity, in part, through activation of Toll-like receptor signaling. *Leukemia.* 2014;28(9):1851-1860.
37. Wilson A, Laurenti E, Oser G, et al. Hematopoietic stem cells reversibly switch from dormancy to self-renewal during homeostasis and repair. *Cell.* 2008;135(6):1118-1129.
38. Winkler IG, Wiercinska E, Barbier V, Nowlan B, Bonig H, Levesque JP. Mobilization of hematopoietic stem cells with highest self-renewal by G-CSF precedes clonogenic cell mobilization peak. *Exp Hematol.* 2016;44(4):303-314.e1.
39. You X, Zhou Y, Chang YI, et al. Gata2 +9.5 enhancer regulates adult hematopoietic stem cell self-renewal and T-cell development. *Blood Adv.* 2022;6(4):1095-1099.
40. Kubes P, Jutila M, Payne D. Therapeutic potential of inhibiting leukocyte rolling in ischemia/reperfusion. *J Clin Invest.* 1995;95(6):2510-2519.
41. Petty JM, Lenox CC, Weiss DJ, Poynter ME, Suratt BT. Crosstalk between CXCR4/stromal derived factor-1 and VLA-4/VCAM-1 pathways regulates neutrophil retention in the bone marrow. *J Immunol.* 2009;182(1):604-612.
42. Falanga A, Marchetti M, Evangelista V, et al. Neutrophil activation and hemostatic changes in healthy donors receiving granulocyte colony-stimulating factor. *Blood.* 1999;93(8):2506-2514.
43. Semerad CL, Christopher MJ, Liu F, et al. G-CSF potently inhibits osteoblast activity and CXCL12 mRNA expression in the bone marrow. *Blood.* 2005;106(9):3020-3027.
44. Bellucci R, De Propriis MS, Buccisano F, et al. Modulation of VLA-4 and L-selectin expression on normal CD34+ cells during mobilization with G-CSF. *Bone Marrow Transplant.* 1999;23(1):1-8.
45. Roberts AW, Foote S, Alexander WS, Scott C, Robb L, Metcalf D. Genetic influences determining progenitor cell mobilization and leukocytosis induced by granulocyte colony-stimulating factor. *Blood.* 1997;89(8):2736-2744.
46. Cohen KS, Cheng S, Larson MG, et al. Circulating CD34(+) progenitor cell frequency is associated with clinical and genetic factors. *Blood.* 2013;121(8):e50-56.
47. Lopez de Lapuente Portilla A, Ekdahl L, Cafaro C, et al. Genome-wide association study on 13 167 individuals identifies regulators of blood CD34+cell levels. *Blood.* 2022;139(11):1659-1669.
48. Garciaz S, Sfumato P, Granata A, et al. Analysis of a large single institution cohort of related donors fails to detect a relation between SDF1/CXCR4 or VCAM/VLA4 genetic polymorphisms and the level of hematopoietic progenitor cell mobilization in response to G-CSF. *PLoS One.* 2020;15(3):e0228878.
49. Martin-Antonio B, Carmona M, Falantes J, et al. Impact of constitutional polymorphisms in VCAM1 and CD44 on CD34+ cell collection yield after administration of granulocyte colony-stimulating factor to healthy donors. *Haematologica.* 2011;96(1):102-109.
50. Schulz M, Karpova D, Spohn G, et al. Variant rs1801157 in the 3'UTR of SDF1-1B does not explain variability of healthy-donor G-CSF responsiveness. *PLoS One.* 2015;10(3):e0121859.
51. Lenk J, Bornhauser M, Kramer M, et al. Sex and body mass index but not CXCL12 801 G/A polymorphism determine the efficacy of hematopoietic cell mobilization: a study in healthy volunteer donors. *Biol Blood Marrow Transplant.* 2013;19(10):1517-1521.
52. Benboubker L, Watier H, Carion A, et al. Association between the SDF1-3'A allele and high levels of CD34(+) progenitor cells mobilized into peripheral blood in humans. *Br J Haematol.* 2001;113(1):247-250.

53. Mishima S, Matsuda C, Ishihara T, Nagase M, Taketani T, Nagai A. Single nucleotide polymorphisms of the DGKB and VCAM1 genes are associated with granulocyte colony stimulating factor-mediated peripheral blood stem cell mobilization. *Transfus Apher Sci.* 2017;56(2):154-159.
54. Maciejewski-Duval A, Meuris F, Bignon A, et al. Altered chemotactic response to CXCL12 in patients carrying GATA2 mutations. *J Leukoc Biol.* 2016; 99(6):1065-1076.
55. DiPersio JF, Micallef IN, Stiff PJ, et al. Phase III prospective randomized double-blind placebo-controlled trial of plerixafor plus granulocyte colony-stimulating factor compared with placebo plus granulocyte colony-stimulating factor for autologous stem-cell mobilization and transplantation for patients with non-Hodgkin's lymphoma. *J Clin Oncol.* 2009;27(28):4767-4773.
56. Giralt S, Costa L, Schriber J, et al. Optimizing autologous stem cell mobilization strategies to improve patient outcomes: consensus guidelines and recommendations. *Biol Blood Marrow Transplant.* 2014;20(3):295-308.
57. Pulliam AC, Hobson MJ, Ciccone SL, et al. AMD3100 synergizes with G-CSF to mobilize repopulating stem cells in Fanconi anemia knockout mice. *Exp Hematol.* 2008;36(9):1084-1090.
58. Milsom MD, Lee AW, Zheng Y, Cancelas JA. *Fanca*^{-/-} hematopoietic stem cells demonstrate a mobilization defect which can be overcome by administration of the Rac inhibitor NSC23766. *Haematologica.* 2009;94(7):1011-1015.
59. Wang TF, Liou YS, Chang HH, et al. Correlation of body mass index and proinflammatory cytokine levels with hematopoietic stem cell mobilization. *J Clin Med.* 2022;11(14):4169.
60. Takai J, Shimada T, Nakamura T, Engel JD, Moriguchi T. *Gata2* heterozygous mutant mice exhibit reduced inflammatory responses and impaired bacterial clearance. *iScience.* 2021;24(8):102836.
61. Tran VL, Liu P, Katsumura KR, et al. Restricting genomic actions of innate immune mediators on fetal hematopoietic progenitor cells. *iScience.* 2023; 26(4):106297.