

Phillips NS, Mulrooney DA, Williams AM, et al. Neurocognitive impairment associated with chronic morbidity in long-term survivors of Hodgkin Lymphoma. *Blood Adv.* 2023;7(23):7270-7278.

Page 7273: in **Table 1** under "Survivors N = 204," in the row "Diagnosed with second malignancy," the value should read "48 (24.0)," not "51 (25.0)." The corrected **Table 1** is shown below:

**Table 1. Characteristics of survivors of HL and community controls**

	<b>Survivors N = 204</b>	<b>Controls N = 205</b>	<b>P value</b>
	<b>n (%)</b>	<b>n (%)</b>	
<b>Sex</b>			
Male	97 (47.5)	99 (48.3)	0.88
Female	107 (52.5)	106 (51.7)	
<b>Race</b>			
White	172 (84.3)	176 (85.9)	0.66
Other	32 (15.7)	29 (14.1)	
<b>Smoking Status</b>			
Current	39 (19.1)	29 (14.3)	0.37
Former	35 (17.2)	41 (20.2)	
Never	130 (63.7)	133 (65.5)	
<b>Body mass index</b>			
Underweight	7 (3.4)	3 (1.5)	0.23
Normal weight	51 (25.0)	62 (30.2)	
Overweight	73 (35.8)	60 (29.3)	
Obese	73 (35.8)	80 (39.0)	
Age at assessment (y)	Mean (SD)		
	36.6 (8.01)	36.7 (9.17)	0.84
	Median (min-max)		
Age at diagnosis (y)	14.7 (3.0-22.6)		
Time since diagnosis (y)	20.6 (10.7-45.6)		
Anthracyclines (n = 156) mg/m <sup>2</sup>	157.2 (91.0-400.0)*		
IV methotrexate (n = 85) mg/m <sup>2</sup>	123.2 (20.3-167.7)		
Bleomycin (n = 58) mg/m <sup>2</sup>	35.6 (5.0-95.0)		
Corticosteroids (n = 140) mg/m <sup>2</sup>	2240 (250-4480)		
Vincristine (n=182) mg/m <sup>2</sup>	45.9 (9.2-151.2)		
Cyclophosphamides (n=113) mg/m <sup>2</sup>	3655 (1332-22 889)†		
Chest Radiation (n=204) Gy	26.0 (15.0-45.0)		
Neck Radiation (n=204) Gy	26.0 (15.0-51.0)		
	N (%)		
Relapsed	14 (6.9)		
Diagnosed with second malignancy	48 (24.0)		

Max, maximum; min, minimum; SD, standard deviation.

\*Anthracycline equivalent dose.

†Cyclophosphamide equivalent dose.

Page 7274: in **Table 2** under the “Survivors Mean (SD)” and “Controls Mean (SD)” columns, the authors misreported the means and standard deviations of the neurocognitive test at the hundredth decimal place. The corrected **Table 2** is shown below:

**Table 2. Mean age-adjusted neurocognitive z scores of survivors and controls**

Domain	Survivors	Controls	FDR P
	Mean (SD)	Mean (SD)	
<b>Attention</b>			
Focused	0.35 (0.95)	0.54 (0.84)	0.061
Span	0.04 (1.01)	0.22 (0.92)	0.085
Sustained	-0.10 (1.35)	0.23 (0.89)	<b>0.011</b>
Variability	-0.23 (1.11)	-0.03 (1.05)	0.089
<b>Processing Speed</b>			
Reaction time	-0.16 (1.17)	-0.04 (0.99)	0.29
Visuomotor	0.13 (0.96)	0.40 (0.94)	<b>0.011</b>
Visual	0.27 (0.98)	0.62 (0.97)	<b>0.002</b>
Motor	-0.38 (1.05)	-0.01 (0.90)	<b>0.001</b>
<b>Memory</b>			
Verbal learning	-0.08 (1.04)	0.31 (1.02)	<b>0.001</b>
Short-term recall	-0.24 (1.01)	0.12 (1.04)	<b>0.002</b>
Long-term recall	-0.30 (1.14)	0.08 (1.10)	<b>0.002</b>
Selective reminding	-0.18 (1.11)	-0.03 (0.96)	0.17
<b>Executive Function</b>			
Working memory	-0.08 (0.93)	0.05 (0.86)	0.16
Self-monitoring	-0.00 (1.14)	0.20 (0.95)	0.085
Cognitive flexibility	-0.02 (1.26)	0.26 (1.11)	<b>0.033</b>
Fluency	-0.05 (1.20)	0.03 (1.05)	0.47

Means and SDs presented as age-adjusted z scores, with population mean = 0 and SD = 1.0.

FDR, *P*, false discovery rate–adjusted *P*-values for multiple comparisons between survivors and control; SD, standard deviation.

These errors do not impact the *P* values, statistical significance, or relevant interpretations of the data.

<https://doi.org/10.1182/bloodadvances.2024013111>

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](#), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.