

Risk of infectious adverse events of venetoclax therapy for hematologic malignancies: a systematic review and meta-analysis of RCTs

Connor Prosty,¹ Khaled Katergi,² Alex Nguyen,¹ Owen Dan Luo,¹ Mark Sorin,¹ Vladimir Cherniak,³ Michael Sebag,⁴ Koray Demir,⁵ Emily G. McDonald,^{3,6,7} Todd C. Lee,⁶⁻⁸ and Matthew P. Cheng⁶⁻⁸

¹Faculty of Medicine, McGill University, Montréal, QC, Canada; ²Faculty of Medicine, Université de Montréal, Montréal, QC, Canada; ³Division of General Internal Medicine, Department of Medicine, McGill University Health Centre, Montréal, QC, Canada; ⁴Division of Hematology, Department of Medicine, McGill University, Montréal, QC, Canada; ⁵Division of Infectious Diseases, Department of Medicine, Université de Montréal, Montréal, QC, Canada; ⁶Division of Experimental Medicine, Department of Medicine, McGill University, Montréal, QC, Canada; and ⁷Clinical Practice Assessment Unit and ⁸Division of Infectious Diseases, Department of Medicine, McGill University Health Centre, Montréal, QC, Canada

Venetoclax is a small molecule inhibitor of BCL-2 used in the treatment of acute myelogenous leukemia (AML) and chronic lymphocytic leukemia (CLL). Recent postmarketing studies of ibrutinib, another small molecule inhibitor, suggested that these agents may predispose to opportunistic infections. We sought to systematically review the randomized controlled trial (RCT) evidence of venetoclax to assess whether it predisposes patients to infectious adverse events (IAEs) and neutropenia. We systematically reviewed RCTs comparing venetoclax therapy with active or placebo controls for patients with hematologic malignancies. Data on IAEs and neutropenia were pooled by Bayesian meta-analysis, and we computed the probability of any increased risk ($P[\text{risk ratio (RR)} > 1]$) of IAEs or neutropenic complications. Seven RCTs were included, comprising 2067 patients. In CLL ($n = 1032$), there was a low probability of increased risk of high-grade ($P[\text{RR} > 1] = 71.2\%$) and fatal IAEs ($P[\text{RR} > 1] = 64.5\%$) and high-grade neutropenia ($P[\text{RR} > 1] = 63.4\%$). There were insufficient data to perform a meta-analysis of IAEs in AML; however, 1 trial suggested an increased risk of IAEs with venetoclax. Furthermore, in AML ($n = 642$), venetoclax was associated with a high probability of increased risk of high-grade neutropenia ($P[\text{RR} > 1] = 94.6\%$) and febrile neutropenia ($P[\text{RR} > 1] = 90.6\%$). Our results suggest that venetoclax has a low probability of increased risk of IAEs or neutropenia in CLL. By contrast, there is likely increased risk of high-grade neutropenia and febrile neutropenia in AML. Importantly, our analyses did not identify any specific IAEs that would benefit from routine antimicrobial prophylaxis or pre-emptive testing.

Introduction

Infections remain a significant cause of morbidity and mortality among patients with hematologic malignancies.¹ Various mechanisms contribute to the increased risk of infectious complications in this population including underlying disease activity, immune dysregulation, and iatrogenic

Submitted 19 October 2023; accepted 13 December 2023; prepublished online on *Blood Advances* First Edition 28 December 2023; final version published online 14 February 2024. <https://doi.org/10.1182/bloodadvances.2023011964>.

Data are available on request from the corresponding author, Connor Prosty (connor.prosty@mail.mcgill.ca).

The full-text version of this article contains a data supplement.

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

immunosuppression.² Certain drugs used in the treatment of hematologic malignancies carry a disproportionate risk of opportunistic infections (OIs), such as rituximab and corticosteroids, which are associated with an increased risk of hepatitis B reactivation and *Pneumocystis jirovecii* pneumonia (PJP), respectively.^{3,4} Awareness of the specific infectious complications associated with anticancer drugs is critical to guide clinical vigilance, screening (eg, for tuberculosis and hepatitis B), and prophylactic (eg, antifungal and antiviral) strategies.^{5,6}

Although newer generation small molecule inhibitors have improved progression-free survival in patients with hematologic malignancies,^{7,8} the benefits of these treatments must be weighed against the risk of infectious adverse events (IAEs). Such complications can go unnoticed in initial randomized controlled trials (RCTs) because the trials are not adequately powered to detect rare events and may have shorter durations of follow-up than postmarketing studies. For example, ibrutinib, a Bruton tyrosine kinase inhibitor, was found to be associated with invasive fungal disease (IFD)⁹⁻¹¹ in postmarketing studies. Alternatively, the increased risk of IAEs may only manifest in certain populations or in the context of disease-specific interactions. For example, ibrutinib treatment confers a higher risk of PJP in patients with chronic lymphocytic leukemia (CLL) than in those with Waldenström macroglobulinemia.¹²

Venetoclax is a small molecule inhibitor that targets the anti-apoptotic protein BCL-2.¹³ Venetoclax is approved in the United States alone or in combination with other agents for CLL and acute myelogenous leukemia (AML).^{14,15} The impact of venetoclax on infectious complications may be complex. On one hand, BCL-2 inhibition may increase the risk of invasive pyogenic and fungal infections through neutropenia.^{16,17} On the other hand, data from patients with CLL suggest that venetoclax may mitigate some immunosuppressive effects of CLL,¹⁸ potentially reducing the risk of IAEs.

Previous 2020 reviews on venetoclax did not include an in-depth analysis of infectious complications and were biased by the inclusion of single-arm studies.^{19,20} To address this knowledge gap and inform patient management strategies from an infectious disease perspective, we conducted a systematic review and meta-analysis of RCTs to comprehensively evaluate the IAEs associated with venetoclax treatment for hematologic malignancies.

Methods

Protocol

This systematic review and meta-analysis adhered to a protocol registered a priori on the International Prospective Register of Systematic Reviews (PROSPERO) (CRD42021259416) and followed the guidelines from the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA),²¹ and the Cochrane Handbook for Systematic Reviews of Interventions.²²

Search strategy

A comprehensive search strategy was developed in collaboration with a research librarian. MedLine and Embase via Ovid were

searched from inception to 22 October 2022 for RCTs of adult populations on venetoclax regimens (supplemental Tables 1 and 2). There were no language restrictions; languages other than English and French were translated to English using Google Translate. Additionally, clinicaltrials.gov was searched for completed interventional studies involving venetoclax.

Study inclusion criteria

RCTs of any phase involving venetoclax for the treatment of hematologic malignancies were eligible for inclusion. This included those comparing venetoclax monotherapy or a venetoclax-containing regimen with another regimen regardless of the line of therapy, underlying hematologic malignancy, or use of antimicrobial prophylaxis. Studies were required to report on the incidence of IAEs and/or neutropenia. Studies with <30 patients in a treatment arm, studies limited to pediatric patients, conference abstracts, or gray literature were excluded.

Study selection

Search results were imported into Covidence and deduplicated.²³ Unique articles were then screened for relevance by title and abstract by 2 independent reviewers (C.P. and K.K.). Selected full-text articles were evaluated by 2 independent reviewers (C.P. and K.K.) to confirm they met inclusion criteria. Afterward, the reference lists of the included articles were manually reviewed for additional pertinent studies. When disagreements on the inclusion of an article occurred, they were resolved by consensus or with a third author when necessary.

Quality assessment

Consensus quality assessments were performed by 2 independent reviewers (C.P., A.N., O.D.L., or K.K.) using version 2 of the Cochrane risk-of-bias tool for randomized trials.²⁴ A summary figure of the quality assessments was generated using RevMan 5.41.²⁵

Data extraction

Data were extracted by 2 reviewers (C.P. and K.K.) using a standardized form. Discrepancies were resolved by consensus or arbitrated by a third author when necessary. The following data were extracted: study authors, year of publication, study phase, blinding, malignancy type, eligibility criteria, study arms, median follow-up time (according to the study's definition), antimicrobial prophylaxis, granulocyte colony stimulating factor (G-CSF) use, and timing of IAE data collection. Data on the incidence of neutropenia and IAEs were extracted according to the following groupings: grade 1 to 2, grade 3 to 5, fatal, total, and serious adverse events (SAEs). In addition, details were extracted on OIs and fatal IAEs.

Data analysis

A Bayesian analysis was used because it was thought to provide a more nuanced approach to the assessment of venetoclax's risk of harm than conventional frequentist analyses, which are dichotomized. When outcome data were reported by at least 2 studies, the data were pooled by Bayesian random effects meta-analysis. Bayesian meta-analyses were conducted using the bayesmeta package in R²⁶ using a weakly informative prior, with a population mean of 0 and standard deviation of 0.71 on the log risk ratio (RR)

scale,²⁷ which corresponds to an assumption that the effect size of most interventions in medicine falls between 0.25 and 4. The prior for heterogeneity (Tau) used those proposed by Turner et al²⁸ for the outcome of interest. The probability of any increased risk ($P(RR > 1)$) in the clinical outcomes of interest was computed using the posterior function.²⁶ The primary analyses were conducted separately for the 2 approved indications for venetoclax, AML, and CLL. Additional analyses were performed for other potential indications for venetoclax. Additionally, we pooled the incidence rates of fatal OIs by inverse variance meta-analysis^{29,30} and compared them between treatment groups by Bayesian meta-analysis. Person-time was estimated using the median treatment exposure time; if this was not reported, then the median follow-up time was used. When only a single data point was available for an outcome, the data were instead presented with descriptive statistics.

Results

Search results

Our search strategy returned a total of 2328 results, including 1777 unique articles. A total of 1759 articles were excluded during screening, and the remaining 18 articles were read in full. Among these 18 articles, 7 were included in the systematic review and meta-analysis, and the remaining were excluded (Figure 1). Hand searching of relevant references and a review of clinicaltrials.gov did not yield additional results.

Study and population characteristics

The 7 RCTs meeting inclusion criteria³¹⁻³⁷ were composed of 1190 and 877 patients randomized to a venetoclax-containing and

a comparator regimen, respectively. The hematologic malignancies studied included CLL ($n = 3$),^{31,32,34} AML ($n = 2$),^{33,35} multiple myeloma (MM; $n = 1$),³⁷ and follicular lymphoma (FL; $n = 1$).³⁶ Four studies only enrolled patients with previously untreated disease,³²⁻³⁵ whereas the remaining studies only enrolled patients with relapsed or refractory disease.^{31,36,37} Median follow-up time ranged from 12.0 to 28.1 months. Treatment arms were variable with no 2 studies comparing identical regimens. The majority of patients (62.5%) included were male, and the median age ranged from 61 to 76 years. Only 1 study used different strategies for antimicrobial prophylaxis between the 2 study arms. Although prophylaxis was not included at the initiation of the study, Kumar et al later amended their protocol to administer PJP and antibacterial prophylaxis to all patients assigned to venetoclax.³⁷ G-CSF use was reported in 3 studies^{32,33,36} and ranged from 32.0% to 43.5% and 34.0% to 45.8% among patients assigned to venetoclax and comparators, respectively. None of the studies elaborated on the duration of G-CSF use. A detailed description of the included studies is outlined in Table 1.

Data on the extracted IAEs are presented in supplemental Table 3, and details on the fatal IAEs are presented in supplemental Table 4. Importantly, there was heterogeneity in the reporting of rare and nonfatal IAEs, especially OIs, among the included studies. Only Kumar et al³⁷ reported all adverse events occurring during the study. The other studies applied a minimum threshold prevalence to report the adverse events ranging from >2% to 20% for grade 3/4 adverse events, >1% to 5% for SAEs, and >10% to 20% for the remainder of adverse events.

Quality assessments

Three studies were judged at a low overall risk of bias,^{33,35,37} and 4 had some concerns^{31,32,34,36} (Figure 2). The predominant concern was related to the open-label study design and resultant potential for ascertainment bias in the reporting and/or grading of IAEs or in the differential use of antimicrobial prophylaxis. All 3 CLL trials were at some risk of bias, and both AML trials had a low risk of bias.

CLL

The 3 CLL RCTs were open-label and head-to-head comparisons, comprising a total of 1032 patients.^{31,32,34} Two studies involved patients receiving firstline therapy, and the other involved subsequent-line therapy.

The pooled RR of grade 3 to 5 and fatal IAEs were 1.11 (95% credible interval [CrI], 0.74-1.68; $P[RR > 1] = 71.2\%$) and 1.16 (95% CrI, 0.53-2.57; $P[RR > 1] = 64.5\%$), respectively (Table 2). The risk of high-grade neutropenia (RR = 1.07; 95% CrI, 0.64-1.74; $P[RR > 1] = 63.4\%$) and febrile neutropenia (RR = 0.76; 95% CrI, 0.40-1.49; $P[RR > 1] = 20.5\%$) were also similar between the 2 groups. Similarly, we did not identify a highly probable increased risk of sepsis, pneumonia, upper respiratory tract infections, or cellulitis.

Three fatal OIs were reported in the CLL trial: 2 in the comparator group involving a case of invasive *Scedosporium* infection and sepsis from *Listeria monocytogenes*, and the third fatal OI was a case of fungal pneumonia in a patient treated with venetoclax. The incidences of fatal OIs per 100 person-years were 0.34 (95% confidence interval [CI], 0.09-1.37) and 0.56 (95% CI, 0.18-1.73) for venetoclax




Figure 1. PRISMA diagram.

Table 1. Study characteristics

Author, y	Phase	Design	Malignancy type	Eligibility criteria	Venetoclax regimen, N	Comparator regimen, N	Timing of reporting of AEs	Antimicrobial prophylaxis	G-CSF use	Median follow-up time (mo)
DiNardo, 2020	3	Double blind	AML	Untreated disease and ineligible for standard induction therapy	Venetoclax + azacitidine, N = 286	Placebo + azacitidine, N = 145	30 d after discontinuation of the study drug	Antimicrobial prophylaxis was received by 83% and 81% of patients assigned to venetoclax and comparator, respectively	Venetoclax arm: 32% Comparator arm: NR	20.5
Fischer, 2019	3	Open-label	CLL	Untreated and comorbidities	Venetoclax + Obinutuzumab, N = 216	Chlorambucil + Obinutuzumab, N = 216	28 d after the final dose of venetoclax or 90 d after the final dose of obinutuzumab, whichever is longer. Grade 3 and 4 IAEs were collected until 2 y after the final study drug dose	When clinically indicated, antimicrobial prophylaxis was permitted	Venetoclax arm: 43.5% Comparator arm: 45.8%	28.1
Kater, 2022	3	Open-label	CLL	Untreated and either older (>65 y) or comorbidities	Venetoclax + ibrutinib, N = 106	Chlorambucil + obinutuzumab, N = 105	Unclear	NR	Venetoclax arm: NR Comparator arm: NR	27.7
Kumar, 2020	3	Doubleblind	MM	Relapsed or refractory disease	Venetoclax + bortezomib + dexamethasone N = 194	Placebo + bortezomib + dexamethasone, N = 97	30 d after the final dose of the study drug	All patients received varicella zoster virus prophylaxis. Initially, antibacterial and PJP prophylaxis were given at the discretion of the study investigator; however, later they were required for all patients receiving venetoclax	Venetoclax arm: NR Comparator arm: NR	18.7
Seymour, 2018	3	Open-label	CLL	Relapsed or refractory disease	Venetoclax + rituximab, N = 194	Bendamustine + rituximab, N = 195	28 d after the final dose of the study drug, up to a maximum of 2 y for venetoclax, or 90 d after the final dose of rituximab	Antimicrobial prophylaxis was permitted when clinically indicated	Venetoclax arm: NR Comparator arm: NR	23.8
Wei, 2020	3	Double blind	AML	Previously untreated disease and ineligible for intensive chemotherapy	Venetoclax + cytarabine, N = 143	Placebo + cytarabine, N = 68	30 d after the final dose of the study drug	All subjects with absolute neutrophil counts below 500 were administered antimicrobial prophylaxis	Venetoclax arm: NR Comparator arm: NR	12.0
Zinzani, 2020	2	Open-label	FL	Relapsed or refractory disease	Venetoclax + bendamustine + rituximab, N = 51	Bendamustine + rituximab, N = 51	30 d after the final dose of the study drug or 90 d after the final dose of rituximab	NR	Venetoclax arm: 42.9% Comparator arm: 34.0%	18.2 and 18.4 in the venetoclax and control arms, respectively

NR, not reported.

	Randomization process	Deviations from the intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall risk of bias
DiNardo 2020	+	+	+	+	+	+
Fischer 2019	+	+	+	?	+	?
Kater 2022	+	+	+	?	?	?
Kumar 2020	+	+	+	+	+	+
Seymour 2018	+	+	+	?	+	?
Wei 2020	+	+	+	+	+	+
Zinzani 2020	+	+	+	?	+	?

Figure 2. Cochrane RoB 2 summary. RoB 2, Cochrane risk-of-bias tool for randomized trials version 2.

and comparator regimens (Figure 3), respectively (RR = 0.83; 95% CrI, 0.27-2.58; $P[RR > 1] = 37.6\%$). Only 1 other IFD was reported; Kater et al observed 1 case (0.9%) of nonfatal bronchopulmonary aspergillosis in the venetoclax arm.³⁴ The Kater et al study also reported 1 case each of nonfatal cytomegalovirus pneumonia and PJP among venetoclax recipients but none among controls.³⁴

AML

The 2 included AML RCTs were double-blind and placebo-controlled, using venetoclax as the firstline agent, and comprised 642 patients in total.^{33,35}

There were insufficient data points to compute the risk of total, fatal, and high-grade IAEs as well as fatal OIs in patients with AML. However, in the venetoclax arm of the DiNardo et al³³ trial, there was a higher number of total (239 [83.6%] vs 97 [66.9%]) and high-grade (180 [62.9%] vs 74 [51.0%]) IAEs. Venetoclax was associated with a probable increased risk of high-grade neutropenia (RR = 1.71; 95% CrI, 0.87-3.17; $P[RR > 1] = 94.6\%$) and febrile neutropenia (RR = 1.49; 95% CrI, 0.78-2.60; $P[RR > 1] = 90.6\%$; Table 3). However, the rates of sepsis and pneumonia were comparable. No information on fatal IAEs or OIs was reported in the AML studies.

MM

Only 1 study on MM was included,³⁷ which precluded meta-analysis. Progression-free survival was improved in the venetoclax

arm of the Kumar et al study.³⁷ However, overall survival was poorer in this arm, predominantly driven by an increase in fatal IAEs (8 [4.1%] vs 0 [0.0%]) despite a protocol amendment implementing antibacterial, α herpesvirus, and PJP prophylaxis to the venetoclax arm. Of the 8 fatal IAEs in the venetoclax arm, 5 (2.6%) were unspecified sepsis and 3 (1.5%) were pneumonia. A total of 9 cases (4.6%) of varicella were reported among venetoclax recipients vs 1 (1.0%) among controls. No other OIs were reported. Both high-grade neutropenia (35 [18.0%] vs 7 [7.2%]) and febrile neutropenia (5 [2.6%] vs 0 [0.0%]) were numerically higher in the venetoclax arm.

FL

Only 1 trial on FL was identified, preventing meta-analysis.³⁶ There was 1 episode (2.0%) of fatal pneumonia in the venetoclax arm and no fatal IAEs in the comparator arm. High-grade neutropenia (29 [56.9%] vs 14 [27.5%]) and febrile neutropenia (6 [11.8%] vs 3 [5.9%]) were numerically higher in the venetoclax arm. For OIs, Zinzani et al documented 3 cases (5.9%) of PJP in the venetoclax arm but none in the comparator arm.³⁶

Overall analyses

Pooled analyses irrespective of indication for venetoclax are presented in supplemental Table 5. The pooled RR of total and high-grade IAEs were 1.14 (95% CrI, 0.76-1.66; $P[RR > 1] = 81.1\%$) and 1.16 (95% CrI, 0.86-1.55; $P[RR > 1] = 86.6\%$), respectively. There was a probable increased risk of high-grade neutropenia (RR = 1.44; 95% CrI, 1.01-2.10; $P[RR > 1] = 97.8\%$).

Discussion

This extensive review of the venetoclax RCT data suggests that venetoclax has a low probability of increasing the risk of infection or neutropenia in CLL. There was insufficient evidence to determine whether venetoclax increases the risk of IAEs in AML; however, there was a probable increase in high-grade neutropenia (94.6%) and febrile neutropenia (90.6%).

As demonstrated in early trials of venetoclax,^{16,17} we found a high prevalence of high-grade and febrile neutropenia. In CLL, there was only a 63.4% probability of increased risk of high-grade neutropenia and only a 71.2% probability of association with an increased risk of IAEs. Our findings may be explained by the immunomodulatory effects of venetoclax in CLL. Although venetoclax treatment results in an absolute reduction in T, B, and natural killer cells, it also restores natural killer cell function.¹⁸ Therefore, the immunosuppressive effects of venetoclax-induced neutropenia may be attenuated by a partial recovery of cellular immunity in CLL. In AML, however, a probable increased risk of neutropenia was noted, which may have driven the numerically increased risk of infection observed in the DiNardo et al study.³³

Initial trials of ibrutinib failed to recognize an increased risk of rare but severe IFDs in patients with hematologic malignancies,^{8,38,39} and the first report suggesting the association was published 3 years after its licensure.⁴⁰ Concern over ibrutinib conferring an increased risk of IFDs^{41,42} underscores the importance of scrutinizing the IAEs of novel small molecule inhibitors. Recent observational studies have reported a 4% to 13% incidence of probable or proven IFDs on venetoclax; however, this risk was not modified by antifungal prophylaxis.⁴³⁻⁴⁵ A large study pooling the safety data

Table 2. Pooled CLL overall outcomes

Outcome	Venetoclax, n/N (%)	Comparator, n/N (%)	Number of studies included	RR (95% CrI)	P(RR > 1), %
Total IAE	—	—	—	—	—
Grade 3-5 IAEs	101/516 (19.6%)	92/516 (17.8%)	3	1.11 (0.74-1.68)	71.2
Fatal IAEs	10/516 (1.9%)	8/516 (1.6%)	3	1.16 (0.53-2.57)	64.5
Fatal OI*	1/516 (0.2%)	2/516 (0.4%)	3	0.83 (0.27-2.58)	37.6
Grade 1-2 neutropenia	23/516 (4.5%)	38/516 (7.4%)	3	0.66 (0.37-1.21)	8.6
Grade 3-5 neutropenia	261/516 (50.6%)	228/516 (44.2%)	3	1.07 (0.64-1.74)	63.4
Total neutropenia	284/516 (55.0%)	266/516 (51.6%)	3	1.02 (0.63-1.63)	54.1
SAE neutropenia	—	—	—	—	—
Grade 1-2 febrile neutropenia	—	—	—	—	—
Grade 3-5 febrile neutropenia	20/516 (3.9%)	29/516 (5.6%)	3	0.76 (0.40-1.49)	20.5
Total febrile neutropenia	—	—	—	—	—
SAE febrile neutropenia	19/516 (3.7%)	27/516 (5.2%)	3	0.76 (0.39-1.50)	20.9
SAE sepsis	7/516 (1.4%)	7/516 (1.4%)	3	1.00 (0.39-2.58)	50.0
Grade 1-2 pneumonia	9/300 (3%)	11/300 (3.7%)	2	0.88 (0.39-1.96)	37.2
Grade 3-5 pneumonia	30/516 (5.8%)	30/516 (5.8%)	3	1.01 (0.58-1.74)	50.9
Total pneumonia	29/300 (9.7%)	29/300 (9.7%)	2	0.93 (0.52-1.67)	39.7
SAE pneumonia	32/516 (6.2%)	30/516 (5.8%)	3	1.06 (0.62-1.81)	58.2
Total upper respiratory tract infections	56/300 (18.7%)	43/300 (14.3%)	2	1.23 (0.69-2.08)	78.5
Total nasopharyngitis	—	—	—	—	—
Total bronchitis	—	—	—	—	—
Total urinary tract infections	—	—	—	—	—
SAE cellulitis	4/322 (1.2%)	0/321 (0.0%)	2	1.54 (0.47-5.03)	76.3

Missing data (—) indicate <2 studies reporting the outcome.

All 3 CLL studies had some risk of bias on the Cochrane risk-of-bias tool for randomized trials version 2 (RoB 2) grading.

*The RR was computed using the incidence.


**Figure 3. Incidence of fatal OIs in CLL.** CI, confidence interval.

Table 3. Pooled AML overall outcomes

Outcome	Venetoclax, n/N (%)	Comparator, n/N (%)	Number of studies included	RR (95% CrI)	P(RR>1), %
Total IAE	—	—	—	—	—
Grade 3-5 IAEs	—	—	—	—	—
Fatal IAEs	—	—	—	—	—
Fatal OI	—	—	—	—	—
Grade 1-2 neutropenia	—	—	—	—	—
Grade 3-5 neutropenia	185/429 (43.1%)	52/213 (24.4%)	2	1.71 (0.87-3.17)	94.6
Total neutropenia	—	—	—	—	—
SAE neutropenia	17/429 (4.0%)	3/213 (1.4%)	2	1.62 (0.60-4.26)	83.3
Grade 1-2 febrile neutropenia	—	—	—	—	—
Grade 3-5 febrile neutropenia	163/429 (38.0%)	47/213 (22.1%)	2	1.49 (0.78-2.60)	90.6
Total febrile neutropenia	—	—	—	—	—
SAE febrile neutropenia	107/429 (24.9%)	27/213 (12.7%)	2	1.53 (0.71-2.97)	87.8
SAE sepsis	24/429 (5.6%)	16/213 (7.5%)	2	0.80 (0.41-1.59)	25.5
Grade 1-2 pneumonia	—	—	—	—	—
Grade 3-5 pneumonia	—	—	—	—	—
Total pneumonia	94/429 (21.9%)	50/213 (23.5%)	2	0.96 (0.59-1.63)	43.3
SAE pneumonia	65/429 (15.2%)	39/213 (18.3%)	2	0.87 (0.51-1.56)	29.8
Total upper respiratory tract infections	—	—	—	—	—
Total nasopharyngitis	—	—	—	—	—
Total bronchitis	—	—	—	—	—
Total urinary tract infections	—	—	—	—	—
SAE cellulitis	—	—	—	—	—

Missing data (—) indicate <2 studies reporting the outcome.

Both AML studies had low risk of bias on the Cochrane risk-of-bias tool for randomized trials version 2 (RoB 2) grading of indication for venetoclax.

from 3 venetoclax CLL phase 1/2 trials found a lower prevalence of OIs of 3.1%, with a median time to OI of 4.5 months.⁴⁶ Our results confirm a low incidence of fatal OIs in CLL. Furthermore, we did not identify specific infectious complications that would benefit from specific antimicrobial prophylaxis or screening measures beyond the current standard of care.

Although small molecule inhibitors significantly improve various oncologic outcomes (eg, progression- and event-free survival), it is important to weigh these benefits against the risk of infectious complications and overall survival. Although venetoclax improved progression-free survival in the MM study by Kumar et al,³⁷ it increased the risk of all-cause mortality, powered predominantly by an increase in fatal IAEs. Therefore, in MM, the risk-benefit profile of venetoclax appears unfavorable. In contrast, in CLL, venetoclax appears more beneficial given that our results found a low probability of increased risk of IAEs or neutropenia, and the 3 included trials demonstrated improved progression-free survival with venetoclax as well as an overall mortality benefit in the Seymour et al study.^{31,32,34} In AML, our results suggested a probable increased risk of neutropenia, and 1 study suggested an increased risk of infection, but both AML trials demonstrated improved oncologic outcomes with 1 finding an overall mortality benefit with venetoclax.^{33,35}

To our knowledge, this systematic review is the first investigating the impact of venetoclax on IAEs using RCT data. We used a

comprehensive search strategy, and no additional articles were identified through hand searching. The inclusion of only RCTs limited the risk of selection, detection, and attrition bias that confound observational studies. Moreover, none of the included RCTs were considered at a high risk of bias. Despite these strengths, our results must be interpreted within the included studies' contexts. First, our study may be underpowered to detect differences in rare IAEs, and the included studies were inconsistent about reporting infrequent IAEs. Second, because there are few RCTs published on venetoclax, RCTs were included regardless of underlying malignancy, line of therapy, and concomitant chemotherapy regimen, which introduces some interstudy heterogeneity. Third, studies were pooled regardless of the dose or duration of venetoclax, which may obscure dose- or duration-dependent toxicities. Fourth, we were unable to perform time-dependent analyses because these data were unavailable, which could introduce bias from a competing risk of malignancy-specific mortality. Fifth, an inherent limitation of including open-label studies is the possibility of ascertainment bias in the reporting/grading of IAEs and the differential use of antimicrobial prophylaxis or G-CSF. Sixth, there was significant heterogeneity in the reporting and definitions of IAEs, and many studies did not provide sufficient detail for more granular analyses. Lastly, multiple hypothesis testing increases the risk of type 1 error, and these results must be interpreted with this consideration. Nevertheless, our study is the most current and comprehensive assessment of IAEs of venetoclax to date, and our

probabilistic analysis provides some insight beyond the traditional *P* value–driven dichotomous analysis.

Conclusions

Venetoclax had a low probability of increased risk of infection or neutropenia in CLL. In AML, however, a probable increased risk of high-grade and febrile neutropenia was noted, with 1 study suggesting an increased risk of IAEs. The risk of OIs was overall low. Our analyses did not identify any specific IAEs that would necessitate additional antimicrobial prophylaxis or pre-emptive testing. Nevertheless, we recommend that postmarketing studies continue surveillance of the IAEs associated with venetoclax treatment.

Acknowledgment

The authors thank Ibtisam Mahmoud, research librarian, for her assistance in designing the search strategy.

This work had no sources of funding.

Authorship

Contribution: M.P.C., K.D., V.C., and C.P. conceptualized the study and investigated the data; M.P.C., K.D., V.C., C.P., and T.C.L. designed the methodology; C.P., K.D., and V.C. validated the data; C.P. and T.C.L. provided formal analysis; C.P. and M.P.C. provided resources; C.P., K.D., V.C., A.N., O.D.L., M. Sorin, and K.K. curated the data; C.P., M.P.C., E.G.M., T.C.L., M. Sebag, and K.D. wrote the original draft; C.P. provided visualization; M.P.C. supervised the project; C.P., M.P.C., and K.D. administered the project; and all authors reviewed and edited the manuscript.

Conflict-of-interest disclosure: M.P.C. reports operating grants from the Canadian Institutes of Health Research during the conduct of the study and salary support from the Fonds de

Recherche du Québec – Santé, both unrelated to this work; personal fees from GEn1E Lifesciences and Nomic Bio as a member of the scientific advisory board; personal fees from AstraZeneca as a scientific consultant; research support from Cidara Therapeutics, Scynexis, Inc, and Amplyx Pharmaceuticals during the conduct of the study but outside the submitted work; is the cofounder of Kanvas Biosciences Inc and owns equity in the company; and has pending patents, including: (1) methods for detecting tissue damage, graft-versus-host disease, and infections using cell-free DNA profiling, (2) methods for assessing the severity and progression of severe acute respiratory syndrome coronavirus 2 infections using cell-free DNA pending, and (3) rapid identification of antimicrobial resistance and other microbial phenotypes using highly multiplexed fluorescence in situ hybridization. M. Sebag is on the advisory boards of Janssen, Pfizer, Amgen, Novartis, Gilead, Bristol Myers Squibb, Focus Therapeutics, and Sanofi, and reports funding from the Canadian Institutes of Health Research and Janssen Pharmaceuticals. E.G.M. and T.C.L. have operating grants from the Canadian Institutes of Health Research and receive salary support from the Fonds de Recherche du Québec – Santé outside of this work. The remaining authors declare no competing financial interests.

ORCID profiles: C.P., [0000-0001-6790-220X](https://orcid.org/0000-0001-6790-220X); A.N., [0000-0001-9773-3515](https://orcid.org/0000-0001-9773-3515); O.D.L., [0000-0002-4222-6265](https://orcid.org/0000-0002-4222-6265); E.G.M., [0000-0003-0783-0624](https://orcid.org/0000-0003-0783-0624); T.C.L., [0000-0002-2267-4239](https://orcid.org/0000-0002-2267-4239).

Correspondence: Connor Prosty, Infectious Diseases, McGill University, 1001 Decarie Blvd E5-1820, Montréal, QC H4A 3J1, Canada; email: connor.prosty@mail.mcgill.ca; and Matthew P. Cheng, Infectious Diseases, McGill University, 1001 Decarie Blvd EM3.3218, Montréal, QC H4A 3J1, Canada; email: matthew.cheng@mcgill.ca.

References

1. Chen L, Zheng Y, Yu K, et al. Changing causes of death in persons with haematological cancers 1975-2016. *Leukemia*. 2022;36(7):1850-1860.
2. Zembower TR. Epidemiology of infections in cancer patients. *Cancer Treat Res*. 2014;161:43-89.
3. Yale SH, Limper AH. Pneumocystis carinii pneumonia in patients without acquired immunodeficiency syndrome: associated illness and prior corticosteroid therapy. *Mayo Clin Proc*. 1996;71(1):5-13.
4. Evens AM, Jovanovic BD, Su YC, et al. Rituximab-associated hepatitis B virus (HBV) reactivation in lymphoproliferative diseases: meta-analysis and examination of FDA safety reports. *Ann Oncol*. 2011;22(5):1170-1180.
5. Tsutsumi Y, Yamamoto Y, Ito S, et al. Hepatitis B virus reactivation with a rituximab-containing regimen. *World J Hepatol*. 2015;7(21):2344-2351.
6. Baden LR, Swaminathan S, Angarone M, et al. Prevention and treatment of cancer-related infections, version 2.2016, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Cancer Netw*. 2016;14(7):882-913.
7. Woyach JA, Ruppert AS, Heerema NA, et al. Ibrutinib regimens versus chemoimmunotherapy in older patients with untreated CLL. *N Engl J Med*. 2018;379(26):2517-2528.
8. Wang ML, Jurczak W, Jerkeman M, et al. Ibrutinib plus bendamustine and rituximab in untreated mantle-cell lymphoma. *N Engl J Med*. 2022;386(26):2482-2494.
9. Maschmeyer G, De Greef J, Mellingshoff SC, et al. Infections associated with immunotherapeutic and molecular targeted agents in hematology and oncology. A position paper by the European Conference on Infections in Leukemia (ECIL). *Leukemia*. 2019;33(4):844-862.
10. Ball S, Das A, Vutthikraivit W, et al. Risk of infection associated with ibrutinib in patients with B-cell malignancies: a systematic review and meta-analysis of randomized controlled trials. *Clin Lymphoma Myeloma Leuk*. 2020;20(2):87-97.e5.
11. Maffei R, Maccaferri M, Arletti L, et al. Immunomodulatory effect of ibrutinib: reducing the barrier against fungal infections. *Blood Rev*. 2020;40:100635.
12. Ryan CE, Cheng MP, Issa NC, Brown JR, Davids MS. Pneumocystis jirovecii pneumonia and institutional prophylaxis practices in CLL patients treated with BTK inhibitors. *Blood Adv*. 2020;4(7):1458-1463.

13. Souers AJ, Levenson JD, Boghaert ER, et al. ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med*. 2013;19(2):202-208.
14. *FDA Grants Regular Approval to Venetoclax in Combination for Untreated Acute Myeloid Leukemia*. US Food and Drug Administration; 2020.
15. *FDA Approves Venetoclax for CLL and SLL*. US Food and Drug Administration; 2019.
16. Roberts AW, Davids MS, Pagel JM, et al. Targeting BCL2 with venetoclax in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2016;374(4):311-322.
17. Stilgenbauer S, Eichhorst B, Schetelig J, et al. Venetoclax in relapsed or refractory chronic lymphocytic leukaemia with 17p deletion: a multicentre, open-label, phase 2 study. *Lancet Oncol*. 2016;17(6):768-778.
18. de Weerdt I, Hofland T, de Boer R, et al. Distinct immune composition in lymph node and peripheral blood of CLL patients is reshaped during venetoclax treatment. *Blood Adv*. 2019;3(17):2642-2652.
19. Li Q, Cheng L, Shen K, et al. Efficacy and safety of Bcl-2 inhibitor venetoclax in hematological malignancy: a systematic review and meta-analysis of clinical trials. *Front Pharmacol*. 2019;10:697.
20. Bewersdorf JP, Giri S, Wang R, et al. Venetoclax as monotherapy and in combination with hypomethylating agents or low dose cytarabine in relapsed and treatment refractory acute myeloid leukemia: a systematic review and meta-analysis. *Haematologica*. 2020;105(11):2659-2663.
21. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
22. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.4 (updated August 2023)*. Cochrane; 2023. Accessed 2 March 2023. <https://training.cochrane.org/handbook>
23. Covidence. Systematic review software, Veritas Health Innovation, Melbourne, Australia. Accessed 2 March 2023. <https://www.covidence.org/>
24. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:l4898.
25. *Review Manager (RevMan). Version 5.4.1*. The Cochrane Collaboration; 2020.
26. bayesmeta: Bayesian Random-Effects Meta-Analysis and Meta-Regression. Version 3.2. *CRAN*. 2022. Accessed 2 March 2023. <https://CRAN.R-project.org/package=bayesmeta>
27. Zampieri FG, Casey JD, Shankar-Hari M, Harrell FE, Harhay MO. Using Bayesian methods to augment the interpretation of critical care trials. An overview of theory and example reanalysis of the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial. *Am J Respir Crit Care Med*. 2021;203(5):543-552.
28. Turner RM, Jackson D, Wei Y, Thompson SG, Higgins JPT. Predictive distributions for between-study heterogeneity and simple methods for their application in Bayesian meta-analysis. *Stat Med*. 2015;34(6):984-998.
29. meta: General Package for Meta-Analysis. Version 5.2-0. *CRAN*. 2022. Accessed 21 March 2022. <https://CRAN.R-project.org/package=meta>
30. Viechtbauer W. Conducting meta-analyses in R with the metafor Package. *J Stat Softw*. 2010;36(3):1-48.
31. Seymour JF, Kipps TJ, Eichhorst B, et al. Venetoclax-rituximab in relapsed or refractory chronic lymphocytic leukemia. *N Engl J Med*. 2018;378(12):1107-1120.
32. Fischer K, Al-Sawaf O, Bahlo J, et al. Venetoclax and obinutuzumab in patients with CLL and coexisting conditions. *N Engl J Med*. 2019;380(23):2225-2236.
33. DiNardo CD, Jonas BA, Pullarkat V, et al. Azacitidine and venetoclax in previously untreated acute myeloid leukemia. *N Engl J Med*. 2020;383(7):617-629.
34. Kater AP, Owen C, Moreno C, et al. Fixed-duration ibrutinib-venetoclax in patients with chronic lymphocytic leukemia and comorbidities. *NEJM Evidence*. 2022;1(7):EVIDoa2200006.
35. Wei AH, Montesinos P, Ivanov V, et al. Venetoclax plus LDAC for newly diagnosed AML ineligible for intensive chemotherapy: a phase 3 randomized placebo-controlled trial. *Blood*. 2020;135(24):2137-2145.
36. Zinzani PL, Flinn IW, Yuen SLS, et al. Venetoclax-rituximab with or without bendamustine vs bendamustine-rituximab in relapsed/refractory follicular lymphoma. *Blood*. 2020;136(23):2628-2637.
37. Kumar SK, Harrison SJ, Cavo M, et al. Venetoclax or placebo in combination with bortezomib and dexamethasone in patients with relapsed or refractory multiple myeloma (BELLINI): a randomised, double-blind, multicentre, phase 3 trial. *Lancet Oncol*. 2020;21(12):1630-1642.
38. Byrd JC, Brown JR, O'Brien S, et al. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med*. 2014;371(3):213-223.
39. Burger JA, Tedeschi A, Barr PM, et al. Ibrutinib as initial therapy for patients with chronic lymphocytic leukemia. *N Engl J Med*. 2015;373(25):2425-2437.
40. Ibrutinib for chronic lymphocytic leukemia. *N Engl J Med*. 2016;374(16):1592-1595.
41. Ghez D, Calleja A, Protin C, et al. Early-onset invasive aspergillosis and other fungal infections in patients treated with ibrutinib. *Blood*. 2018;131(17):1955-1959.
42. Ruchlemer R, Ben-Ami R, Bar-Meir M, et al. Ibrutinib-associated invasive fungal diseases in patients with chronic lymphocytic leukaemia and non-Hodgkin lymphoma: an observational study. *Mycoses*. 2019;62(12):1140-1147.

43. Chen EC, Liu Y, Harris CE, et al. Outcomes of antifungal prophylaxis for newly diagnosed AML patients treated with a hypomethylating agent and venetoclax. *Leuk Lymphoma*. 2022;63(8):1934-1941.
44. On S, Rath CG, Lan M, et al. Characterisation of infections in patients with acute myeloid leukaemia receiving venetoclax and a hypomethylating agent. *Br J Haematol*. 2022;197(1):63-70.
45. Aldoss I, Dadwal S, Zhang J, et al. Invasive fungal infections in acute myeloid leukemia treated with venetoclax and hypomethylating agents. *Blood Adv*. 2019;3(23):4043-4049.
46. Davids MS, Hallek M, Wierda W, et al. Comprehensive safety analysis of venetoclax monotherapy for patients with relapsed/refractory chronic lymphocytic leukemia. *Clin Cancer Res*. 2018;24(18):4371-4379.