

Mutations highly specific for secondary AML are associated with poor outcomes in ELN favorable risk *NPM1*-mutated AML

Onyee Chan,¹ Najla Al Ali,¹ Hammad Tashkandi,² Austin Ellis,² Somedeb Ball,¹ Justin Grenet,³ Caroline Hana,⁴ Yehuda Deutsch,⁴ Ling Zhang,² Mohammad Hussaini,² Jinming Song,² Seongseok Yun,¹ Chetasi Talati,¹ Andrew Kuykendall,¹ Eric Padron,¹ Alison Walker,¹ Gail Roboz,³ Pinkal Desai,³ David Sallman,¹ Kendra Sweet,¹ Rami Komrokji,¹ and Jeffrey Lancet¹

¹Department of Malignant Hematology and ²Department of Hematopathology, Moffitt Cancer Center, Tampa, FL; ³Division of Hematology and Oncology, Weill Cornell Medical College, New York, NY; and ⁴Department of Malignant Hematology and Cellular Therapy at Memorial Healthcare System, Moffitt Cancer Center, Pembroke Pines, FL

Key Points

- Patients with favorable-risk AML who had *NPM1* mutation have poor prognosis if they also harbor mutations highly specific for sMut.
- Patients with sMut had better outcomes if they achieved *NPM1* MRD negativity.

Acute myeloid leukemia (AML) is a heterogeneous malignancy with outcomes largely predicted by genetic abnormalities. Mutations of *NPM1* are common in AML, occurring in ~30% of cases, and generally considered a favorable risk factor. Mutations highly specific for secondary AML (sMut) have been shown to confer poor prognosis, but the overall impact of these mutations in the setting of favorable-risk AML defined by mutant *NPM1* remains unclear. In this multicenter study of patients with AML (n = 233) with *NPM1* mutation at diagnosis, we observed that patients with sMut had worse overall survival (OS) than those without sMut (15.3 vs 43.7 months; $P = .002$). Importantly, this finding persisted in the European LeukemiaNet (ELN) 2017–defined favorable risk subset (14.7 months vs not reached; $P < .0001$). Among patients who achieved *NPM1* measurable residual disease (MRD) negativity, longer OS was observed in the entire cohort ($P = .015$) as well as in both the sMut subset (MRD negative: median OS (mOS) 73.9 months vs MRD positive: 12.3 months; $P = .0170$) and sMut ELN 2017–favorable subset (MRD negative: mOS 27.3 vs MRD positive: 10.5 months; $P = .009$). Co-occurrence of sMut and mutant *NPM1* confers a poor prognosis in AML.

Introduction

Mutated nucleophosmin (*NPM1*) is one of the most common molecular abnormalities in acute myeloid leukemia (AML), occurring in ~30% of cases regardless of cytogenetics.¹ Wild-type *NPM1* contains 12 exons and encodes for a ubiquitous multifunctional protein that shuttles between the nucleus and cytoplasm. Some of the major functions of *NPM1* include ribosome biogenesis, p53-dependent stress response, and genomic stability maintenance.² Aberrant cytoplasmic dislocation is the hallmark feature in *NPM1*-mutated AML, a discovery leading to the recognition of *NPM1*-mutated AML as a distinct entity within the 2016 World Health Organization classification.³ Mutated *NPM1* is enriched in AML with normal cytogenetics and is associated with higher response rates to induction chemotherapy and better overall survival (OS) than many other AML subtypes.^{4,5} Besides *CEBPA*-bZip mutations, mutant *NPM1* is the only other mutation that can define favorable-risk disease under the European LeukemiaNet (ELN) 2017 and 2022 criteria.^{6,7} It is well documented that certain gene-gene interactions may affect the prognostic impact of a single gene mutation. For example, in ELN 2022, co-occurrence of *NPM1* and *FLT3-ITD* mutations defines intermediate-risk disease rather than favorable-risk disease

Submitted 11 July 2023; accepted 6 December 2023; prepublished online on *Blood Advances* First Edition 3 January 2024; final version published online 28 February 2024. <https://doi.org/10.1182/bloodadvances.2023011173>.

Publication-related data are available upon reasonable request from the corresponding author, Onyee Chan (onyee.chan@moffitt.org).

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

(seen in cases with *NPM1* mutation and wild-type *FLT3*). A recent study observed that outcomes of *NPM1* and *WT1*-co-mutated AML in patients aged <60 years resembled outcomes in patients at adverse risk.⁸ Another study showed the combination of *NPM1*, *DNMT3A*, and *IDH* mutations led to an inferior OS.⁹ In ELN 2022, the presence of mutations highly specific for secondary AML (sAML), including *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and/or *ZRSR2* (sMut)-defined adverse risk.^{6,10} Older patients with de novo AML (dAML) with these genetic alterations share outcomes similar to those with conventional-defined sAML, with lower complete remission (CR) rate and decreased survival.¹⁰ To date, ELN 2022 has not yet classified AML with sMut that co-occurs with favorable-risk molecular features (such as mutant *NPM1*) as adverse.⁶ As such, we set out to study the prognostic impact of sMut co-occurring with mutant *NPM1* in AML, including its prognostic effect among patients who achieve mutant *NPM1*-measurable residual disease (MRD) negativity.

Patients and methods

Patient selection

We performed a retrospective review of 233 patients with AML who had *NPM1* mutation and were treated at the Moffitt Cancer Center (n = 160), Weill Cornell (n = 51), and Memorial Healthcare System (n = 22) between 2013 and 2022. Inclusion was restricted to patients who had next-generation sequencing (NGS) performed at diagnosis. Patients who participated in clinical trials at any point during their treatment course were included. Demographics, disease-specific variables, and clinical outcomes were collected in accordance with the institutional review board-approved protocol. The study was conducted in accordance with the Declaration of Helsinki. dAML is defined as AML developed in patients without clinical history of prior myeloid hematologic neoplasm or exposure to potentially leukemogenic agents. sAML refers to AML arising from an antecedent myeloid hematologic neoplasm (based on clinical history) such as myelodysplastic syndromes or myeloproliferative neoplasm. Therapy-related AML is defined as AML arising from prior cytotoxic, radiation, or immunosuppressive therapy for an unrelated disease. Risk stratification and response to therapy were defined using the ELN 2017 criteria, unless otherwise specified.⁷

Molecular analysis

The majority of somatic mutations were assessed using TruSight Myeloid Sequencing Panel (Illumina, San Diego, CA) that targets 54 genes by NGS with a detection threshold of 5% to determine presence or absence of mutations based on institutional standards. These genes include *ABL1*, *ASXL1*, *ATRX*, *BCOR*, *BCORL1*, *BRAF*, *CALR*, *CBL*, *CBLB*, *CBLC*, *CDKN2A*, *CEBPA*, *CSF3R*, *CUX1*, *DNMT3A*, *ETV6*, *EZH2*, *FBXW7*, *FLT3*, *GATA1*, *GATA2*, *GNAS*, *HRAS*, *IDH1*, *IDH2*, *IKZF1*, *JAK2*, *JAK3*, *KDM6A*, *KIT*, *KMT2A*, *KRAS*, *MPL*, *MYD88*, *NOTCH1*, *NPM1*, *NRAS*, *PDGFRA*, *PHF6*, *PTEN*, *PTPN11*, *RAD21*, *RUNX1*, *SETBP1*, *SF3B1*, *SMC1A*, *SMC3*, *SRSF2*, *STAG2*, *TET2*, *TP53*, *U2AF1*, *WT1*, and *ZRSR2*. In patients sequenced after 2020 at Moffitt, somatic mutations were assessed using 98-gene Moffitt Myeloid Action Panel that targets 98 genes with a detection threshold of 3%. These genes include *ABL1*, *ANKRD26*, *ASXL1*, *ASXL2*, *ATM*,

ATRX, *BCOR*, *BCORL1*, *BRAF*, *BRCC3*, *CALR*, *CBL*, *CBLB*, *CBLC*, *CCND2*, *CDKN2A*, *CEBPA*, *CHEK2*, *CREBBP*, *CSF3R*, *CSMD1*, *CSNK1A1*, *CTCF*, *CUX1*, *DDX41*, *DHX15*, *DNMT3A*, *ELANE*, *ETNK1*, *ETV6*, *EZH2*, *FANCA*, *FANCL*, *FLT3*, *GATA1*, *GATA2*, *GNAS*, *GNB1*, *HNRNPK*, *HRAS*, *IDH1*, *IDH2*, *IKZF1*, *JAK1*, *JAK2*, *JAK3*, *KDM6A*, *KIT*, *KMT2A*, *KRAS*, *LUC7L2*, *MECOM*, *MET*, *MLL2*, *MPL*, *MYC*, *NF1*, *NOTCH1*, *NOTCH2*, *NPM1*, *NRAS*, *PAX5*, *PDGFRA*, *PHF6*, *PIGA*, *PML*, *PPM1D*, *PTPN11*, *RAD21*, *RAF1*, *RB1*, *RBBP6*, *RPS19*, *RTEL1*, *RUNX1*, *SAMD9*, *SAMD9L*, *SBDS*, *SETBP1*, *SF3B1*, *SH2B3*, *SMC1A*, *SMC3*, *SOS1*, *SRP72*, *SRSF2*, *STAG1*, *STAG2*, *STAT3*, *STAT5B*, *TERC*, *TERT*, *TET2*, *TP53*, *U2AF1*, *WT1*, *ZBTB7A*, and *ZRSR2*. A subset of molecular analysis was performed using TruSeq (Illumina, San Diego, CA), which includes 31 common myeloid targeted genes, in patients who were diagnosed before 2015. Older NGS tests only reported presence or absence of the mutation and its location without variant allele frequency (VAF) information. Weill Cornell's NGS panel has a detection limit of 5% VAF. Memorial Healthcare System uses OnkoSight Advanced NGS Myeloid Panel (GenPath, Elmwood Park, NJ) that has a detection limit of 4%. In a subset of the total cohort, *NPM1* MRD testing was performed using a propriety NGS assay by Invivoscribe. The analytic sensitivity of this assay is 1×10^{-5} mutant alleles per total alleles. In another subset of patients, *NPM1* MRD was assessed using quantitative reverse transcription polymerase chain reaction. Using the delta-delta Ct method relative to a plasmid calibrator, *NPM1* variant transcripts are quantified and reported as a normalized ratio to *ABL1* transcripts present in the sample. The limit of detection of this assay is 1:100 000 cells (0.001%). These were performed at a reference laboratory, ARUP Laboratories in Salt Lake City, UT. Timing of MRD collection varied between patients and their treatment course.

Statistical analysis

Clinical variables were summarized using descriptive statistics. Fisher exact test and χ^2 test were used to assess the association between categorical variables. The Wilcoxon rank-sum test was used to compare the 2 groups' continuous variables. A 2-sided *P* value of <5% was considered statistically significant. The OS and relapse-free survival (RFS) were estimated by the Kaplan-Meier method and calculated from the date of diagnosis.

Multivariate analyses were conducted using Cox proportional hazard regression model. Statistical analysis was performed using R version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria).¹¹

Results

Patient characteristics

A total of 233 patients with AML (111 male and 122 female) with *NPM1* mutation were included in this analysis (Table 1). The median age at the time of diagnosis was 63 years (range, 22-86). Clinically, patients with dAML formed the majority at 81.1% (189/233) followed by 13.3% (31/233) sAML, and 4.7% (11/233) therapy-related AML. According to the ELN 2017 criteria, 137 patients (58.8%) had favorable-risk, 68 (29.2%) had intermediate-risk, and 22 (9.4%) had adverse-risk disease. A total of 43 patients (18.5%) had sMut (any ≥ 1 of the following: *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and *ZRSR2*), of whom

Table 1. Baseline characteristics of the study cohort

	All patients, n (%)	sMut, n (%)	No sMut, n (%)	P value
Number of patients	233 (100)	43 (18.5)	190 (81.5)	
Gender				
Male	111 (47.6)	27 (62.8)	84 (44.2)	.029
Female	122 (52.4)	16 (37.2)	106 (55.8)	
Age				
<60 y	97 (41.6)	6 (14.0)	91 (47.9)	<.001
≥60 y	136 (58.4)	37 (86.0)	99 (52.1)	
Median at diagnosis, n (range)	63 (22-86)	68 (34-86)	62 (22-80)	
Type of AML				
De novo	189 (81.1)	35 (81.4)	154 (81.1)	1.000
Secondary	31 (13.3)	5 (11.6)	26 (13.7)	1.000
Therapy-related	11 (4.7)	3 (7.0)	8 (42.0)	.431
Unknown	2 (0.9)	0	2 (1.1)	
ELN 2017 Risk				
Favorable	137 (58.8)	30 (69.8)	107 (56.3)	.124
Intermediate	68 (29.2)	5 (11.6)	63 (33.2)	.005
Adverse	22 (9.4)	7 (16.3)	15 (7.9)	.143
Unknown	6 (2.6)	1 (2.3)	5 (2.6)	
ELN 2022 Risk				
Favorable	119 (51.1)	26 (60.5)	93 (48.9)	.182
Intermediate	88 (37.8)	9 (20.9)	79 (41.6)	.014
Adverse	22 (9.4)	7 (16.3)	15 (7.9)	.143
Unknown	4 (1.7)	1 (2.3)	3 (1.6)	
Cytogenetics				
Favorable	1 (0.4)	0	1 (0.5)	1.000
Intermediate	207 (88.8)	35 (81.4)	172 (90.5)	.106
Adverse	8 (3.4)	2 (4.7)	6 (3.2)	.643
Unknown	17 (7.3)	6 (14.0)	11 (5.8)	
Most common mutations				
<i>DNMT3A</i>	114 (48.9)	15 (34.9)	99 (52.1)	.044
<i>FLT3-ITD</i>	79 (33.9)	11 (25.6)	68 (35.8)	.218
<i>TET2</i>	67 (28.7)	17 (39.5)	50 (26.3)	.095
<i>IDH1</i>	37 (15.8)	9 (20.9)	28 (14.7)	.355
<i>FLT3-TKD</i>	35 (15.0)	4 (9.3)	31 (16.3)	.345
<i>NRAS</i>	29 (12.4)	3 (6.9)	26 (13.7)	.373
<i>IDH2</i>	27 (11.6)	6 (13.9)	21 (11.1)	.600
sMut	43 (18.5)	43 (100)	0	
Mutation burden, average	3.4	4.1	3.3	
Treatment				
Frontline				
Intensive	177 (76.0)	26 (60.5)	151 (79.5)	.016
Nonintensive	50 (21.5)	16 (37.2)	34 (17.9)	.008
No treatment	6 (2.6)	1 (2.3)	5 (2.6)	1.000
Lines of therapy, average	2	2	2	
Allo-HCT	109 (46.8)	15 (34.9)	94 (49.5)	.093

sMut: *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and *ZRSR2*.

30 (69.8%) had ELN 2017–defined favorable-risk disease. When reclassified based on ELN 2022, a total of 19 patients had risk escalation from favorable to intermediate, and 2 patients with previously unknown risk were now intermediate. This is because the updated ELN no longer takes allelic ratio of *FLT3-ITD* into account. The most frequent comutations in our cohort included *DNMT3A* (114/233 [48.9%]), *FLT3-ITD* (79/233 [33.9%]), *TET2* (67/233 [28.7%]), *IDH1* (37/233 [15.8%]), *FLT3-TKD* (35/233 [15%]), *NRAS* (29/233 [12.4%]), and *IDH2* (27/233 [11.6%]) (Figure 1). Among patients with sMut, the most common sMutS were *SRSF2* (15/43 [34.9%]) and *ASXL1* (8/43 [18.6%]). Among patients without sMut, common comutations included *DNMT3A* (99/190 [52.1%]), *FLT3-ITD* (68/190 [35.8%]), *TET2* (50/190 [26.3%]), *IDH1* (28/190 [14.7%]), *FLT3-TKD* (31/190 [16.3%]), *NRAS* (26/190 [13.7%]), and *IDH2* (21/190 [11.1%]). Of note, there were 6 patients with *TP53* mutations, and none of them were categorized as having sMut based on their comutations. The average number of mutations in the sMut and no sMut cohorts was 4.1 and 3.3, respectively (Table 1). *NPM1* VAF was available in 151 patients (64.8%) with a median of 26.2% (range, 2.37%-62%). Among patients with sMut with available *NPM1* VAF (n = 31), median *NPM1* VAF was 19% (range, 3%-49.6%). The majority of patients (177/233 [76%]) received frontline intensive chemotherapy, and 46.8% (109/233) proceeded to allogeneic hematopoietic cell transplantation (allo-HCT). Fewer patients with sMut received frontline intensive chemotherapy than those without (26/43 [60.5%] vs 151/190 [79.5%]; $P = .016$), and more received nonintensive regimens (37.2% [16/43] vs 17.9% [34/190]; $P = .008$). A total of 15 out of 43 (~35%) of the sMut cohort received transplant compared with 94 of 190 (~50%) in those without sMut ($P = .093$).

Responses by secondary mutations status

Patients with sMut achieved a CR/CR with incomplete hematologic recovery (CRI) rate of 69.7% (30/43) compared with 85.3% (162/190) in those without sMut ($P = .025$) (Table 2). Relapse occurred in 34.9% (15/43) and 36.8% (70/190) of sMut and no sMut cohorts, respectively ($P = .862$). There was a higher rate of primary refractory disease in patients with sMut (13/43 [30.2%] vs 20/190 [10.5%]; $P = .003$). Posttreatment MRD testing for mutant *NPM1* was performed in 95 of 233 patients (40.8%) and was negative in 57 of 95 (60%). The proportion of posttreatment MRD negativity between sMut (9/43 [20.9%]) and non-sMut (48/190 [25.3%]) was not significantly different ($P = .695$) (Table 2).

Impact of sMut on survival

The median OS of the entire cohort was 35.5 months with a median follow-up of 52.1 months. Those who harbored sMut (n = 43) had worse OS than those without sMut (n = 190) (15.3 months vs 43.7 months; $P = .002$) (Figure 2A). Median RFS was 22.1 months in sMut (n = 43) and 37.5 months in those without sMut (n = 190) ($P = .335$). Focusing only on patients with ELN 2017 favorable-risk disease (n = 137), OS was 14.7 months vs not reached for those with sMut (n = 30) and those without sMut (n = 107), respectively ($P < .0001$) (Figure 2B). In this same subset of patients with ELN 2017 favorable risk, RFS was 14.7 months in sMut (n = 30) and 37.5 months in no sMut (n = 107) ($P = .124$). Similar findings were observed when patients were reclassified according to ELN 2022. Those with ELN 2022 favorable risk (n = 119) had significantly shorter OS if they had sMut (n = 26) than those with no sMut (n = 93) (14.7 months vs not reached; $P < .0001$). In addition, omitting *FLT3-ITD* from the




Figure 1. Molecular landscape of AML with *NPM1* mutation. *NPM1* is frequently comutated with *DNMT3A* (48.9%), *FLT3-ITD* (33.9%), and *TET2* (28.7%).

Table 2. Outcomes of the study cohort

	All patients n (%)	sMut n (%)	No sMut n (%)	P value
Response				
CR/CRi	192 (82.4)	30 (69.7)	162 (85.3)	.025
PR/NR	38 (16.3)	10 (23.3)	28 (14.7)	.176
Unknown	3 (1.3)	3 (7.0)	0	-
Best <i>NPM1</i> MRD by PCR				
Positive	38 (16.3)	4 (9.3)	34 (17.9)	.251
Negative	57 (24.5)	9 (20.9)	48 (25.3)	.695
Unknown	138 (59.2)	30 (69.8)	108 (56.8)	-
Median OS, mo	35.5	15.3	43.7	

sMut: *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and *ZRSR2*. PCR, polymerase chain reaction.

sMut cohort or the entire cohort did not change the adverse prognostic impact of sMut in all risks or favorable risk only. Univariate analysis demonstrated that age ≥ 60 years at diagnosis, nonintensive frontline treatment (compared with intensive treatment), no treatment, and sMut negatively affected OS (age ≥ 60 years; hazard ratio [HR], 2.63; 95% confidence interval [CI], 1.47-4.69; $P = .001$; nonintensive frontline treatment: HR, 3.67; 95% CI, 2.16-6.24; $P < .001$; no treatment: HR, 57.66; 95% CI, 14.48-229.5; $P < .001$; sMut: HR, 2.97; 95% CI, 1.77-4.96; $P < .001$). Patients who received allo-HCT had better outcomes than those who did not (HR, 0.31; 95% CI, 0.18-0.53; $P < .001$). In multivariate analysis, sMut retained independent negative prognostic significance on OS (HR, 2.95; 95% CI, 1.56-5.59; $P < .001$) along with those who did not get intensive treatment or allo-HCT

(Table 3). Among patients with sMut with available *NPM1* VAF data ($n = 31$), those with *NPM1* VAF $\leq 20\%$ ($n = 16$) had a shorter median OS of 7.9 months compared with 42.4 months ($P = .014$) in those with VAF $>20\%$ ($n = 15$). Among patients ($n = 120$) without sMut, the median *NPM1* VAF was 27.4% (range, 2.37%-62%), and there was no significant difference in OS between those with VAF $\leq 20\%$ ($n = 44$) and those with VAF $>20\%$ ($n = 76$; $P = .39$).

Impact of *NPM1* MRD on outcomes

Patients who achieved MRD negativity ($n = 57$) had longer OS than those with MRD positivity ($n = 38$) (MRD negative median OS (mOS), 73.9 months vs MRD positive mOS, 34.5 months; $P = .015$). Similarly, MRD negativity predicted longer OS in the overall sMut subset, although patient numbers were low ($n = 13$ [9 MRD negative and 4 MRD positive]; mOS, 73.9 vs 12.3 months; $P = .017$). Among patients with favorable-risk sMut, MRD negativity also predicted longer survival ($n = 8$ [6 MRD negative and 2 MRD positive]; mOS, 27.3 vs 10.5 months; $P = .009$).

Effects of age on response and survival in patients with *NPM1*

Lower CR/CRi rates were observed in patients with sMut within both age cohorts (age < 60 years CR/CRi, 5/6 [83.3%] with sMut vs 82/91 [90.1%] without sMut; age ≥ 60 years CR/CRi, 25/37 [67.6%] vs 80/99 [80.8%]) (Table 4). Among patients aged ≥ 60 years, those who harbored sMut ($n = 37$) fared significantly worse than patients without sMut ($n = 99$) (12.6 months vs 26.1 months; $P < .01$) (Figure 3A). In patients aged < 60 years, there was no significant difference in OS between those with sMut ($n = 6$) and those without non-sMut ($n = 91$), although the numbers were quite small. Median OS was not reached for either group. Similarly, in the




Figure 2. OS by sMut status of the entire cohort and favorable risk subset. (A) Median OS was shorter in patients with *NPM1*-mutated AML with sMut than those without sMut (15.3 months vs 43.7 months; $P = .002$). (B) When restricted to only patients with favorable risk, median OS was 14.7 months vs not reached ($P < .0001$).

Table 3. Univariate (log-rank) and multivariate (Cox regression) analyses of patients with *NPM1*+ ELN favorable-risk AML

Variable	OS					
	Univariate			Multivariate		
	HR	95% CI	P value	HR	95% CI	P value
Age ≥60 y at diagnosis	2.63	1.47-4.69	.001	0.94	0.45-1.95	.869
Type of AML*						
sAML	1.05	0.13-8.42	.964	0.83	0.10-7.01	.868
Therapy-related AML	5.84	0.64-53.13	.117	5.70	0.61-52.81	.126
<i>FLT3-ITD</i> status	1.31	0.67-2.58	.434	1.27	0.61-2.64	.528
Mutation burden >3	1.21	0.74-1.98	.438	1.07	0.62-1.87	.810
sMut	2.97	1.77-4.96	<.001	2.95	1.56-5.59	<.001
Frontline treatment†						
Nonintensive	3.67	2.16-6.24	<.001	2.71	1.44-5.13	.002
No treatment	57.66	14.48-229.5	<.001	84.03	16.76-421.3	<.001
Received allo-HCT	0.31	0.18-0.53	<.001	0.36	0.20-0.66	.001

sMut: *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and *ZRSR2*.

*Reference level is de novo AML.

†Reference level is intensive treatment.

ELN 2017 subset of patients with favorable risk, OS was significantly shorter in patients age ≥60 years with sMut (n = 28) than in those without (n = 56) ($P < .005$), and this was not observed among patients aged <60 years ($P = .31$) (Figure 3B). When analyses were repeated using the age cutoff of 70 years for the entire cohort, among those aged <70 years (n = 160), the median OS were not reached and 35 months for those without (n = 137) and those with sMut (n = 23), respectively ($P = .1471$). Among patients aged ≥70 years (n = 73), the median OS were 21.1 and 10 months for those without (n = 53) and those with sMut (n = 20), respectively ($P = .0115$). In the favorable-risk subset, sMut status significantly affected both age cohorts (age <70 years [n = 92]: no sMut [n = 75] median OS, not reached vs sMut [n = 17] median OS, 22.2 months; $P = .0059$; age ≥70 years [n = 45] no sMut [n = 32] median OS, 24.4 months vs sMut [n = 13] median OS, 10 months; $P = .0036$).

Discussion

NPM1 mutation is generally considered a favorable prognostic marker in AML.⁶ Only selected co-occurring mutations such as

FLT3-ITD or *TP53* escalate the risk under ELN 2017. *ASXL1* and *RUNX1*, 2 of the 9 mutations highly specific for sAML, are the only sMut in ELN 2017 that defines adverse-risk disease.⁷ More recent reports, including the seminal study by Lindsley et al, provided strong evidence of a poor prognosis conferred by sMut in AML.^{4,10} Based on this, 1 major update in ELN 2022 is the inclusion of all sMut being categorized as adverse risk. However, it was recommended that sMut should not currently be used as an adverse prognostic marker in the setting of favorable-risk AML subtypes, pending further evidence.⁶

In this multicenter study of 233 patients with mutant *NPM1* at diagnosis, we demonstrated that the presence of co-occurring sMut was strongly associated with adverse prognosis, including within the subset of patients with ELN 2017–defined favorable-risk disease. The facts that we observed an incidence of co-occurring sMut of ~20% in *NPM1* mutant AML and that most of these patients had ELN 2017 favorable-risk AML suggest that a substantial proportion of traditionally defined patients with favorable-risk AML indeed have poor outcomes based on the presence of these secondary-like mutations, negating the potentially beneficial impact of mutant *NPM1*. Within the sMut cohort, the rate of primary refractory disease was higher than in the non-sMut cohort, likely contributing to worse OS in this cohort, a finding consistent with prior studies showing a correlation between sMut and lower CR rates.¹⁰

We next looked at *NPM1* VAF at diagnosis among patients with sMut and those with non-sMut together and separately to see whether it alters survival. Using *NPM1* VAF threshold of 20%, there was no difference in OS among patients without sMut ($P = .39$). Surprisingly, in the sMut group, lower VAF (n = 16) had worse outcome than higher VAF (n = 15) (VAF ≤20% mOS, 7.9 months vs VAF >20% mOS, 42.4 months; $P = .014$). A prior study showed higher *NPM1* VAF (> 40%) conferred worse outcomes in de novo AML.¹² It is unclear whether the lower *NPM1* VAF in this population may have contributed to the worse outcomes observed. It is possible in the setting of sMut, low *NPM1* VAF represents a smaller

Table 4. Response rates by age and sMut

	Patients n (%)	sMut n (%)	No sMut n (%)	P value
Age <60 y at diagnosis	97	6	91	
CR/CRi	87 (89.7)	5 (83.3)	82 (90.1)	1.000
PR/NR	9 (9.3)	0	9 (9.9)	-
Unknown	1 (1.0)	1 (16.7)	0	-
Age ≥60 y at diagnosis	136	37	99	
CR/CRi	105 (77.2)	25 (67.6)	80 (80.8)	.245
PR/NR	29 (21.3)	10 (27.0)	19 (19.2)	-
Unknown	2 (1.5)	2 (5.4)	0	-

sMut: *ASXL1*, *BCOR*, *EZH2*, *RUNX1*, *SF3B1*, *SRSF2*, *STAG2*, *U2AF1*, and *ZRSR2*.

PR/NR: partial response/no response




Figure 3. OS by age and sMut status of the entire cohort and favorable risk subset. (A) Median OS was shorter in older patients (age ≥ 60) with *NPM1*-mutated AML with sMut than those without sMut. (12.6 months vs 26.1 months; $P < .01$). (B) When restricted to only patients with favorable risk, median OS was again significantly shorter in the older, sMut subset.

prognostic role of this favorable-risk mutation and a larger adverse impact from co-occurring mutations. However, our sample size is small precluding more extensive analysis. Further studies are needed to address this question more definitively.

Older age is also known to have an important prognostic role in patients with *NPM1*-mutated AML. For example, among patients aged ≥ 55 years treated with intensive chemotherapy who were enrolled onto Southwest Oncology Group and UK National Cancer Research Institute/Medical Research Council clinical trials, *NPM1*-positive/*FLT3-ITD*-negative genotype was a favorable prognostic factor in patients aged ≤ 65 years but had no significant prognostic impact in those aged > 65 years.¹³ In this study of patients with *NPM1*, those who carried sMut were generally older than patients without sMut, indicating age as a contributor to the worse outcomes among patients with sMut but clearly not the only driver. Consequently, they are also less likely to receive frontline intensive chemotherapy and subsequent allo-HCT. Interestingly, the negative prognostic impact of sMut in patients with mutant *NPM1* was not observed in patients aged < 60 years, raising the possibility that sMut is a less important prognostic factor in younger patients, although the numbers of younger patients with sMut in our study were very low, precluding any definitive conclusions in younger patients exclusively.

MRD in AML is becoming increasingly important for the assessment of response, monitoring, and prognosis.¹⁴ Different methods of measuring MRD exist, and its detection predicts increased risk of

relapse and decreased survival.^{15,16} Ivey et al assessed a large cohort of patients with *NPM1*-mutated AML with MRD data available ($n = 194$) after 2 cycles of chemotherapy using quantitative polymerase chain reaction with a median sensitivity level of 10^{-5} . They found those with undetectable MRD had a significantly lower cumulative incidence of relapse (3-year, 30% vs 80%) and higher OS (3-year, 75% vs 24%) than those with detectable MRD.¹⁶ In this study, patients who achieved *NPM1* MRD negativity had significantly longer OS within the entire cohort as well as within the sMut group itself (mOS, 73.9 vs 12.3 months; $P = .017$). Unfortunately, the number of patients in our data set who had postremission MRD testing was low, because widescale implementation of such testing is relatively recent. A lack of standardization as to the precise timing of MRD testing in our patients also might have affected any interpretation of the ultimate impact of this testing among patients with sMut. Nonetheless, the observation that *NPM1* MRD negativity in patients with sMut suggests the possibility of identifying subsets of patients with favorable outcomes in an otherwise poor-risk disease. Larger scale studies with more standardized MRD testing (and timing of testing) to confirm these observations are warranted.

In summary, our findings demonstrate that the co-occurrence of sMuts is not uncommon in *NPM1*-mutant AML, conferring a significantly detrimental effect on survival of such patients that is not well recognized by the current classification and prognostic systems. Our data not only support the recent ELN 2022 recommendations of sMut as an adverse prognostic indicator but strongly suggest that

the risk classification of AML be further refined to consider the presence of sMut_s as adverse risk in the presence of mutant *NPM1*. Such refinement would likely have significant implications on the management of AML along with clinical trial design, and additional studies should be undertaken to further strengthen and confirm these considerations to ultimately assist in the best utilization of current therapy and development of future therapies for AML.

Acknowledgment

The authors thank the patients and the supporting staff at each of the participating institution.

Authorship

Contribution: J.L. and O.C. designed the study; O.C., N.A., H.T., A.E., S.B., J.G., C.H., Y.D., L.Z., M.H., J.S., S.Y., A.K., E.P., P.D., D.S., K.S., R.K., and J.L. collected and analyzed the data; O.C. wrote the first draft; and all authors critically reviewed the first draft, provided edits, and approved the final version of the manuscript.

Conflict-of-interest disclosure: O.C. reports honoraria for consulting from Bristol Myers Squibb (BMS) and research funding from Jazz Pharmaceuticals. M.H. reports consulting and/or other fees from Adaptive Biotechnologies, Amgen, Aptitude Health, Blueprint Oncology, Celgene, Decibio, Diaceutics, Guidepoint, Seattle Genetics, Stemline, Tegus, Janssen, and BMS. C.T. is a current equity holder in private company, AbbVie. A.K. reports consulting and/or other fees from AbbVie, Blueprint Medicines, Celgene, Constellation, CTI Biopharma, Incyte, Novartis, Prelude, Protagonist, and Sierra Oncology; and grant funding from BMS, Constellation, Novartis, Prelude, Protagonist, and Sierra Oncology. E.P. reports consulting and/or other fees from Blueprint Medicines,

Novartis, and Taiho Oncology. G.R. reports consulting and/or other fees from Actinium, AbbVie, Agios, Amgen, Astellas, AstraZeneca, BMS, Blueprint Medicines, bluebird bio, GlaxoSmithKline, Janssen, Jasper, Jazz, MEI Pharma, Mesoblast, Novartis, Pfizer, Syndax, and Takeda. P.D. reports consulting and/or other fees from Janssen, Servier, AbbVie, and BMS. D.S. reports consulting and/or other fees from Aprea Therapeutics, Avencell Europe GmbH, bluebird bio, BMS, Gilead Sciences, Incyte, Intellia Therapeutics, Janssen Global Services, Jazz Pharmaceuticals, Kite Pharma, Molecular Partners AG, Novartis, Servier Pharmaceuticals, Shattuck Labs, Syndax Pharmaceuticals, Syros Pharmaceuticals, and Takeda Pharmaceutical. K.S. reports consulting and/or other fees from Arog, Astellas Pharma, BerGenBio, BMS, Curis, Gilead Sciences, Mablytics, Novartis, Pfizer Pharmaceuticals, and Incyte. R.K. reports consulting and/or other fees from AbbVie, Acceleron Pharma, Agios Pharmaceuticals, Alexion Pharmaceuticals, BMS, Celgene, CTI Biopharma, Daiichi Sankyo, Geron, Janssen Biotech, Jazz Pharmaceuticals, Novartis Pharma, Pfizer, PharmaEssentia, Servier Pharmaceuticals LLC, Taiho Oncology, and Takeda Oncology. J.L. reports consulting and/or other fees from AbbVie, Boxer Capital, BMS, Jasper Therapeutics, Jazz Pharmaceuticals, Novartis Pharma, and Servier Pharmaceuticals. The remaining authors declare no competing financial interests.

ORCID profiles: O.C., [0000-0003-2872-1961](https://orcid.org/0000-0003-2872-1961); H.T., [0000-0002-2498-9111](https://orcid.org/0000-0002-2498-9111); C.H., [0000-0002-6663-3535](https://orcid.org/0000-0002-6663-3535); L.Z., [0000-0002-1463-1458](https://orcid.org/0000-0002-1463-1458); A.K., [0000-0002-9040-7415](https://orcid.org/0000-0002-9040-7415); G.R., [0000-0002-0384-3658](https://orcid.org/0000-0002-0384-3658); D.S., [0000-0003-0504-8233](https://orcid.org/0000-0003-0504-8233); R.K., [0000-0002-1876-5269](https://orcid.org/0000-0002-1876-5269).

Correspondence: Onyee Chan, Department of Malignant Hematology, Moffitt Cancer Center, 12902 Magnolia Dr, Tampa, FL 33612; email: onyee.chan@moffitt.org.

References

1. Falini B, Brunetti L, Sportoletti P, Martelli MP. NPM1-mutated acute myeloid leukemia: from bench to bedside. *Blood*. 2020;136(15):1707-1721.
2. Heath EM, Chan SM, Minden MD, Murphy T, Shlush LI, Schimmer AD. Biological and clinical consequences of NPM1 mutations in AML. *Leukemia*. 2017;31(4):798-807.
3. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-2405.
4. Papaemmanuil E, Gerstung M, Bullinger L, et al. Genomic classification and prognosis in acute myeloid leukemia. *N Engl J Med*. 2016;374(23):2209-2221.
5. Thiede C, Koch S, Creutzig E, et al. Prevalence and prognostic impact of NPM1 mutations in 1485 adult patients with acute myeloid leukemia (AML). *Blood*. 2006;107(10):4011-4020.
6. Dohner H, Wei AH, Appelbaum FR, et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022;140(12):1345-1377.
7. Dohner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017;129(4):424-447.
8. Eisfeld AK, Kohlschmidt J, Mims A, et al. Additional gene mutations may refine the 2017 European LeukemiaNet classification in adult patients with de novo acute myeloid leukemia aged <60 years. *Leukemia*. 2020;34(12):3215-3227.
9. Dunlap JB, Leonard J, Rosenberg M, et al. The combination of NPM1, DNMT3A, and IDH1/2 mutations leads to inferior overall survival in AML. *Am J Hematol*. 2019;94(8):913-920.
10. Lindsley RC, Mar BG, Mazzola E, et al. Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood*. 2015;125(9):1367-1376.
11. R Core Team. R: A Language and Environment for Statistical Computing. 4.0.2. R Foundation for Statistical Computing; 2020.
12. Patel SS, Kuo FC, Gibson CJ, et al. High NPM1-mutant allele burden at diagnosis predicts unfavorable outcomes in de novo AML. *Blood*. 2018;131(25):2816-2825.

13. Ostronoff F, Othus M, Lazenby M, et al. Prognostic significance of NPM1 mutations in the absence of FLT3-internal tandem duplication in older patients with acute myeloid leukemia: a SWOG and UK National Cancer Research Institute/Medical Research Council report. *J Clin Oncol*. 2015;33(10):1157-1164.
14. Heuser M, Freeman SD, Ossenkoppele GJ, et al. 2021 Update on MRD in acute myeloid leukemia: a consensus document from the European LeukemiaNet MRD Working Party. *Blood*. 2021;138(26):2753-2767.
15. Schuurhuis GJ, Heuser M, Freeman S, et al. Minimal/measurable residual disease in AML: a consensus document from the European LeukemiaNet MRD Working Party. *Blood*. 2018;131(12):1275-1291.
16. Ivey A, Hills RK, Simpson MA, et al; UK National Cancer Research Institute AML Working Group. Assessment of minimal residual disease in standard-risk AML. *N Engl J Med*. 2016;374(5):422-433.