

# A phase 1 study of the irreversible FLT3 inhibitor FF-10101 in relapsed or refractory acute myeloid leukemia

Tracking no: ADV-2023-010619R2

Mark Levis (Johns Hopkins University, United States) Alexander Perl (Abramson Cancer Center, University of Pennsylvania, United States) Gary Schiller (David Geffen School of Medicine at UCLA, United States) Amir Fathi (Massachusetts General Hospital / Harvard Medical School, United States) Gail Roboz (Weill Medical College of Cornell University, United States) Eunice Wang (Roswell Park Comprehensive Cancer Center, United States) Jessica Altman (Northwestern University, United States) Trivikram Rajkhowa (Johns Hopkins University, United States) Makoto Ando (Fujifilm, Japan) Takeaki Suzuki (Fujifilm Pharmaceuticals, USA, United States) Ruth Subach (Fujifilm Pharmaceuticals, USA, United States) Gary Maier (Fujifilm Pharmaceuticals, USA, United States) Timothy Madden (Fujifilm Pharmaceuticals, USA, United States) Mary Johansen (FUJIFILM Pharmaceuticals U.S.A., Inc., United States) Kin Cheung (Fujifilm Pharmaceuticals, USA, United States) Michael Kurman (FUJIFILM Pharmaceuticals U.S.A., Inc., United States) Catherine Smith (University of California (San Francisco), United States)

## Abstract:

FLT3 tyrosine kinase inhibitors (TKIs) have clinical efficacy for patients with FLT3-mutated AML (acute myeloid leukemia), but their impact is limited by resistance in the setting of monotherapy and by tolerability problems when used in combination therapies. FF-10101 is a novel compound that covalently binds to a cysteine residue near the active site of FLT3, irreversibly inhibiting receptor signaling. It is effective against most FLT3 activating mutations, and unlike other inhibitors is minimally vulnerable to resistance induced by FLT3 ligand (FL). We conducted a phase 1 dose escalation study of oral FF-10101 (NCT03194685) in patients with relapsed and/or refractory AML, the majority of whom harbored FLT3-activating mutations and/or had prior exposure to FLT3 inhibitors. Fifty-four participants enrolled in cohorts ranging from 10 to 225 mg per day and 50 to 100 mg twice daily (BID). The dose limiting toxicities (DLT) were diarrhea and QT prolongation. Among 40 response-evaluable participants, the composite complete response rate was 10%, and the overall response rate (including partial responses) was 12.5%, including patients who had progressed on gilteritinib. 56% of participants had prior exposure to FLT3 inhibitors. The recommended phase 2 dose (RP2D) was 75 mg BID. FF-10101 potentially represents a next-generation advance in the management of FLT3-mutated AML.-

**Conflict of interest:** COI declared - see note

**COI notes:** M.L. received consultancy fees from Abbvie, Astellas, Amgen, Bristol-Myers-Squibb, Daiichi-Sankyo, Jazz, Pfizer, and Kite, and research support from FujiFilm. G.S. received research support from FujiFilm. A.T.F. received research support from Servier, Bristol-Myers-Squibb, and Abbvie, and consulting fees from Servier, Bristol-Myers-Squibb, Abbvie, Genentech, Amgen, Takeda, Astellas, Mablytics, EnClear, Orum, Forma, Daiichi Sankyo, Pfizer, Rigel, Autolus, Menarini,, and Remix. G.R. received research support from Janssen, and consultancy fees from Abbvie, Amgen, Argenx, Astra Zeneca, Bluebird Bio, Blueprint Medicines, Bristol-Myers Squibb, Caribou Biosciences, Celgene, Daiichi Sankyo, Ellipses Pharma, GlaxoSmithKline, Janssen, Jasper Pharmaceuticals, Jazz Pharmaceuticals, Molecular Partners, Novartis, Pfizer, Roche, Syndax, and Takeda. E.S.W. received consultancy fees from Abbvie, Astellas, BMS, Daiichi Sankyo, Genentech, Gilead, GlaxoSmithKline, Janssen, Jazz, Kite, Kura, Novartis, NuProbe, Pfizer, Rigel, Sellas, and Sumitomo Pharma. A.P. received consultancy fees from Abbvie, Astellas, BerGenBio, Daiichi Sankyo, Genentech, Immunogen, Foghorn, Actinium, and Forma, and research support from Abbvie, Astellas, Daiichi Sankyo, FujiFilm, and Syndax. J.A. received consultancy fees from AbbVie, Astellas Pharma, BioSight, Bluebird Bio, Curio, Daiichi Sankyo, Gilead, Kura Oncology, Kymera, Stemline Therapeutics, and Syros and research funding from AbbVie, Agios, ALX Oncology, Amgen, Amphivena, Aprea AB, Aptose Biosciences, Astellas Pharma, BioSight, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Cyclacel, Fujifilm, Immunogen, Kartos Therapeutics, Kura Oncology, Loxo, and Pfizer. T.R. has no conflicts of interest to declare. M.A., T.S., R.A.S., G.M., T.M., M.J., K.C., and M.K. are employees of FujiFilm, whose product (FF-10101) is described in this work. C.C.S received research funding from FujiFilm (clinical trial), Astellas (clinical trial) AbbVie, BMS/Celgene (clinical trial), Erasca, Revolution Medicines, and Zentaris (clinical trial), and has served on a board of advisory committee for Astellas Pharma, Daiichi Sankyo, and Genentech.

**Preprint server:** No;

**Author contributions and disclosures:** M.L. designed the study, enrolled trial participants, analyzed the data, and wrote the manuscript. A.E.P. G. S., A.T.F., G.R., E.S.W., J.A., and C.S. enrolled trial participants and edited the manuscript. T.R. performed laboratory experiments. M.A., T.S., R.A.S., G.M., T.M., M.J., and K.C. contributed to study design, analyzed the data, and edited the manuscript. M.K. was the medical monitor and edited the manuscript.

**Non-author contributions and disclosures:** No;

**Agreement to Share Publication-Related Data and Data Sharing Statement:** Email to corresponding author

**Clinical trial registration information (if any):** NCT03194685  
<https://clinicaltrials.gov/ct2/show/NCT03194685>

# A phase 1 study of the irreversible FLT3 inhibitor FF-10101 in relapsed or refractory acute myeloid leukemia

Mark Levis<sup>1\*</sup>, Alexander Perl<sup>2</sup>, Gary Schiller<sup>3</sup>, Amir T. Fathi<sup>4</sup>, Gail Roboz<sup>5</sup>, Eunice S. Wang<sup>6</sup>, Jessica Altman<sup>7</sup>, Trivikram Rajkhowa<sup>1</sup>, Makoto Ando<sup>8</sup>, Takeaki Suzuki<sup>9</sup>, Ruth Ann Subach<sup>9</sup>, Gary Maier<sup>9</sup>, Timothy Madden<sup>9</sup>, Mary Johansen<sup>9</sup>, Kin Cheung<sup>9</sup>, Michael Kurman<sup>9</sup>, Catherine Smith<sup>10</sup>

1. Department of Oncology, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD, USA
2. Department of Medicine, Division of Hematology/Oncology, Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA, USA
3. Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA
4. Massachusetts General Hospital Cancer Center, Harvard Medical School, Boston, MA, USA
5. Weill Cornell Medicine and the New York Presbyterian Hospital, New York, NY, USA
6. Roswell Park Comprehensive Cancer Center, Buffalo, NY, USA
7. Department of Medicine, Division of Hematology/Oncology, Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Chicago, IL, USA
8. FUJIFILM Corporation, Tokyo, Japan
9. FUJIFILM Pharmaceuticals U.S.A., Inc., Cambridge, MA, USA
10. Department of Medicine, Division of Hematology/Oncology, Helen Diller Comprehensive Cancer Center, University of California at San Francisco, San Francisco, CA, USA

Running title: Phase 1 study of FF-10101

Word count:

Abstract- 204

Text- 3361

\*Corresponding Author:

Mark Levis MD PhD  
Johns Hopkins University  
1650 Orleans Street, Room 2M44  
Baltimore, MD 21287  
levisma@jhmi.edu

## Key Points:

FF-10101 is an irreversible inhibitor of FLT3 active against most types of FLT3 activating mutations

FF-10101 induced responses in patients who had progressed on other FLT3 inhibitors, as well as in some patients with wild type FLT3.

## Abstract

FLT3 tyrosine kinase inhibitors (TKIs) have clinical efficacy for patients with FLT3-mutated AML (acute myeloid leukemia), but their impact is limited by resistance in the setting of monotherapy and by tolerability problems when used in combination therapies. FF-10101 is a novel compound that covalently binds to a cysteine residue near the active site of FLT3, irreversibly inhibiting receptor signaling. It is effective against most FLT3 activating mutations, and unlike other inhibitors is minimally vulnerable to resistance induced by FLT3 ligand (FL). We conducted a phase 1 dose escalation study of oral FF-10101 (NCT03194685) in patients with relapsed and/or refractory AML, the majority of whom harbored FLT3-activating mutations and/or had prior exposure to FLT3 inhibitors. Fifty-four participants enrolled in cohorts ranging from 10 to 225 mg per day and 50 to 100 mg twice daily (BID). The dose limiting toxicities (DLT) were diarrhea and QT prolongation. Among 40 response-evaluable participants, the composite complete response rate was 10%, and the overall response rate (including partial responses) was 12.5%, including patients who had progressed on gilteritinib. 56% of participants had prior exposure to FLT3 inhibitors. The recommended phase 2 dose (RP2D) was 75 mg BID. FF-10101 potentially represents a next-generation advance in the management of FLT3-mutated AML.

## Introduction

Activating mutations of the *FLT3* gene are among the most common genetic perturbations in acute myeloid leukemia (AML) and the most common of these, internal tandem duplication (ITD) mutations, are associated with a worse prognosis.<sup>1-3</sup> A number of different FLT3 inhibitors have been shown to improve survival in patients with FLT3-mutated AML when incorporated into induction and consolidation chemotherapy, as monotherapy for relapsed/refractory disease, and as post-transplant maintenance.<sup>4-8</sup> While FLT3 inhibitors are now a part of the standard-of-care approach to AML, responses in the relapsed/refractory setting are still sub-optimal.<sup>5,9</sup> A number of resistance mechanisms have emerged,<sup>10-13</sup> and tolerability (myelosuppression, QT-prolongation, gastrointestinal, and skin toxicities) is still problematic, particularly when the drugs are given in combination therapy.<sup>14-17</sup> Given that FLT3 inhibition has clearly improved outcomes in AML, identifying a FLT3 inhibitor that is both more tolerable and less affected by resistance mechanisms offers an opportunity to improve outcomes yet further.

FF-10101 is a novel type 1 FLT3 inhibitor that contains an acryloyl group that forms a covalent bond with cysteine 695 near the ATP-binding pocket of FLT3, irreversibly inhibiting the receptor.<sup>18</sup> FF-10101 has potent *in vitro* inhibitory activity against *FLT3* internal tandem duplication (*FLT3*-ITD) and *FLT3* tyrosine kinase domain (*FLT3*-TKD) mutations, as well as against the gatekeeper, non-canonical, and extracellular activating mutations that are associated with resistance to other FLT3 inhibitors.<sup>19</sup> In addition, its irreversible binding renders the drug minimally affected by the binding of FLT3 ligand to the receptor, another mechanism of resistance to FLT3 inhibitors.<sup>18</sup> We conducted a phase 1 first-in-human study to identify a dose

of FF-10101 that could achieve sustained *in vivo* FLT3 inhibition in patients with relapsed and/or refractory (R/R) AML regardless of FLT3 mutation status in a safe and tolerable fashion.

## Methods

### *Study design*

This was a phase 1, dose-escalation study of oral FF-10101 conducted at eight centers in the U.S. Eligible patients were age  $\geq 18$  years of age with relapsed and/or refractory AML and unlikely to benefit from standard therapy (particularly approved FLT3 inhibitors) according to the local treating physician, and after consensus from all other investigators via teleconferences. The primary objectives were to define the dose limiting toxicities, determine a maximum tolerated dose, and identify a dose or doses for expansion based on *in vivo* target inhibition, pharmacokinetics, and tolerability. The original design was a phase 1 3 x 3 dose escalation study in patients with R/R AML, regardless of *FLT3* mutation status, in order to identify a recommended phase 2 dose (RP2D). This would be followed by expansion cohorts restricted to patients with FLT3 mutant R/R AML. However, the COVID-19 pandemic began (March 2020) just as the final participants in the dose-escalation phase were enrolled and the RP2D was being identified. The industry sponsor made the decision to halt the study at that time.

The trial schema is shown in Figure 1. Dose escalation of FF-10101 started at a single daily dose of 10 mg and continued up to 225 mg per day in 28-day cycles. If no dose-limiting toxicities

(DLTs) were observed in the first three patients, the subsequent three patients were treated at the next dose level, after agreement between the medical monitor and the participating investigators. If one DLT occurred at a specific dose level, three more patients were enrolled at the same dose level. If the three additional patients did not experience a DLT, the next dose level was initiated. If two or more DLTs occurred in a cohort of three to six people, the next lower dose level was declared to be the maximum tolerated dose. A DLT was defined as any grade 3 or greater non-hematologic toxicity considered at least possibly related to FF-10101 and not clearly resulting from the underlying leukemia, except for grade 3 nausea and vomiting that reverted to grade 2 or less within 72 hours, infection, bleeding or other expected direct complications of cytopenias due to underlying leukemia, grade 3 asthenia, fever, anorexia, or constipation, or alopecia of any grade. In addition, grade 3 or greater elevations of creatine kinase (CK) in the absence of clinical signs of myositis were not considered DLTs, as such changes may be due to reduced enzyme clearance caused by the drug inhibiting the function of the CSF1 receptor (like FLT3, a class III tyrosine kinase receptor).<sup>20-22</sup> To ensure there was no cardiac damage in association with elevated CK, troponin and B-type natriuretic peptide was monitored. All such changes were classified as adverse events (AEs) of special interest for review by the safety committee. Finally, differentiation syndrome was also designated as an AE of special interest (as an on-target effect), not considered a DLT.

While each dose cohort was assessed for safety and tolerability, each was also evaluated for in vivo FLT3 inhibition using an ex vivo surrogate assay, the plasma inhibitory activity (PIA) assay for FLT3.<sup>23</sup> As the trial progressed, in response to PIA assay results and pharmacokinetic data, new cohorts were opened in which participants were dosed twice daily (every 12 hours). The

study was conducted in compliance with the Declaration of Helsinki and approved by institutional review boards at each site. All participants provided informed consent. Bone marrow aspirates were collected prior to start of therapy and on day 1 of cycle 2 and beyond. Response was evaluated (in participants receiving at least 75% of doses in at least one cycle of study drug) using the Revised Recommendations of the International Working Group for AML,<sup>24</sup> with the modification that complete remission with incomplete hematologic recovery (CRh), defined as complete remission (CR) but with absolute neutrophil count (ANC) between 500/uL and 1000/uL and platelets above 50,000/uL, was included. Adverse events (AEs) were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events version 4.03. Following data lock, the primary data was directly analyzed by the corresponding author, who then wrote the manuscript without assistance from non-authors.

### *Data Sharing*

Access to deidentified participant data is provided after a proposal has been approved by an independent review committee identified for this purpose and after receipt of a signed data sharing agreement. Data and documents, including the study protocol, statistical analysis plan, clinical study report, and blank or annotated case report forms, will be provided in a secure data sharing environment. The earliest availability of data will be two years after first regulatory approval of the initial indication.

Contact: [fphucontact@fujifilm.com](mailto:fphucontact@fujifilm.com).

### *Cell lines, primary patient samples, and turnover studies*



*FLT3*-ITD-expressing cell lines Molm14 and MV411,<sup>25,26</sup> and primary AML samples were cultured as described.<sup>11,27,28</sup> *FLT3* immunoprecipitation and immunoblotting was carried out as described.<sup>11,27,28</sup> Primary blasts collected from patients with AML were provided by the Sidney Kimmel Comprehensive Cancer Center Hematologic Malignancies Cell Bank Protocol. All participants gave informed consent according to the Declaration of Helsinki under a protocol approved by the Institutional Review Board at Johns Hopkins. Blasts were isolated and viably cryopreserved as described.<sup>11,27,28</sup> FF-10101 was provided by FUJIFILM Corporation and dissolved in dimethyl sulfoxide (DMSO) at stock concentrations of 10 mM. Stocks were aliquoted into 10  $\mu$ L volumes, stored at -80°C and thawed immediately before use. For turnover studies, cell lines or blasts were cultured in the presence or absence of 50  $\mu$ g/mL cycloheximide (Sigma-Aldrich, St. Louis, MO) prior to *FLT3* immunoblotting analysis.

#### *Pharmacokinetic and pharmacodynamic studies*

At pre-determined time points during the first and second cycles, whole blood was collected into lithium-heparin tubes. Plasma was separated from the cellular fraction and stored frozen. Plasma samples were analyzed for FF-10101 concentration with a validated method, and in parallel, for PIA and *FLT3* ligand (FL) concentration, as described.<sup>11,27</sup> When feasible, circulating blasts were collected and analyzed for total and phosphorylated *FLT3* (P-*FLT3*) to directly compare with PIA results at simultaneous time points, as described.<sup>11,27</sup>

#### *Statistical analysis*

The dose-toxicity association and determination of the MTD was determined using a two parameter Bayesian logistical regression model.<sup>29</sup> All adverse events, clinical laboratory data, and pharmacokinetic and pharmacodynamic data were summarized using descriptive statistics.

## Results

### *Trial participants*

Between August 2017 and February 2021, 54 participants were enrolled and treated. The database was locked on October 27, 2021. The characteristics of the participants are displayed in Table 1. The median number of prior therapies was 3 (range 1-7), and 30 of 54 participants (55.6%) harbored a FLT3 mutation at the time of enrollment. An equal proportion (30 out of 54; 56.6%) had previously been treated with a FLT3 inhibitor, but in some participants, a previously-detected *FLT3* mutation was no longer detectable at enrollment. Other than FLT3-ITD and FLT3-TKD (D835) mutations, only two non-canonical FLT3 mutations (V491L and A680V) were detected in the participants. No “gatekeeper” mutations (at residue F691) were detected.

### *Pharmacokinetics and pharmacodynamics*

The IC<sub>50</sub> (concentration inhibiting 50% of baseline activity) of FF-10101 against FLT3 in 100% human plasma is estimated to be 20 nM (Supplementary Figure 1). Pharmacokinetic analysis indicated that following oral dosing FF-10101 has a mean half-life ( $t_{1/2}$ ) of approximately 10 hours, with absorption unaffected by food. The PIA assay was originally developed as an *ex-vivo* surrogate of FLT3 inhibition using reversible FLT3 inhibitors. While it is expected to accurately reflect FLT3 inhibition based on the concentration of inhibitor in any given plasma sample, an irreversible inhibitor is expected to have inhibitory effects after the drug has cleared from the system- depending on the turnover or  $t_{1/2}$  of the target receptor. We carried out a series of experiments to determine the  $t_{1/2}$  of both the wild type and mutant FLT3 receptor in cell lines

and in primary blasts isolated from patients (Figure 2A). The  $t_{1/2}$  of FLT3 in Molm-14 and MV4-11 cells, as well as in primary samples ranged from 0.9-5.2 hours, which is consistent with previously published studies. The turnover was only modestly decreased in the presence of FLT3 inhibition, with an approximately 50% increase in the  $t_{1/2}$  of FLT3 (*data not shown*). To confirm that the PIA assay was an appropriate surrogate for *in vivo* FLT3 inhibition by FF-10101, we collected whole blood from a participant receiving 100 mg per day FF-10101 who had circulating blasts at enrollment. We divided the sample in two, and half was used to directly isolate and analyze P-FLT3 from the circulating blasts while the other half was used to isolate plasma for the PIA assay. The results of these two parallel assays are shown in Figure 2B. Of note, total FLT3 in the primary blasts was markedly up-regulated (more so than in the Molm14 cells), presumably reflecting either increased production, decreased turnover, or both. Regardless, based on the equivalent degree of P-FLT3 inhibition in blasts versus in Molm14 cells assessed by PIA (Figure 2B, *upper panels*), we concluded that the PIA assay was an adequate surrogate for *in vivo* FLT3 inhibition by FF-10101. Based on the short half-life of the FLT3 receptor and the level of P-FLT3 visible at 8 hours after dosing, we also concluded that dosing of FF-10101 needed to be more than once daily to allow for inhibition of newly synthesized receptors. The protocol was subsequently amended to open additional cohorts with twice daily (BID) dosing.

PIA assay results for the escalating doses of FF-10101 are displayed in Figure 3A, with representative immunoblots for each dose level shown in Supplementary Figure 2. Inhibition of phosphorylated FLT3 was most effective at dose levels of 150 mg daily, 225 mg daily, 50 mg twice daily, and 75 mg twice daily. The plasma concentration of FF-10101 correlated with the

PIA results (Figure 3B), with concentrations greater than 100 ng/mL associated with more complete inhibition. Steady state concentrations of FF-10101 (after dosing on day 1 of cycle 2) are displayed in Figures 3C-D. Maximal trough suppression of P-FLT3 activity (e.g., < 5% of baseline) was achieved at FF-10101 concentrations > 200 ng/mL. Concomitant azole use was allowed after the first week of treatment with FF-10101, and more intensive PK and ECG monitoring was performed when azoles were used. However, only 10 participants across 7 dosing cohorts were treated with azoles (voriconazole, posaconazole, or isavuconazonium), and so while no apparent azole effect on FF-10101 concentration was observed, the sample number is too small to draw any firm conclusions about this drug interaction.

### *Safety and tolerability*

53 of 54 (98.1%) participants reported at least one treatment emergent adverse event (TEAE). The most frequent related TEAEs are summarized in Table 2, and Supplementary Table 1 displays all grade 3 or greater TEAEs reported for all cohorts. Six participants died while on study treatment from causes deemed by the local investigator to be unrelated to study drug: 4 from sepsis/infection, 1 from intracranial hemorrhage, and 1 from pulmonary embolus- all reported as not related to study drug. Differentiation syndrome, which has been reported in association with FLT3 inhibitor therapy,<sup>30,31</sup> occurred in 4 participants, 2 in the 75 mg BID cohort (one of whom was *FLT3* WT) 1 in the 100 QD cohort and 1 in the 100 mg BID cohort.

CK elevation was an adverse event of special interest (AESI) and occurred in 16 (29.6%) of participants. In no case of CK elevation was there any sign of myositis. Clearance of the CK

enzyme is thought to be cleared at least in part by hepatic Kupffer cells (monocyte-derived) that are in turn regulated by the CSF1R receptor.<sup>21</sup> CSF1R is a type III receptor tyrosine kinase (RTK),<sup>20</sup> structurally homologous to FLT3. Elevation of CK appears to be a class effect of type III RTK inhibitors, and has been observed in clinical studies of other FLT3 inhibitors.<sup>32,33</sup> Given the prevalence of elevated CK, as an additional safety precaution the protocol was amended to monitor cardiac ejection fraction throughout therapy. Echocardiograms were obtained at baseline and at the start of each subsequent cycle of therapy. Follow up echocardiograms were obtained from 38/54 (70%). The most common reason for no follow up echocardiogram was failure to complete one cycle of therapy. Ten of these 38 participants were admitted with adverse events (9/10 fever/sepsis and 1 with hemorrhage and differentiation syndrome) when the follow up echocardiogram was obtained, so these were not included in this evaluation. Of the 28 participants with follow up echocardiograms obtained on day 1 of cycle 2, ejection fractions (EFs) fluctuated above and below the baseline value (Supplementary Figure 3). A single participant, an 84-year-old male with a prior history of coronary artery disease, atrial fibrillation, and daunorubicin exposure, experienced a decrease in EF to below 50%.

Diarrhea is a common AE in any phase 1 AML trial. However, there was a clear association between FF-10101 dose and diarrhea, in that 1621 (76.2%) participants treated with 100mg/dose or greater (e.g., cohorts 100 mg QD, 150 mg QD, 225 mg QD, and 100 mg BID) reported grade 1-3 diarrhea, while only 733 (21.2%) participants dose with less than 100 mg/dose reported diarrhea. The frequency with which diarrhea was reported as an AE according to dose level is depicted in Supplementary Figure 4. While only 4 participants had grade 3 diarrhea, this AE rendered these higher doses poorly tolerated over weeks of therapy.

Regarding DLTs, one participant (out of seven) in the 50 mg BID cohort experienced a grade 3 maculopapular rash possibly due to FF-10101. Two out of 5 participants treated with 100 mg BID experienced DLTs. The first was a 50-year-old female who developed grade 3 diarrhea, and was subsequently admitted on cycle 1 day 8 with urosepsis and bacteremia. The study drug was stopped at that time. An echocardiogram obtained 7 days later showed a left ventricular ejection fraction of 26% (compared with 52% at baseline). On review, the cardiomyopathy was deemed stress cardiomyopathy in the setting of sepsis. Follow up echocardiograms showed recovery to a normal ejection fraction. The second participant in the 100 mg BID cohort to experience a DLT was a 68-year-old woman who reported grade 1 diarrhea and vomiting on cycle 1 day 9. She was noted to have grade 3 QT prolongation, considered probably related to FF-10101 and the study drug was stopped. She also developed grade 3 troponin increase, which was deemed not related to study drug. The participant was admitted with sepsis and progressive disease two days later and expired on cycle 1 day 11. Autopsy revealed extensive extramedullary leukemia but no evidence of cardiomyopathy. The two confirmed DLTs for this cohort, therefore, were grade 3 diarrhea and grade 3 QT prolongation, respectively.

The maximum tolerated dose (MTD) was determined to be 75 mg BID. Per protocol, the 100 mg QD, 150 mg QD, and 225 mg QD dosing levels were tolerated, but from a practical standpoint the investigators concluded that the high prevalence of grade 1-2 diarrhea rendered those dose levels generally less tolerable.

## *Efficacy*

Forty out of 54 total participants were evaluable for response. Fourteen participants did not receive at least 75% of study drug during the first cycle of therapy and were withdrawn from the study (mostly due to infection or other adverse events). These were evaluated only for safety. Among the 40 response-evaluable participants, there was 1 CR (in a FLT3-ITD participant previously treated with midostaurin), 1 CRh (in a participant with wild-type FLT3 never previously treated with a FLT3 inhibitor), 2 CRp (one in a participant with a FLT3-ITD mutation not previously treated with a FLT3 inhibitor and one in a participant with wild-type FLT3 previously treated with gilteritinib), and 1 PR (FLT3 wild type, previously treated with gilteritinib). The composite CR rate (CR + CRh + CRp) was 4/40 (10%), and the overall response rate (CR + CRh + CRp + PR) was 5/40 (12.5%). The ORR in participants with FLT3 mutations was 9.5% (2 out of 21) and in participants previously treated with a FLT3 inhibitor, it was 13.6% (3 out of 22). One participant whose disease lacked a FLT3 mutation and never had one previously achieved a CRh. Detailed characteristics of the five responding participants are summarized in Supplementary Table 2.

Of the 40 response-evaluable participants, 14 had circulating blasts at the end of the first cycle, and an evaluation bone marrow biopsy was deferred. For the remaining 26 participants, the percentage of marrow blasts after the first cycle was compared with the baseline marrow. The results are displayed in Figure 4.

In the original plan of the protocol, once FLT3-inhibitory doses were identified, expansion cohorts of participants with FLT3 mutations were to be treated with the identified doses of FF-10101. This did not take place as the sponsor has suspended development at this time.

However, a total of 11 participants with a history of *FLT3*-mutated AML were treated with doses of FF-10101 determined from PIA analysis to inhibit FLT3 *in vivo* in sustained fashion (150 mg QD, 225 mg QD, 50 mg BID, and 75 mg BID) and were evaluable for response. Amongst these 10 participants, there was 1 CR, 2 CRp, and 1 PR. Three out of 4 of these responders had been previously treated with FLT3 inhibitors.

## Discussion

In this first-in-human phase 1 study, the MTD of FF-10101 was determined to be 75 mg BID. This was also the dosing regimen that yielded the most effective and consistent suppression of FLT3 activity in tolerable fashion, and therefore is the RP2D. In pre-clinical models, FF-10101 overcomes many of the resistance mechanisms that have been observed with other FLT3 inhibitors,<sup>19</sup> and from this initial study, the drug does appear to have, in some cases, clinical activity after the failure of approved FLT3 inhibitors. These data were generated prior to the widespread use of gilteritinib, and so any conclusions about efficacy in the gilteritinib-resistance setting are very preliminary. This will require a larger study formally designed to test this concept, and clinical resistance to FF-10101 itself would be expected to eventually emerge, most likely from mutation of cysteine 695.<sup>19</sup> Importantly, we observed responses in participants with AML lacking detectable FLT3-activating mutations, a phenomenon seen with quizartinib but much less so with gilteritinib.<sup>14,34</sup> This may reflect both the irreversibility of FF-10101 as well



as its activity against the wild type receptor, which is frequently over-expressed in AML cells and requires FLT3 ligand for activation.<sup>35,36</sup>

FLT3 inhibitors are now a component of the standard-of-care management of *FLT3*-mutated AML at all stages of the disease. There are currently 4 FLT3 inhibitors in common clinical use around the world- sorafenib, midostaurin, gilteritinib, and quizartinib- and a fifth- crenolanib- is in late stages of development.<sup>37</sup> As these drugs have been incorporated into conventional treatment regimens, including combinations with induction chemotherapy, hypomethylating agents, venetoclax, and as maintenance, a number of problems have emerged. Sorafenib does not have regulatory approval for AML, and is often poorly tolerated by patients.<sup>38,39</sup> It is also inactive against *FLT3*-TKD mutations, which act as a mechanism of resistance.<sup>12</sup> Midostaurin has no single agent activity,<sup>40</sup> and is likewise poorly tolerated in general, with most patients requiring pre-treatment with an anti-emetic before every dose, and with a high cessation rate in maintenance.<sup>15,41</sup> Gilteritinib and quizartinib are well tolerated on a day-to-day basis, but both have exceptionally long terminal half-lives in patients.<sup>42,43</sup> In the setting of relapse/refractory *FLT3*-mutated AML, this long half-life was deemed advantageous for single-agent FLT3 inhibitor therapy, as inhibition of FLT3 needs to be continuous over days in order to induce a meaningful anti-leukemic effect.<sup>27</sup> However, when administered in conjunction with myelosuppressive regimens, both gilteritinib and quizartinib are associated with prolongation of aplasia.<sup>17,44,45</sup> With its short terminal half-life, FF-10101 achieves a potent, sustained inhibition of FLT3 that appears very tolerable and that can be terminated abruptly. This would allow greater flexibility in its incorporation into myelosuppressive treatment regimens. The only consistent toxicity we observed was diarrhea, which only was notable at doses above the RP2D.

Thus, while in a narrow sense FF-10101 might have a role in the treatment of FLT3-mutated AML that is resistant to quizartinib or gilteritinib, its properties of being potent but with a short half-life potentially make this more ideal drug to use in combination regimens. However, a direct head-to-head comparison of FF-10101 with either gilteritinib or quizartinib in the relapsed/refractory setting, in the form of a phase 2 randomized study, could offer more insight into the potential of this compound for the treatment of FLT3-mutated AML.

In summary, FF-10101 is a well-tolerated, oral, irreversible FLT3 inhibitor that can inhibit FLT3 in AML patients and has activity in patients previously treated with FLT3 inhibitors. This agent potentially represents the next stage in evolution of clinically important FLT3 inhibitors.

#### Authorship

M.L. designed the study, enrolled trial participants, analyzed the data, and wrote the manuscript. A.E.P. G. S., A.T.F., G.R., E.S.W., J.A., and C.S. enrolled trial participants and edited the manuscript. T.R. performed laboratory experiments. M.A., T.S., R.A.S., G.M., T.M., M.J., and K.C. contributed to study design, analyzed the data, and edited the manuscript. M.K. was the medical monitor and edited the manuscript.

#### Conflict of Interest Disclosures

M.L. received consultancy fees from Abbvie, Astellas, Amgen, Bristol-Myers-Squibb, Daiichi-Sankyo, Jazz, Pfizer, and Kite, and research support from Fujifilm. G.S. received research support from Fujifilm. A.T.F. received research support from Servier, Bristol-Myers-Squibb, and Abbvie, and consulting fees from Servier, Bristol-Myers-Squibb, Abbvie, Genentech, Amgen, Takeda, Astellas, Mablytics, EnClear, Orum, Forma, Daiichi Sankyo, Pfizer, Rigel, Autolus, Menarini, and Remix. G.R. received research support from Janssen, and consultancy fees from Abbvie, Amgen, Argenx, Astra Zeneca, Bluebird Bio, Blueprint Medicines, Bristol-Myers Squibb, Caribou Biosciences, Celgene, Daiichi Sankyo, Ellipses Pharma, GlaxoSmithKline, Janssen, Jasper Pharmaceuticals, Jazz Pharmaceuticals, Molecular Partners, Novartis, Pfizer, Roche, Syndax, and Takeda. E.S.W. received consultancy fees from Abbvie, Astellas, BMS, Daiichi Sankyo, Genentech, Gilead, GlaxoSmithKline, Janssen, Jazz, Kite, Kura, Novartis, NuProbe, Pfizer, Rigel, Sellas, and Sumitomo Pharma. A.P. received consultancy fees from Abbvie, Astellas, BerGenBio, Daiichi Sankyo, Genentech, Immunogen, Foghorn, Actinium, and Forma, and research support from Abbvie, Astellas, Daiichi Sankyo, Fujifilm, and Syndax. J.A. received consultancy fees from AbbVie, Astellas Pharma, BioSight, Bluebird Bio, Curio, Daiichi Sankyo, Gilead, Kura Oncology, Kymera, Stemline Therapeutics, and Syros and research funding from AbbVie, Agios, ALX Oncology, Amgen, Amphivena, Aprea AB, Aptose Biosciences, Astellas Pharma, BioSight, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, Cyclacel, Fujifilm, Immunogen, Kartos Therapeutics, Kura Oncology, Loxo, and Pfizer. T.R. has no conflicts of interest to declare. M.A., T.S., R.A.S., G.M., T.M., M.J., K.C., and M.K. are employees of Fujifilm, whose product (FF-10101) is described in this work. C.C.S received research funding from Fujifilm (clinical trial), Astellas (clinical trial) AbbVie, BMS/Celgene

(clinical trial), Erasca, Revolution Medicines, and Zentalis (clinical trial), and has served on a board of advisory committee for Astellas Pharma, Daichi Sankyo, and Genentech.

### Acknowledgements

This work was supported by FUJIFILM Pharmaceuticals, USA, Inc. Cambridge, MA and funded by the Newly extended TEchnology transfer Program (NexTEP), Japan Science and Technology Agency (JST), Grant Number JPMJTT15N2.

## References

1. Frohling S, Schlenk RF, Breitnick J, et al. Prognostic significance of activating FLT3 mutations in younger adults (16 to 60 years) with acute myeloid leukemia and normal cytogenetics: a study of the AML Study Group Ulm. *Blood*. 2002;100(13):4372-4380.
2. Chevallier P, Prebet T, Pigneux A, et al. Influence of NPM1 and FLT3-ITD status on outcome in relapsed/refractory AML patients receiving salvage therapy including gemtuzumab ozogamicin. *Leukemia*. 2010;24(2):467-469.
3. Papaemmanuil E, Gerstung M, Bullinger L, et al. Genomic Classification and Prognosis in Acute Myeloid Leukemia. *N Engl J Med*. 2016;374(23):2209-2221.
4. Stone RM, Mandrekar SJ, Sanford BL, et al. Midostaurin plus Chemotherapy for Acute Myeloid Leukemia with a FLT3 Mutation. *N Engl J Med*. 2017;377(5):454-464.
5. Perl AE, Martinelli G, Cortes JE, et al. Gilteritinib or Chemotherapy for Relapsed or Refractory FLT3-Mutated AML. *N Engl J Med*. 2019;381(18):1728-1740.
6. Burchardt A, Bug G, Finke J, et al. Sorafenib As Maintenance Therapy Post Allogeneic Stem Cell Transplantation for FLT3-ITD Positive AML: Results from the Randomized, Double-Blind, Placebo-Controlled Multicentre Sormain Trial. *Blood*. 2018;132:661a.
7. Cortes JE, Khaled S, Martinelli G, et al. Quizartinib versus salvage chemotherapy in relapsed or refractory FLT3-ITD acute myeloid leukaemia (QuANTUM-R): a multicentre, randomised, controlled, open-label, phase 3 trial. *Lancet Oncol*. 2019.
8. Xuan L, Wang Y, Huang F, et al. Sorafenib maintenance in patients with FLT3-ITD acute myeloid leukaemia undergoing allogeneic haematopoietic stem-cell transplantation: an open-label, multicentre, randomised phase 3 trial. *Lancet Oncol*. 2020;21(9):1201-1212.
9. Cortes J, Perl AE, Dohner H, et al. Quizartinib, an FLT3 inhibitor, as monotherapy in patients with relapsed or refractory acute myeloid leukaemia: an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2018.
10. Smith CC, Wang Q, Chin CS, et al. Validation of ITD mutations in FLT3 as a therapeutic target in human acute myeloid leukaemia. *Nature*. 2012;485(7397):260-263.
11. Sato T, Yang X, Knapper S, et al. FLT3 ligand impedes the efficacy of FLT3 inhibitors in vitro and in vivo. *Blood*. 2011;117(12):3286-3293.
12. Man CH, Fung TK, Ho C, et al. Sorafenib treatment of FLT3-ITD(+) acute myeloid leukemia: favorable initial outcome and mechanisms of subsequent nonresponsiveness associated with the emergence of a D835 mutation. *Blood*. 2012;119(22):5133-5143.
13. McMahon CM, Ferng T, Canaani J, et al. Clonal Selection with RAS Pathway Activation Mediates Secondary Clinical Resistance to Selective FLT3 Inhibition in Acute Myeloid Leukemia. *Cancer Discov*. 2019;9(8):1050-1063.
14. Cortes JE, Kantarjian H, Foran JM, et al. Phase I Study of Quizartinib Administered Daily to Patients With Relapsed or Refractory Acute Myeloid Leukemia Irrespective of FMS-Like Tyrosine Kinase 3-Internal Tandem Duplication Status. *J Clin Oncol*. 2013;31(29):3681-3687.
15. Dohner H, Weber D, Krzykalla J, et al. Midostaurin plus intensive chemotherapy for younger and older patients with AML and FLT3 internal tandem duplications. *Blood Adv*. 2022;6(18):5345-5355.

16. Rollig C, Serve H, Huttmann A, et al. Addition of sorafenib versus placebo to standard therapy in patients aged 60 years or younger with newly diagnosed acute myeloid leukaemia (SORAML): a multicentre, phase 2, randomised controlled trial. *Lancet Oncol*. 2015;16(16):1691-1699.
17. Daver N, Perl AE, Maly J, et al. Venetoclax Plus Gilteritinib for FLT3-Mutated Relapsed/Refractory Acute Myeloid Leukemia. *J Clin Oncol*. 2022;40(35):4048-4059.
18. Yamaura T, Nakatani T, Uda K, et al. A novel irreversible FLT3 inhibitor, FF-10101, shows excellent efficacy against AML cells with FLT3 mutations. *Blood*. 2018;131(4):426-438.
19. Ferng TT, Terada D, Ando M, et al. The Irreversible FLT3 Inhibitor FF-10101 Is Active Against a Diversity of FLT3 Inhibitor Resistance Mechanisms. *Mol Cancer Ther*. 2022;21(5):844-854.
20. Blume-Jensen P, Hunter T. Oncogenic kinase signalling. *Nature*. 2001;411(6835):355-365.
21. Radi ZA, Koza-Taylor PH, Bell RR, et al. Increased serum enzyme levels associated with kupffer cell reduction with no signs of hepatic or skeletal muscle injury. *Am J Pathol*. 2011;179(1):240-247.
22. Pognan F, Buono C, Couttet P, Galarneau JR, Timsit Y, Wolf A. Liver enzyme delayed clearance in rat treated by CSF1 receptor specific antagonist Sotuletinib. *Curr Res Toxicol*. 2022;3:100091.
23. Levis M, Brown P, Smith BD, et al. Plasma inhibitory activity (PIA): a pharmacodynamic assay reveals insights into the basis for cytotoxic response to FLT3 inhibitors. *Blood*. 2006;108(10):3477-3483.
24. Cheson BD, Bennett JM, Kopecky KJ, et al. Revised recommendations of the International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. *J Clin Oncol*. 2003;21(24):4642-4649.
25. Levis M, Allebach J, Tse KF, et al. A FLT3-targeted tyrosine kinase inhibitor is cytotoxic to leukemia cells in vitro and in vivo. *Blood*. 2002;99(11):3885-3891.
26. Kelly LM, Yu JC, Boulton CL, et al. CT53518, a novel selective FLT3 antagonist for the treatment of acute myelogenous leukemia (AML). *Cancer Cell*. 2002;1(5):421-432.
27. Pratz KW, Cortes J, Roboz GJ, et al. A pharmacodynamic study of the FLT3 inhibitor KW-2449 yields insight into the basis for clinical response. *Blood*. 2009;113(17):3938-3946.
28. Pratz KW, Sato T, Murphy KM, Stine A, Rajkhowa T, Levis M. FLT3-mutant allelic burden and clinical status are predictive of response to FLT3 inhibitors in AML. *Blood*. 2010;115(7):1425-1432.
29. Yin G, Li Y, Ji Y. Bayesian dose-finding in phase I/II clinical trials using toxicity and efficacy odds ratios. *Biometrics*. 2006;62(3):777-784.
30. Sexauer A, Perl A, Yang X, et al. Terminal myeloid differentiation in vivo is induced by FLT3 inhibition in FLT3/ITD AML. *Blood*. 2012;120(20):4205-4214.
31. Fathi AT, Stein EM, DiNardo CD, Levis MJ, Montesinos P, de Botton S. Differentiation syndrome with lower-intensity treatments for acute myeloid leukemia. *Am J Hematol*. 2021;96(6):735-746.
32. Smith CC, Levis MJ, Frankfurt O, et al. A phase 1/2 study of the oral FLT3 inhibitor pexidartinib in relapsed/refractory FLT3-ITD-mutant acute myeloid leukemia. *Blood Adv*. 2020;4(8):1711-1721.

33. Boal LH, Glod J, Spencer M, et al. Pediatric PK/PD Phase I Trial of Pexidartinib in Relapsed and Refractory Leukemias and Solid Tumors Including Neurofibromatosis Type I-Related Plexiform Neurofibromas. *Clin Cancer Res.* 2020;26(23):6112-6121.
34. Perl AE, Altman JK, Cortes J, et al. Selective inhibition of FLT3 by gilteritinib in relapsed or refractory acute myeloid leukaemia: a multicentre, first-in-human, open-label, phase 1-2 study. *Lancet Oncol.* 2017;18(8):1061-1075.
35. Carow CE, Levenstein M, Kaufmann SH, et al. Expression of the hematopoietic growth factor receptor FLT3 (STK-1/Flk2) in human leukemias. *Blood.* 1996;87(3):1089-1096.
36. Ozeki K, Kiyoi H, Hirose Y, et al. Biologic and clinical significance of the FLT3 transcript level in acute myeloid leukemia. *Blood.* 2004;103(5):1901-1908.
37. Ambinder AJ, Levis M. Potential targeting of FLT3 acute myeloid leukemia. *Haematologica.* 2021;106(3):671-681.
38. Serve H, Krug U, Wagner R, et al. Sorafenib in combination with intensive chemotherapy in elderly patients with acute myeloid leukemia: results from a randomized, placebo-controlled trial. *J Clin Oncol.* 2013;31(25):3110-3118.
39. Pratz KW, Rudek MA, Smith BD, et al. A Prospective Study of Peritransplant Sorafenib for Patients with FLT3-ITD Acute Myeloid Leukemia Undergoing Allogeneic Transplantation. *Biol Blood Marrow Transplant.* 2020;26(2):300-306.
40. Fischer T, Stone RM, Deangelo DJ, et al. Phase IIB trial of oral Midostaurin (PKC412), the FMS-like tyrosine kinase 3 receptor (FLT3) and multi-targeted kinase inhibitor, in patients with acute myeloid leukemia and high-risk myelodysplastic syndrome with either wild-type or mutated FLT3. *J Clin Oncol.* 2010;28(28):4339-4345.
41. Maziarz RT, Levis M, Patnaik MM, et al. Midostaurin after allogeneic stem cell transplant in patients with FLT3-internal tandem duplication-positive acute myeloid leukemia. *Bone Marrow Transplant.* 2021;56(5):1180-1189.
42. Sanga M, James J, Marini J, Gammon G, Hale C, Li J. An open-label, single-dose, phase 1 study of the absorption, metabolism and excretion of quizartinib, a highly selective and potent FLT3 tyrosine kinase inhibitor, in healthy male subjects, for the treatment of acute myeloid leukemia. *Xenobiotica.* 2017;47(10):856-869.
43. James AJ, Smith CC, Litzow M, et al. Pharmacokinetic Profile of Gilteritinib: A Novel FLT-3 Tyrosine Kinase Inhibitor. *Clin Pharmacokinet.* 2020;59(10):1273-1290.
44. Erba HP, Montesinos P, Kim HJ, et al. Quizartinib plus chemotherapy in newly diagnosed patients with FLT3-internal-tandem-duplication-positive acute myeloid leukaemia (QuANTUM-First): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet.* 2023;401(10388):1571-1583.
45. Pratz KW, Cherry M, Altman JK, et al. Gilteritinib in Combination With Induction and Consolidation Chemotherapy and as Maintenance Therapy: A Phase IB Study in Patients With Newly Diagnosed AML. *J Clin Oncol.* 2023:JCO2202721.

Table 1. Demographics and baseline characteristics of trial participants.

Characteristic	Participants (54 total)
Mean age in years (range)	60 (21-84)
Gender	
Male	27 (50%)
Female	27 (50%)
ECOG performance status	
0	7 (13%)
1	41 (76%)
2	6 (11%)
Number of prior treatment regimens	
Median	3
Range	1-7
Prior treatment with intensive therapy*	41 (75.9%)
Prior treatment with venetoclax	21 (38.9%)
Prior Allogeneic transplant	6 (11%)
ELN 2022 classification	
Favorable	0
Intermediate	27 (50%)
Adverse	13 (24%)
Insufficient data to classify	14 (26%)
FLT3 mutation status at enrollment	
FLT3 wild type	24 (44.4%)
FLT3-ITD	24 (44%)
FLT3-TKD	4 (7%)
FLT3 non-canonical	2 (4%)
Prior FLT3 inhibitor treatment	
None	24 (44%)
Midostaurin	5 (9%)
Sorafenib	4 (7%)
Gilteritinib	8 (15%)
Multiple	13 (24%)
Total with prior FLT3 inhibitor	30 (56%)

ECOG: Eastern Cooperative Oncology Group. ELN: European LeukemiaNet. ITD: Internal tandem duplication. TKD: tyrosine kinase domain. FLT3 non-canonical mutations refer to putative activating mutations of FLT3 that are neither ITD nor TKD. The two such mutations listed were A680V and V491L.

\*Intensive therapy is defined as infusional cytarabine combined with daunorubicin or idarubicin (7+3), CPX-351, high-dose cytarabine, or fludarabine-cytarabine-idarubicin (Ida-FLAG).



Table 2. Treatment-emergent adverse events (TEAEs) possibly, probably, or definitely related to study drug occurring in 20% or more of participants.

Preferred term	Grade 1-2 (%)	Grade $\geq$ 3 (%)	All grades (%)
Nausea	19 (35.2)	0 (0)	19 (35.2)
Creatine kinase increase	11 (20.4)	5 (9.3)	16 (29.6)
Febrile neutropenia		1 (1.9%)	1 (1.9%)
Diarrhea	13 (24.1)	2 (3.7)	15 (27.8)
Aspartate aminotransferase elevation	10 (18.5)	4 (7.4)	14 (25.9)

## Figure Legends

Figure 1. Trial schema. Dose escalation of FF-10101 was carried out in cohorts of three participants from 10 mg daily up to 225 mg daily. If any participant experienced a grade 3 or higher adverse event that was possibly, probably, or definitely attributed to FF-10101, three additional participants were added to the cohort. After 100 mg per day was deemed safe and tolerable, cohorts receiving twice daily dosing were opened in parallel.

Figure 2. Turnover of the FLT3 receptor. A. Molm14 and MV411 cells (each harboring a FLT3-ITD mutation) and primary AML samples (two from patients with FLT3-ITD mutations, one from a patient with wild type FLT3) were cultured in the presence or absence of 50 ug/mL cycloheximide. At designated time points, cells were harvested and analyzed for total FLT3 expression by immunoblotting. Densitometric results were analyzed by regression analysis after linear conversion to estimate a half-life ( $t_{1/2}$ ). B. Whole blood was collected at designated time points after dosing from a participant receiving 100 mg per day FF-10101 who had circulating blasts at enrollment. For each time point, half of the sample was used to directly isolate and analyze P-FLT3 from the circulating blasts while the other half was used to isolate plasma for the PIA assay using Molm14 cells.

Figure 3. Pharmacodynamics and pharmacokinetics. A. PIA results (expressed as the percentage of phosphorylated FLT3 relative to baseline) from trough (pre-dose) time points on days 8 and 15 of cycle 1 are plotted according to dose level. The solid black line indicates the mean. B.

PIA results plotted against plasma concentration of FF-10101. Whole blood collected at designated time points throughout cycle 1 of each dose level was analyzed for both FF-10101 concentration by mass spectrometry as well as to quantify the percentage of phosphorylated FLT3 relative to baseline using the PIA assay. C. Plasma concentration of FF-10101 after dosing on cycle 2 day 1 for daily dosing cohorts. D. Plasma concentration of FF-10101 after dosing on cycle 2 day 1 for twice daily dosing cohorts.

Figure 4. Waterfall plot for best change in marrow blast percentage. Shown are changes in absolute blast percentages in bone marrow aspirates at the time of best response compared with baseline. Note that 14 participants who did not have response assessment marrow biopsy because of circulating blasts are not included in this plot. Participants with no FLT3 mutation who had prior treatment with FLT3 inhibitors (black bars with stars) had FLT3 mutations detected previously, but not at the time of enrollment on this protocol.

Figure 1





Figure 1

Figure 2

A.



B.




Figure 2

Figure 3






Figure 3

Figure 4




Figure 4