

American Society of Hematology 2021 L Street NW, Suite 900, Washington, DC 20036 Phone: 202-776-0544 | Fax 202-776-0545 bloodadvances@hematology.org

## Tet2 is a tumor suppressor in the pre-leukemic phase of T-cell acute lymphoblastic leukemia

Tracking no: ADV-2023-011970R2

Stien De Coninck (Ghent University, Belgium) Juliette Roels (Ghent University, Belgium) Beatrice Lintermans (Ghent University, Belgium) Sara T'Sas (Ghent University, Belgium) Tom Taghon (Ghent University, Belgium) David Curtis (Alfred Health, Australia) Tim Pieters (Ghent University, Belgium) Steven Goossens (Cancer Research Institute Ghent (CRIG), Belgium) Pieter Van Vlierberghe (Ghent University, Belgium)

#### Abstract:

TET2-mediated DNA demethylation plays a pivotal role in regulating pre-leukemic clonal expansion in acute myeloid leukemia (AML), where TET2 mutations are also linked to AML progression. However, its function in other types of leukemias, including T-cell acute lymphoblastic leukemia (T-ALL), remains unclear. Here, we used two different T-ALL mouse models to study the possible tumor suppressor role of Tet2 in pre-leukemic T-ALL. Overexpression of Tet2 resulted in a mild but significant increase in T-ALL latency in the immature CD2-Lmo2tg T-ALL mouse model, but no effect on survival was observed in the mature Lck-Cretg/+ Ptenfl/lf T-ALL mouse model. In contrast to the pre-leukemic thymocytes from CD2-Lmo2tg mice, Lck-Cretg/+ Ptenfl/fl thymi do not display self-renewal suggesting that the anti-leukemic effect of Tet2 occurs mainly in the pre-leukemic phase of T-ALL. In conclusion, we demonstrated that the Tet2 tumor suppressor function is dependent on the differentiation stage of T-ALL and limited to the pre-leukemic phase.

Conflict of interest: No COI declared

COI notes:

Preprint server: No;

Author contributions and disclosures: Conceptualization: SDC, JR, DC, SG, PVV; Methodology: SDC, BL, STS; Formal analysis: SDC, JR; Investigation: SDC, BL, STS; Resources: TT, DC, TP, SG, PVV; Data curation: SDC; Writing original draft: SDC, JR, TP, SG, PVV; Writing - review and editing: SDC, JR, BL, STS, TT, DC TP, SGS; Visualization: SDC; Supervision: SG, PVV; Project administration: SDC, SG, PVV; Funding acquisition: SG, PVV.

#### Non-author contributions and disclosures: No;

Agreement to Share Publication-Related Data and Data Sharing Statement: For original data, please contact steven.goossens@ugent.be

Clinical trial registration information (if any):

# *Tet2* is a tumor suppressor in the preleukemic phase of T-cell acute lymphoblastic leukemia

Stien De Coninck<sup>1,2</sup>, Juliette Roels<sup>1,2</sup>, Béatrice Lintermans<sup>1,2</sup>, Sara T'Sas<sup>1,2,3</sup>, Tom Taghon<sup>4</sup>, David J. Curtis<sup>5,6</sup>, Tim Pieters<sup>1,2,3,7</sup>, Steven Goossens<sup>2,3,\*</sup>, Pieter Van Vlierberghe<sup>1,2,\*</sup>

<sup>1</sup> Lab of Normal and Malignant Hematopoiesis, Department of Biomolecular Medicine, Ghent University, 9000 Ghent, Belgium

<sup>2</sup> Cancer Research Institute Ghent (CRIG), 9000 Ghent, Belgium

<sup>3</sup> Unit for Translational Research in Oncology, Department of Diagnostic Sciences, Ghent University, 9000 Ghent, Belgium

<sup>4</sup> Taghon Laboratory, Department of Diagnostic Sciences, Ghent University, 9000 Ghent, Belgium

<sup>5</sup> Australian Centre for Blood Diseases, Central Clinical School, Monash University, Melbourne, VIC, 3004, Australia

<sup>6</sup> Department of Clinical Haematology, Alfred Hospital, Prahran, VIC, 3181, Australia
<sup>7</sup> Lab for research in oncogenesis and resistance to therapy, Department of Biomolecular Medicine, Ghent University, Ghent, Belgium

\* Authors contributed equally to this work

Running Title: *Tet2* in preleukemic T-ALL

For original data, please contact steven.goossens@ugent.be

## Acknowledgments

We thank Dr. Utpal Davé for providing us with the *CD2-Lmo2*<sup>tg</sup> mice.

SDC and JR was supported by a starting fellowship from Kom op tegen Kanker (Stand up to Cancer; the Flemish Cancer Society). The Goossens and Van Vlierberghe labs are supported by the Research Foundation Flanders (FWO-G049920N, SBO-S002322N), Ghent University start-up funds (STA/201909/007) and a Flanders interuniversity consortium grant (BOF23/IBF/042).

T-cell acute lymphoblastic leukemia (T-ALL) is a rare but aggressive leukemia affecting the T-cell lineage. During the malignant transformation of immature T-cells, a clonal expansion of progenitor cells is selected via the gradual accumulation of advantageous genetic mutations and epigenetic changes (1). Alterations in the DNA methylome are suggested as one of the potential initiating mechanisms that allow this gradual oncogenic transformation to occur (2). During DNA methylation, a methyl group is covalently attached to a cytosine in CpG-rich dinucleotide sequences by DNA methyltransferases (DNMTs) (3). It is a plastic and reversible process, where active demethylation is catalyzed by the ten-eleven translocation (TET) oxydase enzymes (4,5). Tet2 knockouts are not only predisposed to myeloid malignancies, but are also prone to develop lymphoid malignancies, including T-ALL (6,7). Although TET2 mutations in T-ALL are rare (0.7%), TET2 is transcriptionally repressed or silenced in 71% and 17% of T-ALL, respectively. This silencing is often associated with hypermethylation of the TET2 promotor (8,9). Based on these observations, restoration of the TET2 tumor suppressor function was proposed as a potential effective therapeutic strategy.

Long-lived pre-leukemic stem cells (pre-LSCs) were uncovered as an initiating event in various blood-born cancers (10). Pre-LSCs are phenotypically normal, but molecularly poised to transform. Their self-renewing capacity allows clonal expansion and subsequent acquisition of extra oncogenic driver mutations, eventually leading to the onset of a full-blown leukemia. Of interest, these pre-LSCs have a competitive advantage compared to untransformed progenitors and are more chemo/radioresistant (11-12). As a result, they are considered to reflect the population that potentially can give rise to relapse (13-15). In T-ALL, the existence of a pre-LSC population has first been described and identified in a spontaneous T-ALL mouse model overexpressing LMO2, CD2-Lmo2<sup>tg/+</sup> mice (16-18). Unlike the thymus of normal mice, which is continuously replenished by progenitors from the bone marrow, the preleukemic thymus of CD2-Lmo2<sup>tg/+</sup> mice is self-sustaining from young age and many months before leukemia development. Similar thymocyte transplantation experiments, to demonstrate presence of a pre-LSC self-renewing thymocyte population, have not been reported for other spontaneous T-ALL mouse models. To investigate presence/absence of pre-LSCs in an unrelated second T-ALL model, thymocytes from *Lck-cre<sup>tg/+</sup>Pten<sup>tl/fl</sup>* mice (19) or their Cre-negative wildtype (WT) littermate controls were transplanted into sublethal irradiated syngeneic Ly5.1 (CD45.1+) recipients. CD2-Lmo2<sup>tg/+</sup> thymocytes and WT bone marrow cells were used as a positive control. Via flow cytometric analysis, we analyzed how many of the donor versus recipient cells contributed to the repopulation of the thymus 6 weeks after transplantation. In contrast to CD45.2+ CD2-Lmo2<sup>tg/+</sup> thymocytes, none of the CD45.2+ Lck-cre<sup>tg/+</sup>Pten<sup>fl/fl</sup> thymocytes could be detected in the recipient mice, demonstrating the absence of a preleukemic phase in the Lck-cre<sup>tg/+</sup>Pten<sup>fl/fl</sup> model (Fig 1A).

Currently, detailed molecular insights are lacking in how thymocytes can gain this aberrant self-renewal and whether altered DNA methylation plays a pivotal role in this initiation event. Of interest, in our previous study (20) we already demonstrated that  $CD2-Lmo2^{tg/+}$  and  $Lck-Cre^{tg/+}Pten^{f/fl}$  murine T-ALLs have significantly different DNA methylation profiles. To investigate a potential role for *Tet2* in pre-LSCs, we started with an expression analysis for *Tet2* in thymocytes isolated from preleukemic *CD2-Lmo2<sup>tg/+</sup>* and WT littermate mice at the age of 8 and 24 weeks. *Tet2* mRNA levels

were significantly downregulated after 8 weeks and almost completely absent after 24 weeks (Fig 1B). A similar *Tet2* mRNA expression analysis was performed in preleukemic *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup>* thymocytes. Since these mice develop T-ALL faster (Supplemental Fig 1), we could only collect thymocytes of non-leukemic *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup>* mice at the age of 8 weeks. In contrast to the *CD2-Lmo2<sup>tg/+</sup>* mice, no difference in *Tet2* levels could be observed between WT and *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup>* mice (Fig 1C). Based on these findings, we hypothesized that a gradual loss of *Tet2* during the preleukemic expansion of self-renewing *CD2-Lmo2<sup>tg/+</sup>* pre-LSCs may contribute to the progression of overt leukemia in this model.

To functionally investigate the *Tet2* tumor suppressor role in T-ALL initiation, we developed a conditional Tet2 gain-of-function mouse model using our previously reported transgenic pipeline (Fig 2A) (21). We crossed this newly developed R26-*Tet2<sup>tg</sup>* with the *CD2-iCre* transgenic line (22), expressing the *Cre* recombinase from the common lymphoid progenitor stage onward. In the resulting R26-Tet2<sup>tg/tg</sup>;CD2*iCre* mice. Cre-mediated deletion of the floxed transcriptional stop cassette enabled expression of a bicistronic mRNA encoding for TET2 and an eGFP/luciferase fusion reporter in lymphoid progenitors and their progeny (Fig 2A). Next, we crossed the R26-Tet2<sup>tg</sup> with the CD2-Lmo2<sup>tg/+</sup> T-ALL model (18) and generated an aging cohort of CD2-Lmo2<sup>tg/+</sup> animals without (Cre<sup>-</sup>, CD2-Lmo2<sup>tg/+</sup>R26-Tet2<sup>tg/tg</sup>, hereafter named Tet2 overexpression (CD2-iCre<sup>tg/+</sup>CD2-Lmo2<sup>tg/+</sup>R26-Tet2<sup>tg/tg</sup>, *Lmo2*) and with hereafter named *Lmo2-Tet2*) to evaluate the effects of TET2 gain on T-ALL initiation. Mice were sacrificed when signs of hematological malignancy were observed. Lmo2-Tet2 mice showed a significant delay in T-ALL formation (median survival of 266 days) compared to Lmo2 mice (median survival of 201 days), which confirmed the role of TET2 as tumor suppressor in T-ALL (Fig 2B). Tet2 overexpression in Lmo2-Tet2 leukemia samples was confirmed by qPCR (Fig 2C) and was in line with gain of GFP (Supplemental Fig 2A) and loss of the transcriptional stop cassette (Supplemental Fig 2B). Luciferase activity (Supplemental Fig 2C-D) confirmed the conditional expression of the transgene transcript in *Lmo2-Tet2* mice. Endogenous Tet2 mRNA levels did not differ between both groups (Supplemental Fig 2E), indicating that the observed increase in T-ALL latency is exclusively caused by the Rosa26-driven Tet2 overexpression.

Next, we wondered if TET2 could act as a tumor suppressor in a mouse model that lacks pre-LSCs. For this, we crossed the R26- $Tet2^{tg/tg}$  mice with the Lck-cre<sup>tg/+</sup>Pten<sup>fl/fl</sup> T-ALL model and generated an aging cohort of mice with (Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup> R26- $Tet2^{tg/tg}$ , hereafter named *Pten-Tet2*) or without *Tet2* overexpression (Lck- $Cre^{tg/+}Pten^{fl/fl}R26$ - $Tet2^{wt/wt}$ , hereafter named *Pten)*. No significant difference in overall survival was observed (Figure 2D), although gain of *Tet2* mRNA expression was confirmed (Figure 2E) in obtained leukemia samples. Similar as for the *LMO2* aging cohort, the increased *Tet2* mRNA levels corresponded with gain of the *GFP-Luciferase* reporter gene and loss of the STOP codon (Supplemental Fig 3A-D) in the *Pten-Tet2* mice. Endogenous *Tet2* levels did not differ between both groups (Supplemental Fig 3E).

Currently, the tumor suppressor role of TET2 in T-ALL has only been studied in knockout models and rescue experiments using vitamin C or 5-azacitidine (9). Here, for the first time, we investigated the effects of gain of TET2 in T-ALL making use of genetically overexpressing mouse models. We demonstrated a tumor suppressor role of TET2 in the preleukemic phase of T-ALL, while no effects were seen on T-ALL

maintenance. The difference in the effect of *Tet2* overexpression between the *Pten* and *Lmo2* T-ALL mouse models can be correlated with our previous observations and reported differences in the DNA methylation profile of these two models (20). The observed decrease in *Tet2* mRNA levels in preleukemic *Lmo2* mice, is associated with an increase in methylation at CpG islands, while this is not seen in the *Pten* model (20). Based on these findings, we hypothesize that a critical TET2 threshold is necessary to protect preleukemic cells from obtaining secondary mutations that in the end will lead to the formation of T-ALL.

All in vivo experiments were approved by the ethical committee for animal experimentation of the Faculty of Medicine and Health Sciences of Ghent University.

### Authorship

Conceptualization: SDC, JR, DC, SG, PVV; Methodology: SDC, BL, STS; Formal analysis: SDC, JR; Investigation: SDC, BL, STS; Resources: TT, DC, TP, SG, PVV; Data curation: SDC; Writing original draft: SDC, JR, TP, SG, PVV; Writing - review and editing: SDC, JR, BL, STS, TT, DC TP, SGS; Visualization: SDC; Supervision: SG, PVV; Project administration: SDC, SG, PVV; Funding acquisition: SG, PVV.

#### The authors declare no conflict of interest.

#### References

1. Joni Van der Meulen NVR, Pieter Van Vlierberghe, Frank Speleman. The epigenetic landscape of T-cell acute lymphoblastic leukemia. The International Journal of Biochemistry & Cell Biology. 2014;53:547-557.

2. Gabriella Ficz JGG. Loss of 5-hydroxymethylcytosine in cancer: cause or consequence? Genomics. 2014;104(5):352-357.

3. Aniket V Gore BMW. DNA methylation in hematopoietic development and disease. Experimental Hematology. 2016;44(9):783-790.

4. Kasper Dindler Rasmussen KH. Role of TET enzymes in DNA methylation, development, and cancer. Genes & Development. 2016;30(7).

5. Jungeun An ARMK. TET family dioxygenases and DNA demethylation in stem cells and cancers. Experimental & Molecular Medicin. 2017;49:e323.

6. Feng Pan TSW, Zhigang Zhao, Rui Gao, Hideki Makishima, Guangbo Qu, Li Lin, Miao Yu, Janice R. Ortega, Jiapeng Wang, Aziz Nazha, Li Chen, Bing Yao, Can Liu, Shi Chen, Ophelia Weeks, Hongyu Ni, Brittany Lynn Phillips, Suming Huang, Jianlong Wang, Chuan He, Guo-Min Li, Tomas Radivoyevitch, Iannis Aifantis, Jaroslaw P. Maciejewski, Feng-Chun Yang, Peng Jin & Mingjiang Xu. Tet2 loss leads to hypermutagenicity in haematopoietic stem/progenitor cells. Nature Communications. 2017;8:15102.

7. Cyril Quivoron LC, Véronique Della Valle, Cécile K Lopez, Isabelle Plo, Orianne Wagner-Ballon, Marcio Do Cruzeiro, Francois Delhommeau, Bertrand Arnulf, Marc-

Henri Stern, Lucy Godley, Paule Opolon, Hervé Tilly, Eric Solary, Yannis Duffourd, Philippe Dessen, Hélène Merle-Beral, Florence Nguyen-Khac, Michaëla Fontenay, William Vainchenker, Christian Bastard, Thomas Mercher, Olivier A Bernard. TET2 inactivation results in pleiotropic hematopoietic abnormalities in mouse and is a recurrent event during human lymphomagenesis. Cancer Cell. 2011;20(1):25-38.

8. Ambroise Marçais LW, Julie Bruneau, Katia Hanssens, Vahid Asnafi, Philippe Gaulard, Felipe Suarez, Patrice Dubreuil, Antoine Gessain, Olivier Hermine and Claudine Pique. Adult T cell leukemia aggressivenness correlates with loss of both 5-hydroxymethylcytosine and TET2 expression. Oncotarget 2017;8(32):52256-52268.

9. Maike Bensberg OR, Aida Selimović, Cathrine Lagerwall, Mikael Benson, Mika Gustafsson, Hartmut Vogt, Antonio Lentini and Colm E. Nestor. TET2 as a tumor suppressor and therapeutic target in T-cell acute lymphoblastic leukemia. PNAS. 2021;118(34):e2110758118.

10. Aparna Vasanthakumar LAG. On the Origin of Leukemic Species. Cell Stem Cell. 2014;14(4):421-422.

11. Cedric S. Tremblay, J.S., Sung Kai Chiu, Nicholas C. Wong, Kirill Tsyganov, Sarah Ghotb, Alison N. Graham, Feng Yan, Andrew A. Guirguis, Stefan E. Sonderegger, Nicole Lee, Paul Kalitsis, John Reynolds, Stephen B. Ting, David R. Powell, Stephen M. Jane & David J. Curtis. Restricted cell cycle is essential for clonal evolution and therapeutic resistance of pre-leukemic stem cells. Nature Communications, 2018;9(1): 3535.

12. B J Shields, R.A., C Nasa, C Bogue, W S Alexander, M P McCormack, Hhex regulates Kit to promote radioresistance of self-renewing thymocytes in Lmo2-transgenic mice. Leukemia, 2015;29(4): 27-938.

13. M Ryan Corces-Zimmerman W-JH, Irving L Weissman, Bruno C Medeiros, Ravindra Majeti. Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators and persist in remission. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(7):2548-2553.

14. Carmen Vicente JC. The origin of relapse in pediatric T-cell acute lymphoblastic leukemia. Haematologica. 2015;100(11):1373-1375.

15. Yoshikane Kikushige TM. Pre-malignant lymphoid cells arise from hematopoietic stem/progenitor cells in chronic lymphocytic leukemia. International journal of hematology. 2015;102(5):528-535.

16. Matthew P McCormack LFY, Sumitha Vasudevan, Carolyn A de Graaf, Rosalind Codrington, Terence H Rabbitts, Stephen M Jane, David J Curtis. The Lmo2 Oncogene Initiates Leukemia in Mice by Inducing Thymocyte Self-Renewal. Science. 2010;327(5967):879-883.

17. Cedric S Tremblay DJC. The clonal evolution of leukemic stem cells in T-cell acute lymphoblastic leukemia. Current opinion in hematology. 2014;21(4):320-325.

18. Stephen Smith RT, Charnise Goodings, Susan Cleveland, Elizabeth Mathias, J Andrew Hardaway, Natalina Elliott, Yajun Yi, Xi Chen, James Downing, Charles Mullighan, Deborah A Swing, Lino Tessarollo, Liqi Li, Paul Love, Nancy A Jenkins, Neal G Copeland, Mary Ann Thompson, Yang Du, Utpal P Davé. LIM domain only-2 (LMO2) induces T-cell leukemia by two distinct pathways. PLoS One. 2014;9(1):e85883.

19. T J Hagenbeek HS. T-cell lymphomas in T-cell-specific Pten-deficient mice originate in the thymus. Leukemia. 2008;22(3):608-619.

20. Juliette Roels MT, Bronisława Szarzyńska, Mattias Landfors, Stien De Coninck, Lisa Demoen, Lien Provez, Anna Kuchmiy, Steven Strubbe, Lindy Reunes, Tim Pieters, Filip Matthijssens, Wouter Van Loocke, Büşra Erarslan-Uysal, Paulina Richter-Pechańska, Ken Declerck, Tim Lammens, Barbara De Moerloose, Dieter Deforce, Filip Van Nieuwerburgh, Laurence C. Cheung, Rishi S. Kotecha, Marc R. Mansour, Bart Ghesquière, Guy Van Camp, Wim Vanden Berghe, Jerzy R. Kowalczyk, Tomasz Szczepański, Utpal P. Davé, Andreas E. Kulozik, Steven Goossens, David J. Curtis, Tom Taghon, Małgorzata Dawidowska, Sofie Degerman, Pieter Van Vlierberghe. Aging of Preleukemic Thymocytes Drives CpG Island Hypermethylation in T-cell Acute Lymphoblastic Leukemia Blood Cancer Discovery. 2020;1(3):274-289.

21. Tim Pieters STS, Lisa Demoen, André Almeida, Lieven Haenebalcke, Filip Matthijssens, Kelly Lemeire, Jinke D'Hont, Frederique Van Rockeghem, Tino Hochepied, Beatrice Lintermans, Lindy Reunes, Tim Lammens, Geert Berx, Jody J. Haigh, Steven Goossens & Pieter Van Vlierberghe. Novel strategy for rapid functional in vivo validation of oncogenic drivers in haematological malignancies. Scientific Reports. 2019;9:10577.

22. Jasper de Boer AW, George Skavdis, Nicola Harker, Mark Coles, Mauro Tolaini, Trisha Norton, Keith Williams, Kathleen Roderick, Alexandre J Potocnik, Dimitris Kioussis. Transgenic mice with hematopoietic and lymphoid specific expression of Cre. European journal of immunology. 2003;33(2):314-325.

### Figure legends

#### Figure 1

- (A) %CD45.2 in thymi of sublethal irradiated CD45.1 mice injected with bone marrow of wild type mice (n=4), thymus of wild type mice (n=4), thymus of CD2-Lmo2<sup>tg/+</sup> mice (n=4) and thymus of Lck-Cre<sup>tg/+</sup> Pten<sup>fl/f</sup> mice (n=4)
- (B) Normalized *Tet2* mRNA expression in thymi of WT (n=4) and *CD2-Lmo2*<sup>tg/+</sup> (n=4) mice at 8 (p=0.0286) and 24 weeks (p=0.0286) of age
- (C)Normalized *Tet*2 mRNA expression in thymi of WT (n=4) and *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/ff</sup>* (n=4) mice at 8 weeks of age (p=0.2900)

#### Figure 2

- (A) Schematic overview of breeding scheme resulting in *CD2-iCre<sup>tg/+</sup>;R26-Tet2<sup>tg/tg</sup>* mice
- (B) Aging cohort of *CD2-Lmo2<sup>tg/+</sup>* (=*Lmo2*) (n=13, median survival 201 days) and *CD2-iCre<sup>tg/+</sup>CD2-Lmo2<sup>tg/+</sup>R26-Tet2<sup>tg/tg</sup>* (=*Lmo2-Tet2*) (n=13, median survival 266 days) mice (p=0.0226)

- (C) *Tet2* mRNA expression in thymi of *Lmo2* (n=12) and *Lmo2-Tet2* mice (n=11)(p=0.0045)
- (D) Aging cohort of *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup>R26-Tet2<sup>wt/wt</sup>* (=*Pten*) (n=11, median survival 75 days) and *Lck-Cre<sup>tg/+</sup>Pten<sup>fl/fl</sup>R26-Tet2<sup>tg/tg</sup>* (=*Pten-Tet2*) (n=11, median survial 87 days) mice (p=0.4508)
- (E) *Tet2* mRNA expression in thymi of *Pten* (n=10) and *Pten-Tet2* (n=10) mice (p=0.0138)



Downloaded from http://ashpublications.net/bloodadvances/article-pdf/doi/10.1182/bloodadvances.2023011970/2220001/bloodadvances.2023011970.pdf by guest on 27 May 2024

