

Check for updates

Acute lymphoblastic leukemia in older adults: curtain call for conventional chemotherapy?

Marlise R. Luskin

Dana-Farber Cancer Institute, Department of Medical Oncology, Division of Leukemia, Boston, MA

Unlike younger adults with acute lymphoblastic leukemia (ALL), older adults are rarely cured due to a combination of intrinsic disease resistance and treatment-related toxicities. Novel therapeutics such as inotuzumab ozogamicin, blinatumomab, venetoclax, and ABL kinase inhibitors have high activity in ALL and are well tolerated by older adults. Frontline treatment regimens for older adults using novel therapeutics with reduction or omission of conventional chemotherapy are being developed with early results demonstrating high remission rates and lower toxicity, but long-term efficacy and toxicity data are lacking. Collaboration between academic and pharmaceutical stakeholders is needed to develop clinical trials to define the optimal treatment regimens for older adults with ALL.

LEARNING OBJECTIVES

- Understand the role of novel therapeutics vs CC in initial treatment of older adults with Ph-negative ALL
- Understand the role of tyrosine kinase inhibitor treatment with or without CC or novel therapeutics in
- initial treatment of adults with Ph-positive ALL

Outcomes of older adults in the era of conventional chemotherapy

For most of history, acute lymphoblastic leukemia (ALL) has been an uncompromising, deadly illness regardless of age, comorbidities, or social circumstance. Then, in 1948, Sidney Farber announced 5 temporary remissions among 16 children with acute leukemia treated with the folic acid antagonist, aminopterin.¹ Now, after decades of effort by numerous clinicians, scientists, and cooperative groups, more than 90% of children in resourced settings are cured of ALL with chemotherapy.² Although celebrated, these pediatric ALL chemotherapy regimens are notable for length, complexity, and toxicity.

Traditionally thought of as a pediatric disease, approximately half of ALL diagnoses occur in younger (18–49 years) and older (≥50 years) adults in roughly equal porportions.³ Unfortunately, the outstanding outcomes in children have not been replicated in older cohorts. The standard approach to treating ALL in adults has been conventional chemotherapy (CC) programs adapted from pediatric schedules but with more myelosuppressive agents and a deemphasis on the noncytotoxic agents that feature prominently in pediatric regimens (corticosteroids, vincristine, and asparaginase). Results in young adults have improved due to the use of pediatric-like therapy, with more than 70% now achieving long-term survival, but older adults aged 55 to 60 or more years have consistently fared more poorly, with less than 20% cured (Table 1).⁴⁻⁹ Older patients experience more toxicity, leading to dose reductions, treatment delays, and high rates of early death and treatment-related mortality in remission.^{4-7,11,12} In addition, highrisk genetic features are more common, leading to fewer remissions and frequent relapse.^{10,16} Age-based dose modifications and prospective studies of CC designed specifically for older adults have not improved outcomes.^{4,5,12-14,17} There remains no accepted standard-of-care CC regimen for older adults with ALL.

Clinicians caring for older adults with ALL perceive limited benefit of CC. A 2019 US Medicare analysis revealed that only 53.5% of patients with ALL aged 66 or more years received any treatment within 90 days of diagnosis.¹⁸ A 2017 US Surveillance, Epidemiology, and End Results analysis reported a median overall survival (OS) of just 4 months among adults 60 years or older diagnosed with ALL since 1980, with minimal progress over 3 decades (3-year OS increasing from 10% to 16%).¹⁹ In summary, the chemotherapy regimens that cure children are not a good match for older adults.^{10,16}

Trial	Ade v	N	CR %	Farly mortality %	OS % (95% CI)	Death in CR among those achieving CR %
	Age, 5				03, % (75% 01)	
Adult trials, outcomes by ag	ge					
CALGB 9111 ⁵	All	185	85	8	3 years: 43 (36-50)	8
	≥60	35	77	17	17 (9–31)	—
	30-59	42	84	—	40 (30–51)	—
	<30	39	90	—	57 (46-68)	—
ECOG 2993/UKALL XII ^{7,11}	55-65	100	73	18	5 years: 21 (12-20)	23
	<55	1814	93	4	41 (39–43)	—
Hyper-CVAD ¹²	≥60	122	84	10	5 years: 20	34
	<60	409	92	2	48	7
Prospectively designed for	older adul	ts				
DFCI regimen ¹³	51-75	30	67	13	2 years: 52 (33-68)	—
	60-75	13	76	—	70 (42–86)	
	51-59	17	58	—	31 (10–55)	
GMALL ¹⁴	55-85	268	76	14	5 years: 23	6
	55-65	_	_	7	_	_
	65-75	_	_	14	_	_
	≥75	_	_	37	_	_
PETHEMA ALLOLD07 ¹⁵	>55	56	74	13	32 (at last follow-up, median 11.4 months)	3

Table 1. Outcomes of older adults treated with CC, select trials (see also Gökbuget¹⁰)

DFCI, Dana-Farber Cancer Institute.

Novel approaches to Philadelphia chromosome-negative ALL in older adults

CASE 1

A 78-year-old man with coronary artery disease, hypertension, and hyperlipidemia developed several weeks of fatigue and weight loss. Prior to illness, he visited the gym regularly engaging in vigorous exercise. Laboratory work revealed pancytopenia (white blood cell count, 2.9 K/µL; hemoglobin, 11.2 g/dL; platelets, 86 K/µL) with 7% circulating blasts. He was diagnosed with B-cell acute lymphoblastic leukemia (B-ALL). Immunophenotype was CD45+, CD34+, HLA-DR+, TdT+, CD19+ (variable), CD22+, CD10-, and CD20-. Karyotype was complex with no evidence of the Philadelphia chromosome. Next-generation sequencing revealed a pathogenic mutation in *TP53*. In 2019, he was referred to our academic center.

The development of novel chemotherapy agents—blinatumomab and inotuzumab ozogamicin (IO)—for relapsed B-ALL has led to a reimagining of the treatment for older adults with ALL. IO is a CD22 monoclonal antibody covalently linked to calicheamicin, a cytotoxic agent (antibody-drug conjugate). Blinatumomab is a bispecific T-cell engager targeting CD19 and CD3. Each is now approved by the US Food and Drug Administration (FDA) for relapsed and refractory (R/R) ALL based on phase 3 randomized trial data showing superiority of the novel agent to salvage cytotoxic chemotherapy in the R/R setting.^{20,21} It is hoped that using IO and blinatumomab in the first-line treatment of older adults will allow toxic CC to be de-escalated or omitted, resulting in both improved response rates and less toxicity.

IO plus CC

Investigators at MD Anderson Cancer Center (MDACC) have been testing IO plus mini-hyper-CVD (cyclophosphamide, vincristine, dexamethasone), a lower-intensity CC regimen, in older adults (≥60 years) with untreated Philadelphia chromosome-negative (Ph-) CD22+ B-ALL in a single-center phase 2 trial (NCT01371630, Table 2). In the original design, IO was administered on day 3 of the first 4 of 8 planned CC cycles followed by 36 months of POMP (6-mercaptopurine, vincristine, methotrexate, prednisone) maintenance. In 2018, MDACC investigators reported high response rates (98% overall response rate [ORR], 96% minimal residual disease-negativity [MRD-negativity]) and no early mortality in the first 52 patients treated.²² An update of the trial in 2020 (with 70 patients enrolled) was notable for continued high response rates (98% ORR, 96% MRD negativity, with no early deaths) and an encouraging 3-year complete remission (CR) duration and OS of 79% and 56%, respectively.²³

Notable toxicities associated with the IO plus mini-hyper-CVD regimen have included frequent prolonged thrombocytopenia (81%), hepatotoxicity (17% grade ≥3 hyperbilirubinemia, 9% veno-occlusive disease), and infections (41% during induction, 70% during consolidation).²³ Death in remission due to treatment-related toxicity has been the major challenge. At the most recent update, 34% (24/70) of responding patients had died in CR due to sepsis, veno-occlusive disease, or secondary myeloid malignancies. Mortality was more common in patients 70 years or older who comprised ~40% of enrolled patients. In response to an initial experience with toxicity, protocol modi-

						Regimen-related		
Reference	Regimen	Phase	Ν	Line	Age, y	deaths	Response	Survival
Kantarjian et al (2018) ²² Short et al (2020) ²³ NCT01371630	IO + mini-hyper- CVD (Blina consolidation)	2	70	First	≥60	0% early mortality 34% mortality in remission	98% ORR 88% CR (96% MRD negative)	Continuous CR 3 years: 79% Median CCR NR OS 3 years: 56% Median OS 62 months
EWALL-INO NCT03249870	IO + mild-intensity chemotherapy CC consolidation	2	_	First	≥55	_	_	_
Stelljes et al (2020) ²⁴ GMALL (INITIAL-1) NCT03460522	IO induction CC consolidation	2	36	First	≥56	0% early mortality	100% CR/CRi (78% MRD negative)	EFS 1 year: 87% (95% Cl, 70-100) OS 1 year: 87% (95% Cl, 70-100)
Jain et al (2019) ²⁵ DFCI/MDACC IST <i>NCT0331</i> 9901	Venetoclax + mini-hyper-CVD	1b/2	19	R/R (8) First (11)	All ages ≥60	0% early mortality	91% CR (100% MRD negative)	_
Advani et al (2018) ²⁶ SWOG 1318 NCT02143414	Blina induction Blina consolidation	2	29	First	≥65	0%	66% CR/CRi (92% MRD negative)	DFS 1 year: 56% (95% Cl, 58-90) OS 1 year: 65% (95% Cl, 43-80)
Alliance 041703 NCT03739814	IO induction Blina consolidation	2	_	R/R First	All ages ≥60	Results pending	Results pending	Results pending

Table 2. Novel approaches to Ph- ALL in older adults: published and ongoing trials*

*Results at most recent publication.

Blina, blinatumomab; CCR, continuous complete remission; CRi, complete remission with incomplete count recovery; NR, not reached.

fications have included IO dose fractionation and reduction (to address liver toxicity) and a drastic decrease in the number of recommended CC cycles (originally 8, decreased to 4 for patients aged 60–69 years and to 2 for patients aged \geq 70 years).

The European phase 2 EWALL (European Working Group for Adult ALL)-INO study (NCT03249870) is also exploring IO in combination with low-intensity chemotherapy for untreated patients 55 years or older with CD22+ ALL. Patients receive IO plus lowintensity chemotherapy induction for 2 cycles followed by IOfree chemotherapy consolidation and maintenance.

In contrast to the combination approach taken by MDACC and EWALL, the German Multicenter Study Group for Adult Acute Lymphoblastic Leukemia (GMALL) study group is studying a sequential approach. In the phase 2 INITIAL-1 study (NCT03460522), untreated older adults (aged >55 years) receive 3 cycles of IO monotherapy followed by CC consolidation (Table 2). A 100% CR was reported among the first 31 patients treated (78% MRD negative), with the majority (29/31) able to complete all 3 IO induction cycles with no early deaths.²⁴ Long-term efficacy and toxicity of the regimen are not yet known.

Venetoclax plus CC

The BCL2 inhibitor venetoclax, an agent with demonstrated preclinical and clinical activity in relapsed ALL, is also being studied in the frontline in combination with mini-hyper-CVD (NCT03319901, Table 2).^{27,28} A phase 1b study of venetoclax plus mini-hyper-CVD for relapsed/refractory (R/R) (n=8) and newly diagnosed patients 60 years or older (n=11) with ALL demonstrated safety (no dose-limiting toxicities (DLTs) or early mortality), with venetoclax 600mg daily declared the recommended phase 2 dose.²⁵ Notably, there was no evidence of prolonged cytopenias or liver toxicity. The regimen was particularly effective in newly diagnosed patients, with 91% (10/11) achieving an MRD-negative CR, including 6 patients with a *TP53* mutation. Because 9 of 10 responders were consolidated with allogeneic hematopoietic stem cell transplant (HSCT), there are limited data on late toxicity. The trial has now moved to phase 2 (with venetoclax 400mg) and is currently enrolling adults 60 years or older with newly diagnosed Ph– ALL. A distinct advantage of this regimen is that it can be offered to patients with both B- and T-cell ALL, in contrast to approaches that rely on B-lineage restricted antibodies and antibody–drug conjugates. Venetoclax is not approved by the FDA for ALL.

Closing the curtain on chemotherapy for Ph-ALL?

Given the efficacy of novel agents, a chemotherapy-free approach may be possible. The US National Clinical Trial Network (NCTN) trial Southwest Oncology Group (SWOG) 1318 phase 2 study of blinatumomab induction followed by blinatumomab consolidation for newly diagnosed Ph– B-ALL (aged >65 years) reported a 66% CR rate (92% MRD negative) among the first 29 patients with no early deaths (NCT02143414, Table 2).²⁶ Another phase 2 NCTN trial (Alliance 041703 NCT03739814) is investigating IO induction followed by blinatumomab consolidation (Table 2). This approach seeks to use each novel agent to maximum advantage. IO is active regardless of disease burden, making it an ideal induction agent, whereas blinatumomab is more effective and less toxic in settings of lower disease burden and thus a good agent for consolidation therapy. $^{\rm 15,29}$

Encore for chemotherapy

Blinatumomab and IO achieve high response rates in the R/R setting but with short durability.^{20,21} Although the durability of singleagent novel therapy in the upfront setting is not known, relapses driven by selective pressure would be expected. It is hypothesized that low-intensity CC integrated into induction and/or consolidation (as being pursued by EWALL-INO and GMALL INITIAL-1 studies) may continue to benefit older patients receiving novel agents by applying broader antileukemic pressure and diminishing opportunities for clonal escape. Novel combination approaches (such as being studied in A041703) may offer the best of both worlds via a multipronged attack without chemotherapy toxicities. MDACC has added blinatumomab consolidation to their IO plus mini-hyper-CVD protocol. Venetoclax plus IO will also soon be tested in R/R B-ALL due to nonoverlapping toxicities and preclinical evidence for synergy.³⁰ If shown to be safe, it may be useful as a frontline regimen. For T-cell ALL, a chemotherapy-free option does not yet exist. More research to identify therapeutic vulnerabilities and novel therapeutics for this subtype is needed.

CASE 1 (continued)

The patient was treated with venetoclax plus mini-hyper-CVD on an investigational protocol and achieved an MRD-negative CR after 1 cycle. He completed 4 cycles of venetoclax plus mini-hyper-CVD, after which he was bridged to venetoclax plus POMP maintenance due to toxicity from CC. He continued POMP maintenance for 2 years and remains in an ongoing remission. (See Figure 1 for approach to Ph-ALL in older adults.)

Novel approaches to Ph+ ALL in older adults

CASE 2

A 73-year-old woman with a history of hypertension and rheumatoid arthritis was asymptomatic but found to have an abnormal complete blood count on routine testing: white blood cells (19 K/µL) with 39% circulating blasts that are CD45+, CD34+, HLA-DR+, TdT+, CD19+, CD10+, and CD20-. Karyotype shows t(9;22) with additional abnormalities. Reverse transcription polymerase chain reaction shows p210 transcript more than 50% international scale (IS).

The Philadelphia chromosome is present in approximately half of ALLs in older adults.³¹ Historically associated with an adverse prognosis, the development of ABL kinase inhibitors has resulted in frequent cures in younger adults and in older adults has enabled prolonged remissions, negating the adverse prognosis.³²⁻³⁴ Older adults with ALL are ideal candidates for low-intensity CC and chemotherapy-free tyrosine kinase inhibitor (TKI)-based protocols.

TKI plus CC

The EWALL has studied TKI plus low-intensity CC in adults 55 years or older (NCT028889777, Table 3). Second-generation TKIs were combined with low-intensity induction and consolidation CC in the consecutive EWALL-PH01 (dasatinib) and EWALL-PH02 (nilotinib) protocols. Both protocols resulted in high CR rates (>95%) without induction mortality. Long-term survival was particularly encouraging in the EWALL-PH02, in which allogeneic HSCT was pursued in 30% (24/79) of patients (4-year OS, 47%; 61% and 39% in transplanted vs nontransplanted, respectively).^{35,36} Importantly, the Group for Research on Adult Acute Lymphoblastic Leukemia (GRAAPH)-2005 trial, which enrolled younger adults (18-59 years), demonstrated that reduced-intensity CC induction was associated with fewer induction deaths and equivalent efficacy compared with standard-intensity CC induction, questioning the benefit of intensive CC induction for any patient (regardless of age) with Ph+ ALL in the TKI era.³⁷

Curtain call on chemotherapy for Ph± ALL?

The Italian Gruppo Italiano Malattie EMatologiche dell'Adulto (GIMEMA) group has pioneered a chemotherapy-free induction approach for Ph+ ALL (Table 3). Imatinib plus corticosteroids induced universal hematologic remissions, although with limited depth (only 1 of 29 achieved a complete molecular remission) and durability (median duration of hematologic response of 8 months).³⁸ Subsequently, the more potent TKI dasatinib was combined with corticosteroids, with more patients achieving deep molecular remissions (52.1% achieved BCR-ABL level <10⁻³ by day 85).^{39,40} The original GIMEMA studies were induction-only studies, with subsequent consolidation therapy for each patient dictated by the treating physician. The US NCTN Alliance 10701 phase 2 study (NCT01256398, Table 3) tested a chemotherapy-free dasatinib plus steroid induction followed

- Careful assessment of comorbidities and performance status.
- Formal geriatric assessment if able.
- Enroll in clinical trial of novel agent therapy, wherever possible.
- If treating with conventional chemotherapy, apply dose reductions as appropriate (particularly for asparaginase).
- Ensure CNS prophylaxis is administered per regimen.
- Monitor response per protocol and include MRD assessment (flow cytometry or molecular).
- Consider referring for allogeneic HSCT if fit and high-risk genetics or persistent MRD.

Table 3. Novel approaches to Ph+ ALL in older adults: published and ongoing trials*

Reference	Regimen	Phase	N	Line	Age, y	Regimen- related deaths	Response	Survival
Rousselot et al (2016) ³⁵ EWALL PH-01 <i>NCT028889777</i>	Dasatinib 140mg daily + low- intensity CC	2	71	First	≥55	4% (3/71) induction 12% (6/71) treatment- related mortality in CR	96% CR 65% 3-log reduction in <i>BCR-ABL</i>	27% (95% Cl, 17–37) 5-year EFS 36% (95% Cl, 25–47) 5-year OS 7 received HSCT 75% of relapses T315I
Ottmann et al (2018) ³⁶ EWALL PH-02 NCT028889777	Nilotinib 400mg twice daily + low-intensity CC	2	79	First	≥55	1% (1/79) induction 11 died in CR (6 after HSCT)	94.4% CR	42% 4-year EFS 47% 4-year OS (61% transplanted, 39% nontransplanted) 24 received alloHSCT, 3 received autoHSCT
Vignetti et al (2007) ³⁸ GIMEMA LAL0101-B	Imatinib 800mg daily + prednisone (induction protocol only)	2	29	First	>60	0% induction 2 died in CR	100% CR 1/27 CMR	48% (95% Cl, 28-69) 1-year DFS 74% (95% Cl, 54-94) 1-year OS
Foà et al (2011) ³⁹ GIMEMA LAL1205	Dasatinib 70mg twice daily + prednisone (induction protocol only)	2	53	First	≥18 (22%> 60)	0% induction	100% CR 52.1% 3-log reduction in <i>BCR-ABL</i> by day 85	69% (95% 61–79) 20-month OS 51% (95% Cl, 44–59) 20-month DFS
Martinelli et al (2017) ⁴⁰ GIMEMA LAL1811 NCT01641107	Ponatinib 45mg daily + prednisone (induction proto- col only)	2	42	First	>60 or unfit	0% induction 1 death in follow-up ponatinib	95.5% CR 45% CMR at 24 weeks	87.5% (95% CI, 76.5%- 99.9%) 1-year OS
Luskin et al (2019) ⁴² DFCI IST <i>NCT03595917</i>	Dasatinib 140 + asciminib (dose escalation) (induction protocol only)	1	8	First	>50	0% induction	100% CR	Not reported
Foà et al (2020) ⁴³ GIMEMA LAL2116 NCT02744768	Dasatinib 140 + prednisone Blinatumomab consolidation	2	63	First	≥18 (median 54)	1.6% induction (1 patient) No regimen related deaths in CR	95% CR	With median follow-up 18 months 95% (95% CI, 90–100) OS 88% (95% CI, 80–97) DFS
Wieduwilt et al (2018) ⁴¹ NCTN/Alliance 10701 <i>NCT01256398</i>	Dasatinib 140 + dexametha- sone induction CC consolidation Allo- vs autoHSCT	2	64	First	≥18 (median 60)	0% induction deaths	97% CR	43% 3-year DFS (55% allo, 43% auto, 46% chemo) 55% 3-year OS (63%≤60 years, 49%> 60 years) (75% allo, 71% auto, 55% chemo) Relapse 25% allo, 43% auto, 37% chemo
NCTN/EA9181 NCT04530565	TKI + prednisone induction TKI + hyper- CVAD vs TKI + blinatumomab consolidation	3	_	First	≥18-75	Results pending	Results pending	Results pending
SWOG 1318 NCT02143414	Dasatinib + prednisone induction Dasatinib + blinatumomab consolidation	2	_	First	≥65	Results pending	Results pending	Results pending

Downloaded from http://ashpublications.net/hematology/article-pdf/2021/1/1/7/1852080/7luskin.pdf by guest on 07 May 2024

*Results at most recent publication.

CMR, complete molecular response; DFS, disease-free survival; EFS, event-free survival.

by dasatinib plus CC consolidation and then transplant (allogeneic vs autologous HSCT or chemotherapy based on availability of an human leukocyte antigen (HLA)-matched donor and fitness).⁴¹ Deep remissions were consistently obtained (97% CR, 75% ultimately achieving a complete molecular response) with 3-year OS of 55% (49% in patients aged >60 years).

Although the GRAAPH-2005 study showed that intensive CC induction was not advantageous compared with lower-intensity CC induction in the context of TKIs, the benefit of adding CC to TKI-based consolidation, particularly in patients bridging to allogeneic HSCT, is not known.

Approaches to prevent relapse in Ph± ALL

Despite improvements with TKI-based therapy, relapse remains common, with T315I-driven relapses a particular menace.^{35,39} Approaches to address this issue include use of ponatinib, which is active against T315I; dual ABL kinase inhibition with dasatinib and asciminib; and addition of novel agent consolidation with blinatumomab.

The GIMEMA group has extended its chemotherapy-free induction approach to ponatinib plus steroids in older adults (aged \geq 60 years), with encouraging early results (NCT01641107, Table 3).⁴⁰ Although toxicity appears reasonable with dose reductions (only 15/42 remained at 45mg daily at week 24), the risk of cardiovascular toxicity with ponatinib remains a concern in older patients. The planned EWALL-03 study (NCT04688983) is randomizing patients to imatinib vs ponatinib in combination with low-intensity CC and may provide insight regarding risk/benefit ratio of later-generation TKIs. A "risk-adapted" approach using early response milestones to select which patients require escalation of therapy to more potent (and more toxic) later-generation TKIs would be attractive.

Combination ABL kinase inhibition is being explored in a phase 1 study treating newly diagnosed adults 50 years or older with Ph+ ALL with dasatinib and asciminib, a novel myristoyl binding pocket inhibitor (NCT03595917, Table 3). The hypothesis is that dual ABL inhibition may induce deeper remissions and prevent T315I relapses. Thus far, the regimen appears well tolerated, with asymptomatic amylase and lipase elevations being the most notable adverse event.⁴² This study is ongoing to establish tolerability and recommended dose.

The GIMEMA D-ALBA trial added dasatinib-blinatumomab consolidation to dasatinib-steroid induction with promising

early results (NCT02744768, Table 3).⁴³ The US NCTN SWOG 1318 Cohort 2 (NCT02143414) is studying a similar approach of dasatinib-steroid induction followed by blinatumomab consolidation. A study of ponatinib-steroid induction followed by blinatumomab consolidation is also planned (NCT04722848). An open question is whether blinatumomab consolidation should be applied to all patients or used in a risk-adapted manner. Frequency of central nervous system (CNS) relapse will also need to be monitored over time in patients treated entirely with TKIs and novel agents (with only intrathecal (IT) chemotherapy administered to prevent CNS relapse).

Other active agents in ALL, including IO and venetoclax, also require evaluation as part of treatment of Ph+ ALL. The Alliance 041703 is planning to add an arm for Ph+ patients testing an IO plus ponatinib induction.

The US NCTN is currently accruing to EA9181 (NCT04530565), a phase 3 study of patients up to age 70 years who receive chemotherapy-free induction of TKI plus steroids and then are randomized to TKI (dasatinib or ponatinib) plus hypercyclophosphamide, vincristine, doxorubicin, dexamethasone (CVAD) vs TKI plus blinatumomab consolidation to compare CC vs novel agent consolidation. Eligible patients undergo transplantation when MRD negative, and all patients receive highdose methotrexate for CNS prophylaxis.

CASE 2 (continued)

The patient was treated with dasatinib and asciminib on a clinical trial. She tolerated therapy well initially but had an inadequate response after 3 cycles of therapy (CR but MRD+ by flow cytometry, BCR/ABL 6.3% IS). She required multiple dasatinib dose reductions due to recurrent pleural effusions. She was transitioned to ponatinib 30 mg daily and continues in remission (BCR-ABL p210 transcript undetectable) now 2 years from diagnosis. (See Figure 2 for approach to Ph-ALL in older adults.)

Additional issues in advancing care for older adults with ALL

The outlook for older adults diagnosed with ALL is improving rapidly. Additional issues facing the field are summarized in Figure 3.

- Careful assessment of comorbidities and performance status.
- Formal geriatric assessment if able.
- Enroll in clinical trial of novel agent therapy, wherever possible.
- I prefer a chemotherapy-free TKI-based induction using imatinib or dasatinib, or follow published regimen (Table 3).
- Ensure CNS prophylaxis is administered per regimen.
- Monitor response per protocol and include MRD assessment (BCR-ABL transcript).
- Escalate treatment (later-generation TKI or additional novel agents) if inadequate response.
- Refer for consideration of allogeneic HSCT if fit.

Downloaded from http://ashpublications.net/hematology/article-pdf/2021/1/1/1852080/7luskin.pdf by guest on 07 May 2024

- Assess and study MRD and other surrogate end points.
- Nonrestrictive inclusion criteria.
- Include quality-of-life and geriatric assessment measures.
- Design strategies to increase recruitment of participants from underrepresented groups including nonwhite and rural patients.
- Academic and pharmaceutical industry collaborations.

Figure 3. Important components for ALL clinical trials in older adults.

Curability of older adults

Although survival is poor among older adults treated with CC, a few fit older patients may tolerate and be cured by CC regimens.^{5,11,12,44,45} In contrast, there is no evidence yet that novel agent approaches are curative, although they may induce durable MRD-negative remissions. Patients responding to novel approaches may become eligible for curative-potential allogeneic HSCT, but the role of HSCT is not clear and has not been confirmed to be superior to CC for older adults with Ph- ALL in first CR in the CC era.⁴⁶ The Alliance/CALGB 10701 study has attempted to address this question for Ph+ ALL by assigning adult patients (aged 18-69 years) to reduced intensity conditioning (RIC) allogeneic HSCT vs autologous HSCT based on availability of a fully matched donor (Table 3).41 Although 3-year OS was similar between those receiving allogeneic HSCT and other consolidation approaches, patients assigned to the allogeneic approach were less likely to relapse.

Need for randomized comparison

Results of novel agent-based regimens tested in nonrandomized studies are extremely encouraging, with strong evidence for superiority compared with historical CC regimens based on high response rates and low toxicity. The next step will be to formally compare a novel and a "standard" CC approach in Ph- ALL, and thus the NCTN is designing a randomized phase 2 study of IO plus mini-hyper-CVD (the novel regimen with the most experience to date) vs age-adjusted hyper-CVAD (Alliance 042001). The objective of this trial will be to establish a novel agent-based benchmark regimen to which other novel agent regimens can be subsequently compared (Figure 4).

Surrogate end points and risk-adapted approaches

Early response end points (such as MRD negativity) should be studied to confirm validity as surrogates for survival in older adults. This could facilitate rapid completion of trials as well as development of risk-adapted approaches.

Equity

Novel agents are expensive and likely to increase disparities between resourced and underresourced settings. Advocacy for improved access is needed.

Representative enrollment and geriatric assessment

Almost every older adult has a medical problem other than ALL. Trial eligibility should be designed to allow representative enrollment. Unnecessarily restrictive inclusion criteria that exclude patients based on advanced age, mildly compromised organ function, or "clinically insignificant" concurrent malignancies should be avoided. Geriatric assessment (GA) should be routinely incorporated into ALL trials of older adults (with financial support to ensure completion). GA has been shown to be feasible in cooperative group cancer trials, including in an Alliance trial for acute myeloid leukemia.^{47,48} The phase 2 NCTN trial in development plans to incorporate GA into its study design.

B-cell Ph-Negative Older Adult Approach (US)



Figure 4. Treating Ph-negative B-cell ALL in older adults: future trial landscape.

Quality-of-life assessment

Quality-of-life end points should be given importance. Incremental improvements in efficacy may not be "worth it" to an older adult if it is associated with significant toxicity and treatment complexity. Similarly, new regimens with equivalent responses but better tolerability may warrant further development.

Cooperation to clobber the conventional approach

By far the biggest challenge in adult ALL will be to efficiently design and conduct logical, sequential studies to answer key questions and make sure patient care "keeps up" with scientific advances. In a rare disease, this is no small task. Trials will need to be developed collaboratively, opened widely with equity considered, and conducted efficiently so that questions can be iteratively asked and answered for the betterment of patients. In the United States, a NCTN working group involving representatives from Alliance, ECOG, and SWOG has formed to coordinate efforts.

Conclusion

An older adult diagnosed with ALL in 2021 has more promising treatment options than an adult diagnosed even 5 years ago. The first task is to expeditiously replace the unacceptable traditional CC approaches with novel approaches via practice changing clinical trials. Then we must work to further define the optimal treatment for all patients considering individual (age, comorbidities, and frailty) and disease (immunophenotype, genetics, response milestones) features and personal preferences.

Conflict-of-interest disclosure

Marlise R. Luskin: discloses no relevant conflict of interest.

Off-label drug use

Marlise R. Luskin: only clinical trials discussed.

Correspondence

Marlise R. Luskin, Dana-Farber Cancer Institute, Department of Medical Oncology, Division of Leukemia, 450 Brookline Ave, Dana 2056, Boston, MA 02215; e-mail: marlise_luskin@dfci.harvard.edu.

References

- Farber S, Diamond LK. Temporary remissions in acute leukemia in children produced by folic acid antagonist, 4-aminopteroyl-glutamic acid. N Engl J Med. 1948;238(23):787–793.
- Pui C-H, Yang J-J, Hunger S-P, et al. Childhood acute lymphoblastic leukemia: progress through collaboration. J Clin Oncol. 2015;33(27):2938–2948.
- Surveillance, Epidemiology, and End Results (SEER) Program. Cancer stat facts: leukemia—acute lymphocytic leukemia (ALL). 2020. Accessed 2 July 2021. https://seer.cancer.gov/statfacts/html/alyl.html.
- Larson RA, Dodge RK, Burns CP, et al. A five-drug remission induction regimen with intensive consolidation for adults with acute lymphoblastic leukemia: cancer and leukemia group B study 8811. *Blood*. 1995;85(8):2025-2037.
- Larson RA, Dodge RK, Linker CA, et al. A randomized controlled trial of filgrastim during remission induction and consolidation chemotherapy for adults with acute lymphoblastic leukemia: CALGB study 9111. Blood. 1998;92(5):1556–1564.
- Kantarjian H, Thomas D, O'Brien S, et al. Long-term follow-up results of hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone (hyper-CVAD), a dose-intensive regimen, in adult acute lymphocytic leukemia. Cancer. 2004;101(12):2788–2801.
- Rowe JM, Buck G, Burnett AK, et al; ECOG; MRC/NCRI Adult Leukemia Working Party. Induction therapy for adults with acute lymphoblastic leukemia: results of more than 1500 patients from the international ALL trial: MRC UKALL XII/ECOG E2993. Blood. 2005;106(12):3760–3767.

- DeAngelo DJ, Stevenson KE, Dahlberg SE, et al. Long-term outcome of a pediatric-inspired regimen used for adults aged 18–50 years with newly diagnosed acute lymphoblastic leukemia. *Leukemia*. 2015;29(3):526–534.
- Stock W, Luger SM, Advani AS, et al. A pediatric regimen for older adolescents and young adults with acute lymphoblastic leukemia: results of CALGB 10403. *Blood.* 2019;133(14):1548–1559.
- Gökbuget N. Treatment of older patients with acute lymphoblastic leukemia. Hematology Am Soc Hematol Educ Program. 2016;2016(1):573–579.
- Sive JI, Buck G, Fielding A, et al. Outcomes in older adults with acute lymphoblastic leukaemia (ALL): results from the international MRC UKALL XII/ECOG2993 trial. Br J Hαematol. 2012;157(4):463–471.
- O'Brien S, Thomas DA, Ravandi F, Faderl S, Pierce S, Kantarjian H. Results of the hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone regimen in elderly patients with acute lymphocytic leukemia. *Cancer.* 2008;113(8):2097–2101.
- 13. Fathi AT, DeAngelo DJ, Stevenson KE, et al. Phase 2 study of intensified chemotherapy and allogeneic hematopoietic stem cell transplantation for older patients with acute lymphoblastic leukemia. *Cancer.* 2016;122(15): 2379-2388.
- 14. Gokbuget N, Beck J, Bruggemann M. Moderate intensive chemotherapy including CNS-prophylaxis with liposomal cytarabine is feasible and effective in older patients with Ph-negative acute lymphoblastic leukemia (ALL): results of a prospective trial from the German Multicenter Study Group for Adult ALL (GMALL). *Blood*. 2012;120(21):1493.
- DeAngelo DJ, Advani AS, Marks DI, et al. Inotuzumab ozogamicin for relapsed/refractory acute lymphoblastic leukemia: outcomes by disease burden. *Blood Cancer J.* 2020;10(8):81.
- 16. Gökbuget N. How I treat older patients with ALL. *Blood.* 2013;122(8):1366-1375.
- 17. Ribera J-M, García O, Oriol A, et al; PETHEMA Group, Spanish Society of Hematology. Feasibility and results of subtype-oriented protocols in older adults and fit elderly patients with acute lymphoblastic leukemia: results of three prospective parallel trials from the PETHEMA group. *Leuk Res.* 2016;41:12–20.
- Kim C, Molony JT, Chia VM, Kota VK, Katz AJ, Li S. Patient characteristics, treatment patterns, and mortality in elderly patients newly diagnosed with ALL. *Leuk Lymphoma*. 2019;60(6):1462–1468.
- Geyer MB, Hsu M, Devlin SM, Tallman MS, Douer D, Park JH. Overall survival among older US adults with ALL remains low despite modest improvement since 1980: SEER analysis. *Blood*. 2017;129(13):1878–1881.
- Kantarjian HM, DeAngelo DJ, Stelljes M, et al. Inotuzumab ozogamicin versus standard therapy for acute lymphoblastic leukemia. N Engl J Med. 2016;375(8):740–753.
- Kantarjian H, Stein A, Gökbuget N, et al. Blinatumomab versus chemotherapy for advanced acute lymphoblastic leukemia. N Engl J Med. 2017;376(9): 836–847.
- 22. Kantarjian H, Ravandi F, Short NJ, et al. Inotuzumab ozogamicin in combination with low-intensity chemotherapy for older patients with Philadelphia chromosome-negative acute lymphoblastic leukaemia: a single-arm, phase 2 study. *Lancet Oncol.* 2018;19(2):240–248.
- 23. Short NJ, Kantarjian HM, Ravandi F, et al. Reduced-intensity chemotherapy with mini-hyper-CVD plus inotuzumab ozogamicin, with or without blinatumomab, in older adults with newly diagnosed Philadelphia chromosome-negative acute lymphoblastic leukemia: results from a phase II study. *Blood.* 2020;136(suppl 1):1014.
- 24. Stelljes M, Raffel S, Wasch R, et al. First results of an open label phase II study to evaluate the efficacy and safety of inotuzumab ozogamicin for induction therapy followed by conventional chemotherapy based consolidation and maintenance therapy in patients aged 56 years and older with acute lymphoblastic leukemia (INITIAL-1 trial). *Blood*. 2020;136(suppl 1):267.
- Jain N, Stevenson K, Winer ES, et al. A multicenter phase I study combining venetoclax with mini-hyper-CVD in older adults with untreated and relapsed/refractory acute lymphoblastic leukemia. *Blood*. 2019;134(suppl 1):3867.
- 26. Advani AS, Moseley A, O'Dwyer KM, et al. Results of SWOG 1318: a phase 2 trial of blinatumomab followed by POMP (prednisone, vincristine, methotrexate, 6-mercaptopurine) maintenance in elderly patients with newly diagnosed Philadelphia chromosome negative B-cell acute lymphoblastic leukemia. *Blood*. 2018;132(suppl 1):33.
- Chonghaile TN, Roderick JE, Glenfield C, et al. Maturation stage of T-cell acute lymphoblastic leukemia determines BCL-2 versus BCL-XL dependence and sensitivity to ABT-199. Cancer Discov. 2014;4(9):1074–1087.
- 28. Lacayo NJ, Pullarkat VA, Stock W, et al. Safety and efficacy of venetoclax in combination with navitoclax in adult and pediatric relapsed/refractory

- Gökbuget N, Dombret H, Bonifacio M, et al. Blinatumomab for minimal residual disease in adults with B-cell precursor acute lymphoblastic leukemia. *Blood.* 2018;131(14):1522–1531.
- Kirchhoff H, Karsli U, Schoenherr C, et al. Venetoclax and dexamethasone synergize with inotuzumab ozogamicin-induced DNA damage signaling in B-lineage ALL. *Blood.* 2021;137(19):2657–2661.
- 31. Chiaretti S, Vitale A, Cazzaniga G, et al. Clinico-biological features of 5202 patients with acute lymphoblastic leukemia enrolled in the Italian AIEOP and GIMEMA protocols and stratified in age cohorts. *Haematologica*. 2013;98(11):1702–1710.
- Fielding AK. Curing Ph+ ALL: assessing the relative contributions of chemotherapy, TKIs, and allogeneic stem cell transplant. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):24–29.
- Ravandi F. How I treat Philadelphia chromosome-positive acute lymphoblastic leukemia. Blood. 2019;133(2):130–136.
- 34. Ribera J-M, García O, Fernández-Abellán P, et al; PETHEMA Group. Lack of negative impact of Philadelphia chromosome in older patients with acute lymphoblastic leukaemia in the thyrosine kinase inhibitor era: comparison of two prospective parallel protocols. Br J Haematol. 2012;159(4):485–488.
- Rousselot P, Coudé MM, Gokbuget N, et al; European Working Group on Adult ALL (EWALL) group. Dasatinib and low-intensity chemotherapy in elderly patients with Philadelphia chromosome-positive ALL. *Blood*. 2016;128(6):774–782.
- 36. Ottmann OG, Pfeifer H, Cayuela JM. Nilotinib (Tasigna®) and low intensity chemotherapy for first-line treatment of elderly patients with BCR-ABL1positive acute lymphoblastic leukemia: final results of a prospective multicenter trial (EWALL-PH02). *Blood*. 2018;132(suppl 1):31.
- 37. Chalandon Y, Thomas X, Hayette S, et al; Group for Research on Adult Acute Lymphoblastic Leukemia (GRAALL). Randomized study of reducedintensity chemotherapy combined with imatinib in adults with Ph-positive acute lymphoblastic leukemia. *Blood.* 2015;125(24):3711–3719.
- 38. Vignetti M, Fazi P, Cimino G, et al. Imatinib plus steroids induces complete remissions and prolonged survival in elderly Philadelphia chromosomepositive patients with acute lymphoblastic leukemia without additional chemotherapy: results of the Gruppo Italiano Malattie Ematologiche dell'Adulto (GIMEMA) LAL0201-B protocol. *Blood*. 2007;109(9):3676–3678.
- Foà R, Vitale A, Vignetti M, et al; GIMEMAAcute Leukemia Working Party. Dasatinib as first-line treatment for adult patients with Philadelphia chromosomepositive acute lymphoblastic leukemia. *Blood.* 2011;118(25):6521–6528.

- 40. Martinelli G, Piciocchi A, Papayannidis C, et al. First report of the Gimema LAL1811 phase II prospective study of the combination of steroids with ponatinib as frontline therapy of elderly or unfit patients with Philadelphia chromosome-positive acute lymphoblastic leukemia. *Blood*. 2017;130(suppl 1):99.
- Wieduwilt M, Yin J, Wetzler M, et al. A phase II study of dasatinib and dexamethasone as primary therapy followed by transplantation for adults with newly diagnosed Ph/BCR-ABL-positive acute lymphoblastic leukemia (Ph+ ALL): final results Alliance/CALGB study 10701. *Blood.* 2018;132(suppl 1):309.
- 42. Luskin M, Murakami MA, Stevenson KE, et al. A phase I study of asciminib (ABL001) in combination with dasatinib and prednisone for untreated BCR-ABL1-positive ALL in older adults. *Blood*. 2019;134(suppl 1):3879.
- Foà R, Bassan R, Vitale A, et al; GIMEMA Investigators. Dasatinib-blinatumomab for Ph-positive acute lymphoblastic leukemia in adults. N Engl J Med. 2020;383(17):1613–1623.
- 44. Derman BA, Streck M, Wynne J, et al. Efficacy and toxicity of reduced vs. standard dose pegylated asparaginase in adults with Philadelphia chromosome-negative acute lymphoblastic leukemia. *Leuk Lymphoma*. 2020;61(3):614–622.
- 45. Martell MP, Atenafu EG, Minden MD, et al. Treatment of elderly patients with acute lymphoblastic leukaemia using a paediatric-based protocol. Br J Haematol. 2013;163(4):458–464.
- 46. Wolach O, Stevenson KE, Wadleigh M, et al. Allogeneic transplantation is not superior to chemotherapy in most patients over 40 years of age with Philadelphia-negative acute lymphoblastic leukemia in first remission. Am J Hematol. 2016;91(8):793–799.
- Hurria A, Cirrincione CT, Muss HB, et al. Implementing a geriatric assessment in cooperative group clinical cancer trials: CALGB 360401. J Clin Oncol. 2011;29(10):1290–1296.
- Klepin HD, Ritchie E, Major-Elechi B, et al. Geriatric assessment among older adults receiving intensive therapy for acute myeloid leukemia: report of CALGB 361006 (Alliance). J Geriatr Oncol. 2020;11(1):107–113.

© 2021 by The American Society of Hematology DOI 10.1182/hematology.2021000226